

## HOLOMORPHIC MAPS THAT EXTEND TO AUTOMORPHISMS OF A BALL

WALTER RUDIN<sup>1</sup>

**ABSTRACT.** It is proved, under hypotheses that may be close to minimal, that certain types of biholomorphic maps of subregions of the unit ball in  $\mathbb{C}^n$  have the extension property to which the title alludes.

Let  $B$  (or  $B_n$ , when necessary) denote the open unit ball of  $\mathbb{C}^n$ . Thus  $z = (z_1, \dots, z_n) \in B$  provided that  $|z| < 1$ , where  $|z| = \langle z, z \rangle^{1/2}$  and  $\langle z, w \rangle = \sum z_j \bar{w}_j$ . An *automorphism* of  $B$ , i.e., a member of  $\text{Aut}(B)$ , is, by definition, a holomorphic map of  $B$  onto  $B$  that is one-to-one, and whose inverse is therefore also holomorphic. The sphere that bounds  $B$  is denoted by  $S$ .

The following extension theorem will be proved.

**THEOREM.** Assume that  $n > 1$ , and that

- (a)  $\Omega_1$  and  $\Omega_2$  are connected open subsets of  $B$ ,
- (b) for  $j = 1, 2$ ,  $\Gamma_j$  is an open subset of  $S$  such that  $\Gamma_j \subset \partial\Omega_j$ ,
- (c)  $F$  is a holomorphic one-to-one map of  $\Omega_1$  onto  $\Omega_2$ , and
- (d) there is a point  $\alpha \in \Gamma_1$ , not a limit point of  $B \cap \partial\Omega_1$ , and a sequence  $\{a_i\}$  in  $\Omega_1$ , converging to  $\alpha$ , such that  $\{F(a_i)\}$  converges to a point  $\beta \in \Gamma_2$ , not a limit point of  $B \cap \partial\Omega_2$ .

Then there exists  $\Phi \in \text{Aut}(B)$  such that  $\Phi(z) = F(z)$  for all  $z \in \Omega_1$ .

The relation of this theorem to earlier results will be discussed after its proof.

The proof will use the following well-known facts.

(I) If  $F: B_k \rightarrow B_n$  is holomorphic, and  $F(0) = 0$ , then  $|F(z)| < |z|$  for all  $z \in B_k$ , and the linear operator  $F'(0)$  (the Fréchet derivative of  $F$  at 0) maps  $B_k$  into  $B_n$ .

(II) If, in addition,  $k = n$ , then the Jacobian  $JF$  of  $F$  satisfies  $|(JF)(0)| < 1$ ; equality holds only when  $F$  is a unitary operator on  $\mathbb{C}^n$ .

(III) If  $F \in \text{Aut}(B)$  and  $F(0) = 0$ , then  $F$  is unitary.

Here is a brief indication of how these are proved. For unit vectors  $u$  and  $v$  in  $\mathbb{C}^k$  and  $\mathbb{C}^n$ , respectively, the classical Schwarz lemma applies to the function  $g$  defined by

$$g(\lambda) = \langle F(\lambda u), v \rangle, \quad (\lambda \in \mathbb{C}, |\lambda| < 1). \quad (1)$$

---

Received by the editors January 8, 1980.

AMS (MOS) subject classifications (1970). Primary 32D15.

<sup>1</sup>This research was partially supported by NSF Grant MCS 78-06860, and by the William F. Vilas Trust Estate.

Thus  $|g(\lambda)| < |\lambda|$  for all eligible  $u, v$ , which leads to  $|F(z)| < |z|$ , and  $|g'(0)| < 1$ , which completes (I), since

$$g'(0) = \langle F'(0)u, v \rangle. \tag{2}$$

Since (I) implies that no eigenvalue of  $F'(0)$  exceeds 1 in absolute value, it follows that

$$|(JF)(0)| = |\det F'(0)| < 1. \tag{3}$$

If  $|(JF)(0)| = 1$ , then the linear operator  $F'(0)$  preserves volume, and maps  $B$  into  $B$ , hence is a unitary operator  $U$ . From this it follows easily (by considering iterates of  $U^{-1}F$ ) that  $F = U$ .

To prove (III), apply (II) to  $F$  as well as to  $F^{-1}$ .

The following lemma contains the essence of the proof of the theorem. To state it, we introduce the notation (for  $z \in \mathbb{C}^n$ )

$$D_z = \{\lambda z: \lambda \in \mathbb{C}, \lambda z \in B\}. \tag{4}$$

Thus, when  $z \neq 0$ ,  $D_z$  is the disc that is the intersection with  $B$  of the complex line through 0 and  $z$ .

LEMMA. Assume that

- (i)  $\Omega_1$  and  $\Omega_2$  are connected open sets in  $B$ ,
- (ii)  $0 \in \Omega_1, 0 \in \Omega_2$ .
- (iii)  $F$  is a holomorphic one-to-one map of  $\Omega_1$  onto  $\Omega_2$ , with  $F(0) = 0$ , and
- (iv) there is a nonempty open set  $V \subset \Omega_1$ , such that  $D_z \subset \Omega_1$  and  $D_{F(z)} \subset \Omega_2$  for every  $z \in V$ .

Then there is a unitary transformation  $U$  on  $\mathbb{C}^n$  such that  $F(z) = Uz$  for all  $z \in \Omega_1$ .

PROOF OF THE LEMMA. If  $z \in V$ , then  $D_z$  lies in the domain of  $F$ . Identifying  $D_z$  with  $B_1$ , we see from fact (I) (the case  $k = 1$ ), that  $|w| < |z|$ , where  $w = F(z)$ . But  $D_w$  lies in the domain of  $F^{-1}$ , and the same argument shows that  $|z| < |w|$ . Thus  $|F(z)|^2 = |z|^2$  for all  $z \in V$ . Both of these functions are real-analytic, hence they are equal in all of  $\Omega_1$ . In particular, choosing  $r > 0$  so small that  $rB \subset \Omega_1$ , we see that  $|F(z)| = |z|$  for all  $z \in rB$ . An appropriately scaled version of fact (III) shows now that  $F$  is unitary.

PROOF OF THE THEOREM. Let  $\{a_i\}$  be as in assumption (d), put  $b_i = F(a_i)$ , and choose  $u_i \in S, v_i \in S$ , so that

$$a_i = |a_i|u_i, \quad b_i = |b_i|v_i, \quad (i = 1, 2, 3, \dots). \tag{5}$$

The geometric information contained in (d) shows that there exists  $t < 1$  such that, setting

$$E_t(\xi) = \{z \in B: t < \operatorname{Re}\langle z, \xi \rangle\}, \quad (\xi \in S), \tag{6}$$

we have  $a_i \in E_t(u_i) \subset \Omega_1$ , and  $b_i \in E_t(v_i) \subset \Omega_2$  for all sufficiently large  $i$ , say  $i > i_0$ .

If  $a \in B \setminus \{0\}$ , let  $P$  denote the orthogonal projection of  $\mathbb{C}^n$  onto the one-dimensional subspace spanned by  $a$ , put  $Q = I - P$ , and define

$$\varphi_a(z) = \frac{a - Pz - (1 - |a|^2)^{1/2}Qz}{1 - \langle z, a \rangle}, \quad (z \in \bar{B}). \tag{7}$$

Then (see [4], for instance)  $\varphi_a \in \text{Aut}(B)$  and  $\varphi_a^{-1} = \varphi_a$ . Define

$$G_i = \varphi_{b_i} \circ F \circ \varphi_{a_i}, \quad (i > i_0). \tag{8}$$

Each  $G_i$  is a holomorphic one-to-one map of  $\Omega_1^i = \varphi_{a_i}(\Omega_1)$  onto  $\Omega_2^i = \varphi_{b_i}(\Omega_2)$ , and  $G_i(0) = 0$ .

If  $a = |a|\xi$ , then  $\langle Pz, \xi \rangle = \langle z, \xi \rangle \xi$ , hence

$$\langle \varphi_a(z), \xi \rangle = (|a| - \langle z, \xi \rangle) / (1 - |a|\langle z, \xi \rangle). \tag{9}$$

If  $t < |a|$ , it follows that  $\varphi_a(E_t(\xi))$  contains all  $z \in B$  with

$$\text{Re}\langle z, \xi \rangle < (|a| - t) / (1 - |a|t). \tag{10}$$

Since  $|a_i| \rightarrow 1$  and  $|b_i| \rightarrow 1$ , and since the right side of (10) tends to 1 as  $|a|$  tends to 1, there is a sequence  $\{r_i\}$ ,  $r_i < 1$ , such that  $r_i \rightarrow 1$  as  $i \rightarrow \infty$ , and such that

$$z \in B, \text{Re}\langle z, u_i \rangle < r_i \text{ implies } z \in \Omega_1^i, \tag{11}$$

$$w \in B, \text{Re}\langle z, v_i \rangle < r_i \text{ implies } w \in \Omega_2^i. \tag{12}$$

By (11),  $r_i B \subset \Omega_1^i$ , the domain of  $G_i$ . Since  $G_i(0) = 0$ , fact (II) gives  $|(JG_i)(0)| < r_i^{-n}$ . In the same way, (12) leads to  $|(JG_i^{-1})(0)| < r_i^{-n}$ , so that  $|(JG_i)(0)| > r_i^n$ . A normal family argument shows now that a subsequence of  $\{G_i\}$  converges, uniformly on compact subsets of  $B$ , to a holomorphic map of  $B$  into  $B$  that fixes 0 and whose Jacobian at 0 has absolute value 1. By fact (II), this limit map is unitary. Call it  $U$ .

Let  $V_i$  be the set of all  $p \in B$  such that

$$D_z \subset \Omega_1^i \quad \text{and} \quad D_{Uz} \subset \Omega_2^i \tag{13}$$

for all  $z$  in some neighborhood of  $p$ .

Now fix  $\epsilon$ ,  $0 < \epsilon < 1/10$ . Using (11)–(13), we see that there is an index  $i$ , fixed from now on, such that

$$|G_i(z) - Uz| < \epsilon \quad \text{whenever } |z| < 1 - \epsilon, \tag{14}$$

and such that  $V_i$  contains a ball of radius  $2\epsilon$ , whose center  $p$  satisfies  $|p| < 1 - 3\epsilon$ . To see in more detail that this can indeed be done, note that when  $r_i$  is sufficiently close to 1, there exists a large set of points  $\xi \in S$  such that  $|\langle \xi, u_i \rangle| < r_i$  and  $|\langle \xi, U^{-1}v_i \rangle| < r_i$ . For any such  $\xi$ ,  $D_\xi \subset \Omega_1^i$  and  $D_{U\xi} \subset \Omega_2^i$ , thus  $\lambda\xi \in V_i$  if  $0 < |\lambda| < 1$ .

Thus  $D_z \subset \Omega_1^i$  if  $|z - p| < 2\epsilon$ , and  $D_w \subset \Omega_2^i$  if  $|w - Up| < 2\epsilon$ . If  $|z - p| < \epsilon$ , and  $w = G_i(z)$ , it follows that  $D_w \subset \Omega_2^i$  because

$$|w - Up| < |G_i(z) - Uz| + |z - p| < 2\epsilon. \tag{15}$$

The lemma applies therefore to  $G_i$  and shows that  $G_i$  is (the restriction of) a unitary operator. Since (8) gives

$$F = \varphi_{b_i} \circ G_i \circ \varphi_{a_i}, \tag{16}$$

the theorem is proved.

REMARKS. (i) Let  $\Omega$  be a connected open subset of  $B$  such that  $\bar{\Omega}$  contains an open subset  $\Gamma$  of  $S$ . If  $F$  is a nonconstant  $C^1$ -map of  $\bar{\Omega}$  into  $\bar{B}$  that is holomorphic in  $\Omega$  and carries  $\Gamma$  into  $S$ , then  $F \in \text{Aut}(B)$ . This was proved by Pinčuk [6, p. 381],

who extended an earlier version due to Alexander [1] in which  $C^\infty$  was assumed in place of  $C^1$ .

This Alexander-Pinčuk result is a fairly direct corollary of the present theorem. If  $F \in C^1(\bar{\Omega})$  satisfies the Alexander-Pinčuk hypotheses, it is not hard to show (see Fornaess [3, p. 549] or Pinčuk [6, p. 378]) that  $JF$  vanishes at no point of  $\Gamma$ . The inverse function theorem implies then that the hypotheses of the present theorem hold.

(ii) In Alexander's proof [2] that every proper holomorphic map of  $B$  into  $B$  is in  $\text{Aut}(B)$  when  $n > 1$ , his appeal to Fefferman's theorem can be replaced by the one proved in the present paper. Consequently, there exists now a much more elementary proof of the proper mapping theorem for  $B$ .

(iii) It is quite possible that the present theorem remains true if  $B$  is replaced by strictly pseudoconvex domains with real-analytic boundaries (as Pinčuk did in the  $C^1$ -case [7]), but an entirely different proof would have to be found; Rosay [8] (strengthening a result of Wong [9]) proved that if some boundary point  $\xi$  of a bounded domain  $\Omega \subset \mathbb{C}^n$  is a point of strict pseudoconvexity, and if there exist automorphisms  $T_k$  of  $\Omega$  such that  $\lim_{k \rightarrow \infty} T_k(p) = \xi$  for some  $p \in \Omega$ , then  $\Omega$  is biholomorphically equivalent to  $B$ .

In other strictly pseudoconvex bounded domains there are thus insufficiently many automorphisms to imitate the proof that works in  $B$ .

(iv) If  $\xi \in S$  and  $\Omega = B \cap \{z: |\xi - z| < 1\}$ ; in other words, if  $\Omega = B \cap (\xi + B)$ , then the map  $z \rightarrow \xi - z$  of  $\Omega$  onto  $\Omega$  demonstrates the relevance of the assumptions concerning the location of the points  $\alpha$  and  $\beta$  in our theorem.

#### REFERENCES

1. H. Alexander, *Holomorphic mappings from the ball and polydisc*, Math. Ann. **209** (1974), 249–256.
2. ———, *Proper holomorphic mappings in  $\mathbb{C}^n$* , Indiana Univ. Math. J. **26** (1977), 137–146.
3. J. E. Fornaess, *Embedding strictly pseudoconvex domains in convex domains*, Amer. J. Math. **98** (1976), 529–569.
4. A. Nagel and W. Rudin, *Moebius-invariant function spaces on balls and spheres*, Duke Math. J. **43** (1976), 841–865.
5. S. I. Pinčuk, *On proper holomorphic mappings of strictly pseudoconvex domains*, Siberian Math. J. **15** (1974), 644–649.
6. ———, *On the analytic continuation of holomorphic mappings*, Math. USSR-Sb. **27** (1975), 375–392.
7. ———, *Analytic continuation of mappings along strictly pseudoconvex hypersurfaces*, Soviet Math. Dokl. **18** (1977), 1237–1240.
8. J. P. Rosay, *Sur une caractérisation de la boule parmi les domaines de  $\mathbb{C}^n$  par son groupe d'automorphismes*, Ann. Inst. Fourier **29** (1979), 91–97.
9. B. Wong, *Characterization of the unit ball in  $\mathbb{C}^n$  by its automorphism group*, Invent. Math. **41** (1977), 253–257.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WISCONSIN-MADISON, MADISON, WISCONSIN 53706