

Dirichlet  $L$ -Functions and Primitive Characters

Author(s): Tom M. Apostol

Source: *Proceedings of the American Mathematical Society*, Vol. 31, No. 2 (Feb., 1972), pp. 384-386

Published by: [American Mathematical Society](#)

Stable URL: <http://www.jstor.org/stable/2037537>

Accessed: 22-12-2015 19:52 UTC

---

Your use of the JSTOR archive indicates your acceptance of the Terms & Conditions of Use, available at <http://www.jstor.org/page/info/about/policies/terms.jsp>

JSTOR is a not-for-profit service that helps scholars, researchers, and students discover, use, and build upon a wide range of content in a trusted digital archive. We use information technology and tools to increase productivity and facilitate new forms of scholarship. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).



American Mathematical Society is collaborating with JSTOR to digitize, preserve and extend access to *Proceedings of the American Mathematical Society*.

<http://www.jstor.org>

## DIRICHLET $L$ -FUNCTIONS AND PRIMITIVE CHARACTERS

TOM M. APOSTOL

**ABSTRACT.** It is well known that a Dirichlet  $L$ -function  $L(s, \chi)$  has a functional equation if the character  $\chi$  is primitive. This note proves the converse result. That is, if  $L(s, \chi)$  satisfies the usual functional equation then  $\chi$  is primitive.

**1. Introduction.** For a positive integer  $k$ , let  $\chi$  be any character modulo  $k$ , let  $L(s, \chi)$  denote the  $L$ -function defined for  $R(s) > 1$  by the Dirichlet series

$$L(s, \chi) = \sum_{n=1}^{\infty} \frac{\chi(n)}{n^s},$$

let  $G(n, k)$  denote the Gauss sum

$$G(n, k) = \sum_{h=1}^{k-1} \chi(h) e^{2\pi i n h / k},$$

and let  $G(\chi) = G(1, \chi)$ . It is well known that if  $\chi$  is *primitive* then  $L(s, \chi)$  satisfies the functional equation

$$(1) \quad L(1-s, \chi) = (2\pi)^{-s} \Gamma(s) k^{s-1} \{e^{-is\pi/2} + \chi(-1)e^{is\pi/2}\} G(\chi) L(s, \bar{\chi}).$$

A recent proof is given in [1]. This paper proves the converse result.

**THEOREM 1.** *If  $\chi$  is a character modulo  $k$  and if  $L(s, \chi)$  satisfies the functional equation (1), then  $\chi$  is a primitive character modulo  $k$ .*

The proof is based on two lemmas, each of which gives a necessary and sufficient condition for a character modulo  $k$  to be primitive.

**2. Lemmas.** The first lemma is a restatement of Theorem 1 in [2].

**LEMMA 1.** *A character  $\chi$  modulo  $k$  is primitive if, and only if,*

$$G(n, \chi) = \bar{\chi}(n) G(\chi)$$

*for every integer  $n$ .*

Received by the editors April 22, 1971.

AMS 1970 subject classifications. Primary 10H10; Secondary 30A20.

Key words and phrases. Dirichlet  $L$ -functions, primitive characters, functional equations.

© American Mathematical Society 1972

The next lemma involves the function  $F(x, s)$  defined for each real  $x$  as the analytic continuation of the Dirichlet series

$$F(x, s) = \sum_{n=1}^{\infty} \frac{e^{2\pi i n x}}{n^s}, \quad R(s) > 1.$$

This function was used recently in [1] to give a new representation of  $L(s, \chi)$  for primitive characters.

LEMMA 2. For any character  $\chi$  modulo  $k$  and any complex  $s$ , let

$$L^*(s, \chi) = \sum_{h=1}^{k-1} \chi(h) F\left(\frac{h}{k}, s\right).$$

Then we have

$$(2) \quad L^*(s, \chi) = G(\chi)L(s, \bar{\chi})$$

for all  $s$  if and only if  $\chi$  is primitive.

PROOF. If  $R(s) > 1$  we have

$$(3) \quad L^*(s, \chi) = \sum_{h=1}^{k-1} \chi(h) \sum_{n=1}^{\infty} n^{-s} e^{2\pi i n h/k} = \sum_{n=1}^{\infty} G(n, \chi) n^{-s}$$

and

$$(4) \quad G(\chi)L(s, \bar{\chi}) = \sum_{n=1}^{\infty} G(\chi)\bar{\chi}(n)n^{-s}.$$

If (2) holds for all  $s$  then it also holds for  $R(s) > 1$  and the two Dirichlet series in (3) and (4) have the same coefficients. By Lemma 1 it follows that  $\chi$  is primitive.

Conversely, if  $\chi$  is primitive, Lemma 1 shows that the two functions in (3) and (4) are equal for  $R(s) > 1$  and hence they must be equal for all  $s$ .

3. **Proof of Theorem 1.** We refer to equation (39) in [1] and note that it is valid for every character  $\chi$  modulo  $k$ . This gives us the relation

$$(5) \quad L(1 - s, \chi) = f(s, \chi)L^*(s, \chi)$$

where

$$f(s, \chi) = (2\pi)^{-s} \Gamma(s) k^{s-1} \{e^{-is\pi/2} + \chi(-1)e^{is\pi/2}\}.$$

If  $L(s, \chi)$  satisfies the functional equation (1) we also have

$$(6) \quad L(1 - s, \chi) = f(s, \chi)G(\chi)L(s, \bar{\chi}).$$

From (5) and (6) we find  $G(\chi)L(s, \bar{\chi}) = L^*(s, \chi)$  for all  $s$ . Therefore, by Lemma 2,  $\chi$  is primitive.

## REFERENCES

1. Tom M. Apostol, *Dirichlet  $L$ -functions and character power sums*, J. Number Theory **2** (1970), 223–234. MR **41** #3412.
2. ———, *Euler's  $\varphi$ -function and separable Gauss sums*, Proc. Amer. Math. Soc. **24** (1970), 482–485. MR **41** #1661.

DEPARTMENT OF MATHEMATICS, CALIFORNIA INSTITUTE OF TECHNOLOGY, PASADENA,  
CALIFORNIA 91109