

ON INTEGRAL CONDITIONS FOR THE GENERAL BELTRAMI EQUATIONS

B. Bojarski, V. Gutlyanski and V. Ryazanov

February 18, 2010

Abstract

Under integral restrictions on dilatations, it is proved existence theorems for the degenerate Beltrami equations with two characteristics $\bar{\partial}f = \mu\partial f + \nu\bar{\partial}f$ and, in particular, to the Beltrami equations of the second type $\bar{\partial}f = \nu\bar{\partial}f$ that play a great role in many problems of mathematical physics and to the so-called reduced Beltrami equations $\bar{\partial}f = \lambda \operatorname{Re} \partial f$ that also have significant applications.

2000 Mathematics Subject Classification: Primary 30C65; Secondary 30C75

1 Introduction

The existence problem for the Beltrami equations with two characteristics

$$(1.1) \quad f_{\bar{z}} = \mu(z) \cdot f_z + \nu(z) \cdot \bar{f}_z$$

where $|\mu(z)| + |\nu(z)| < 1$ a.e. was solved first in the case of the bounded dilatations

$$(1.2) \quad K_{\mu,\nu}(z) := \frac{1 + |\mu(z)| + |\nu(z)|}{1 - |\mu(z)| - |\nu(z)|}$$

in [Bo₁], Theorem 5.1.

Recently in [BGR₁], the existence of homeomorphic solutions in $W_{loc}^{1,s}$ for all $s \in [1, 2)$ to the equation (1.1) was stated in the case when $K_{\mu,\nu}$ had a majorant Q in the class BMO, bounded mean oscillation by John–Nirenberg, see [JN]. Note that $L^\infty \subset \text{BMO} \subset L_{loc}^p$ for all $p \in [1, \infty)$. Our last paper [BGR₂] was devoted to the study of more general cases when $K_{\mu,\nu} \in L_{loc}^1$ and when $K_{\mu,\nu}$ had a majorant Q in the class FMO, finite mean oscillation by Ignat’ev–Ryazanov, see [IR]. Moreover, the paper [BGR₂] contained one new criterion of the Lehto type that is the base for the further development of the theory of the degenerate Beltrami equations (1.1) with integral constraints on the dilatation $K_{\mu,\nu}$ in the present paper, see the next section.

In the theory of quasiconformal mappings, it is well-known the role of the Beltrami equations of the first type

$$(1.3) \quad f_{\bar{z}} = \mu(z) \cdot f_z$$

where $f_{\bar{z}} = \bar{\partial}f = (f_x + if_y)/2$, $f_z = \partial f = (f_x - if_y)/2$, $z = x + iy$, and f_x and f_y are partial derivatives of $f = u + iv$ in the variables x and y , respectively, and $\mu : D \rightarrow \mathbb{C}$ is a measurable function with $|\mu(z)| < 1$ a.e., see e.g. [Ah₁], [Bel], [Bo₁] and [LV] where the existence problem was resolved for the uniformly elliptic case when $\|\mu\|_\infty < 1$. The existence problem for degenerate Beltrami equations (1.3) with unbounded dilatations

$$(1.4) \quad K_\mu(z) := \frac{1 + |\mu(z)|}{1 - |\mu(z)|}$$

is currently an active area of research, see e.g. [AIM], [BJ₁]-[BJ₂], [Ch], [Da], [GMSV], [IM], [Le], [MM], [MMV], [MRSY], [MS], [RSY₁]-[RSY₄], [SY], [Tu], and [Ya]. The study of such homeomorphisms was started in the frames of the theory of the so-called mean quasiconformal mappings, see e.g. [Ah₂], [Bil], [Go], [GK], [GR], [Kr], [Kru], [Kud], [Ku], [Per], [Pes], [Rya], [Str] and [UV].

On the other hand, the Beltrami equations of the second type

$$(1.5) \quad f_{\bar{z}} = \nu(z) \cdot \bar{f}_z$$

play a great role in many problems of mathematical physics, see e.g. [KK]. Hence the research of the equations (1.1) is so actual.

Recall that a function $f : D \rightarrow \mathbb{C}$ is **absolutely continuous on lines**, abbr. $f \in \mathbf{ACL}$, if, for every closed rectangle R in D whose sides are parallel to the coordinate axes, $f|_R$ is absolutely continuous on almost all line segments in R which are parallel to the sides of R . In particular, f is ACL (possibly modified on a set of Lebesgue measure zero) if it belongs to the Sobolev class $W_{loc}^{1,1}$ of locally integrable functions with locally integrable first generalized derivatives and, conversely, if $f \in \mathbf{ACL}$ has locally integrable first partial derivatives, then $f \in W_{loc}^{1,1}$, see e.g. 1.2.4 in [MP].

If $f : D \rightarrow \mathbb{C}$ is a homeomorphic ACL solution of the Beltrami equation (1.1) with $K_{\mu,\nu} \in L_{loc}^1(D)$, then $f \in W_{loc}^{1,1}(D)$, furthermore, if $K_{\mu,\nu} \in L_{loc}^p(D)$, $p \in [1, \infty]$, then $f \in W_{loc}^{1,s}(D)$ where $s = 2p/(p+1)$. Indeed, if $f \in \mathbf{ACL}$, then f has partial derivatives f_x and f_y a.e. and, for a sense-preserving ACL homeomorphism $f : D \rightarrow \mathbb{C}$, the Jacobian $J_f(z) = |f_z|^2 - |f_{\bar{z}}|^2$ is nonnegative a.e. and, moreover,

$$(1.6) \quad |\bar{\partial}f| \leq |\partial f| \leq |\partial f| + |\bar{\partial}f| \leq Q^{1/2}(z) \cdot J_f^{1/2}(z) \quad a.e.$$

Recall that if a homeomorphism $f : D \rightarrow \mathbb{C}$ has finite partial derivatives a.e., then

$$(1.7) \quad \int_B J_f(z) \, dx dy \leq |f(B)|$$

for every Borel set $B \subseteq D$, see e.g. Lemma III.3.3 in [LV]. Consequently, applying successively the Hölder inequality and the inequality (1.7) to (1.6), we get that

$$(1.8) \quad \|\partial f\|_s \leq \|K_{\mu,\nu}\|_p^{1/2} \cdot |f(C)|^{1/2}$$

where $\|\cdot\|_s$ and $\|\cdot\|_p$ denote the L^s - and L^p -norm in a compact set $C \subset D$, respectively.

In the classical case when $\|\mu\|_\infty < 1$, equivalently, when $K_\mu \in L^\infty(D)$, every ACL homeomorphic solution f of the Beltrami equation (1.3) is in the class $W_{loc}^{1,2}(D)$ with $f^{-1} \in W_{loc}^{1,2}(f(D))$. In the case $\|\mu\|_\infty = 1$ with $K_\mu \leq Q \in \text{BMO}$, again $f^{-1} \in W_{loc}^{1,2}(f(D))$ and f belongs to $W_{loc}^{1,s}(D)$ for all $1 \leq s < 2$ but already not necessarily to $W_{loc}^{1,2}(D)$. However, there is a number of degenerate Beltrami equations (1.3) for which there exist homeomorphic solutions f of the class $W_{loc}^{1,1}(D)$ with $f^{-1} \in W_{loc}^{1,2}(f(D))$.

Following [BGR₂], we call a homeomorphism $f \in W_{loc}^{1,1}(D)$ a **regular solution** of (1.1) if f satisfies (1.1) and $J_f(z) \neq 0$ a.e. Note that by [HK] $f^{-1} \in W_{loc}^{1,2}(f(D))$ for such solutions if $K_{\mu,\nu} \in L_{loc}^1(D)$.

2 Preliminaries

The following theorem was recently established in the work [BGR₂].

2.1. Theorem. *Let D be a domain in \mathbb{C} and let μ and $\nu : D \rightarrow \mathbb{C}$ be measurable functions with $|\mu(z)| + |\nu(z)| < 1$ a.e. and $K_{\mu,\nu} \in L_{loc}^1(D)$. Suppose that*

$$(2.2) \quad \int_0^{\delta(z_0)} \frac{dr}{rk_{z_0}(r)} = \infty \quad \forall z_0 \in D$$

where $\delta(z_0) < \text{dist}(z_0, \partial D)$ and $k_{z_0}(r)$ is the average of $K_{\mu,\nu}(z)$ over the circle $|z - z_0| = r$. Then the Beltrami equation (1.1) has a regular solution.

In general, in the Beltrami equation theory in the plane as well as in the theory of space mappings, the integral conditions of the Lehto type

$$(2.3) \quad \int_0^1 \frac{dr}{rq(r)} = \infty$$

are often met where the function Q is given say in the unit ball $\mathbb{B}^n = \{x \in \mathbb{R}^n : |x| < 1\}$ and $q(r)$ is the average of the function $Q(z)$ over the sphere $|x| = r$, see e.g. [AIM], [BGR₂], [Ch], [GMSV], [Le], [MRSY], [MS], [Per], [RSY₁]–[RSY₄], [ZO₁] and [ZO₂].

On the other hand, in the theory of mappings called quasiconformal in the mean, conditions of the type

$$(2.4) \quad \int_{\mathbb{B}^n} \Phi(Q(x)) dx < \infty$$

are standard for various characteristics Q of these mappings, see e.g. [Ah₂], [Bil], [Go], [GR], [Kr]–[Ku], [Per], [Rya] and [Str].

In this connection, in the paper [RSY₄] it was established interconnections between a series of integral conditions on the function Φ and between (2.3) and (2.4), cf. also [BJ₂] and [GMSV]. We give here these conditions for Φ under which (2.4) implies (2.3).

Further we use the following notion of the inverse function for monotone functions. For every non-decreasing function $\Phi : [0, \infty] \rightarrow [0, \infty]$, the **inverse function** $\Phi^{-1} : [0, \infty] \rightarrow [0, \infty]$ can be well defined by setting

$$(2.5) \quad \Phi^{-1}(\tau) = \inf_{\Phi(t) \geq \tau} t .$$

As usual, here \inf is equal to ∞ if the set of $t \in [0, \infty]$ such that $\Phi(t) \geq \tau$ is empty. Note that the function Φ^{-1} is non-decreasing, too.

2.6. Remark. It is evident immediately by the definition that

$$(2.7) \quad \Phi^{-1}(\Phi(t)) \leq t \quad \forall t \in [0, \infty]$$

with the equality in (2.7) except intervals of constancy of the function $\Phi(t)$.

Further, the integral in (2.11) is understood as the Lebesgue–Stieltjes integral and the integrals in (2.10) and (2.12)–(2.15) as the ordinary Lebesgue integrals. In (2.10) and (2.11) we complete the definition of integrals by ∞ if $\Phi(t) = \infty$, correspondingly, $H(t) = \infty$, for all $t \geq T \in [0, \infty)$.

2.8. Theorem. Let $\Phi : [0, \infty] \rightarrow [0, \infty]$ be a non-decreasing function and set

$$(2.9) \quad H(t) = \log \Phi(t) .$$

Then the equality

$$(2.10) \quad \int_{\Delta}^{\infty} H'(t) \frac{dt}{t} = \infty$$

implies the equality

$$(2.11) \quad \int_{\Delta}^{\infty} \frac{dH(t)}{t} = \infty$$

and (2.11) is equivalent to

$$(2.12) \quad \int_{\Delta}^{\infty} H(t) \frac{dt}{t^2} = \infty$$

for some $\Delta > 0$, and (2.12) is equivalent to every of the equalities:

$$(2.13) \quad \int_0^{\delta} H\left(\frac{1}{t}\right) dt = \infty$$

for some $\delta > 0$,

$$(2.14) \quad \int_{\Delta_*}^{\infty} \frac{d\eta}{H^{-1}(\eta)} = \infty$$

for some $\Delta_* > H(+0)$,

$$(2.15) \quad \int_{\delta_*}^{\infty} \frac{d\tau}{\tau\Phi^{-1}(\tau)} = \infty$$

for some $\delta_* > \Phi(+0)$.

Moreover, (2.10) is equivalent to (2.11) and hence (2.10)–(2.15) are equivalent each to other if Φ is in addition absolutely continuous. In particular, all the conditions (2.10)–(2.15) are equivalent if Φ is convex and non-decreasing.

It is necessary here to give one more explanation. From the right hand sides in the conditions (2.10)–(2.15) we have in mind $+\infty$. If $\Phi(t) = 0$ for $t \in [0, t_*]$, then $H(t) = -\infty$ for $t \in [0, t_*]$ and we complete the definition $H'(t) = 0$ for $t \in [0, t_*]$. Note, the conditions (2.11) and (2.12) exclude that t_* belongs to the interval of integrability because in the contrary case the left hand sides in (2.11) and (2.12) are either equal to $-\infty$ or indeterminate. Hence we may assume in (2.10)–(2.13) that $\Delta > t_0$ where $t_0 := \sup_{\Phi(t)=0} t$, $t_0 = 0$ if $\Phi(0) > 0$, and $\delta < 1/t_0$, correspondingly.

Finally, we give the connection of the above conditions with the condition of the Lehto type (2.2).

Recall that a function $\psi : [0, \infty] \rightarrow [0, \infty]$ is called **convex** if $\psi(\lambda t_1 + (1-\lambda)t_2) \leq \lambda\psi(t_1) + (1-\lambda)\psi(t_2)$ for all t_1 and $t_2 \in [0, \infty]$ and $\lambda \in [0, 1]$.

In what follows, \mathbb{D} denotes the unit disk in the complex plane \mathbb{C} ,

$$(2.16) \quad \mathbb{D} = \{ z \in \mathbb{C} : |z| < 1 \}.$$

2.17. Theorem. *Let $Q : \mathbb{D} \rightarrow [0, \infty]$ be a measurable function such that*

$$(2.18) \quad \int_{\mathbb{D}} \Phi(Q(z)) \, dx dy < \infty$$

where $\Phi : [0, \infty] \rightarrow [0, \infty]$ is a non-decreasing convex function such that

$$(2.19) \quad \int_{\delta_0}^{\infty} \frac{d\tau}{\tau\Phi^{-1}(\tau)} = \infty$$

for some $\delta_0 > \tau_0 := \Phi(0)$. Then

$$(2.20) \quad \int_0^1 \frac{dr}{rq(r)} = \infty$$

where $q(r)$ is the average of the function $Q(z)$ over the circle $|z| = r$.

Finally, combining Theorems 2.8 and 2.17 we obtain the following conclusion.

2.21. Corollary. *If $\Phi : [0, \infty] \rightarrow [0, \infty]$ is a non-decreasing convex function and Q satisfies the condition (2.18), then every of the conditions (2.10)–(2.15) implies (2.20).*

3 Existence theorems

Immediately on the base of Theorem 2.1 and Corollary 2.21, we obtain the next significant result.

3.1. Theorem. *Let D be a domain in \mathbb{C} and let μ and $\nu : D \rightarrow \mathbb{C}$ be measurable functions with $|\mu(z)| + |\nu(z)| < 1$ a.e. such that*

$$(3.2) \quad \int_D \Phi(K_{\mu,\nu}(z)) \, dx dy < \infty$$

where $\Phi : [0, \infty] \rightarrow [0, \infty]$ is a non-decreasing convex function. If Φ satisfies at least one of the conditions (2.10)–(2.15), then the Beltrami equation (1.1) has a regular solution.

3.3. Remark. The condition (3.2) can be also localized to neighborhoods U_{z_0} of points $z_0 \in D$ with $\Phi = \Phi_{z_0}$ under the same conditions on the functions Φ_{z_0} . If $\infty \in D$, then the condition (3.2) for $K_{\mu,\nu}(z)$ at $\infty \in D$ should be understood as the corresponding condition for $K_{\mu,\nu}(1/\bar{z})$ at 0. The latter condition can also be rewritten explicitly in terms of $K_{\mu,\nu}(z)$ itself after the inverse change of variables $z \mapsto 1/\bar{z}$ in the form

$$(3.4) \quad \int_{U_\infty} \Phi_\infty(K_{\mu,\nu}(z)) \frac{dx dy}{|z|^4} < \infty .$$

If the domain D is unbounded, then it is better to use the global condition

$$(3.5) \quad \int_D \Phi(K_{\mu,\nu}(z)) \frac{dx dy}{(1 + |z|^2)^2} < \infty$$

instead of the condition (3.2). The latter means the integration of the function $\Phi \circ K_{\mu,\nu}$ in the spherical area.

We may assume in the above theorem that the functions $\Phi_{z_0}(t)$ and $\Phi(t)$ are not convex and non-decreasing on the whole segment $[0, \infty]$ but only on a segment $[T, \infty]$ for some $T \in (1, \infty)$. Indeed, every function $\Phi : [0, \infty] \rightarrow [0, \infty]$ which is convex and non-decreasing on a segment $[T, \infty]$, $T \in (0, \infty)$, can be replaced by a non-decreasing convex function $\Phi_T : [0, \infty] \rightarrow [0, \infty]$ in the following way. We set $\Phi_T(t) \equiv 0$ for all $t \in [0, T]$, $\Phi(t) = \varphi(t)$, $t \in [T, T_*]$, and $\Phi_T \equiv \Phi(t)$, $t \in [T_*, \infty]$, where $\tau = \varphi(t)$ is the line passing through the point $(0, T)$ and touching upon the graph of the function $\tau = \Phi(t)$ at a point $(T_*, \Phi(T_*))$, $T_* \geq T$. For such a function we have by the construction that $\Phi_T(t) \leq \Phi(t)$ for all $t \in [1, \infty]$ and $\Phi_T(t) = \Phi(t)$ for all $t \geq T_*$.

The equation of the form

$$(3.6) \quad f_{\bar{z}} = \lambda(z) \operatorname{Re} f_z$$

with $|\lambda(z)| < 1$ a.e. is called the **reduced Beltrami equation**, see e.g. [AIM], [Bo₁]–[Bo₃] and [Vo]. The equation (3.6) can be rewritten as the equation (1.1) with

$$(3.7) \quad \mu(z) = \nu(z) = \frac{\lambda(z)}{2}$$

and then

$$(3.8) \quad K_{\mu,\nu}(z) = K_\lambda(z) := \frac{1 + |\lambda(z)|}{1 - |\lambda(z)|}.$$

Thus, we obtain from Theorem 3.1 the following consequence for the reduced Beltrami equations (3.6).

3.9. Theorem. *Let D be a domain in \mathbb{C} and let λ be a measurable function with $|\lambda(z)| < 1$ a.e. such that*

$$(3.10) \quad \int_D \Phi(K_\lambda(z)) \, dx dy < \infty$$

where $\Phi : [0, \infty] \rightarrow [0, \infty]$ is a non-decreasing convex function. If Φ satisfies at least one of the conditions (2.10)–(2.15), then the reduced Beltrami equation (3.6) has a regular solution.

3.11. Remark. Remarks 3.3 are valid for the reduced Beltrami equation. Moreover, the above results remain true for the case in (1.1) when

$$(3.12) \quad \nu(z) = \mu(z) e^{i\theta(z)}$$

with an arbitrary measurable function $\theta(z) : D \rightarrow \mathbb{R}$ and, in particular, for the equations of the form

$$(3.13) \quad f_{\bar{z}} = \lambda(z) \operatorname{Im} f_z$$

with a measurable coefficient $\lambda : D \rightarrow \mathbb{C}$, $|\lambda(z)| < 1$ a.e., see e.g. [Bo₁]–[Bo₃].

Next, note that Theorem 5.50 from the work [RSY₄] for the Beltrami equations of the first type (1.3) shows that the conditions (2.10)–(2.15) are not only sufficient but also necessary for the general Beltrami equations (1.1) to have regular solutions.

Finally, the same is valid for the reduced Beltrami equations (3.6) because the examples in the mentioned theorem had the form

$$f(z) = \frac{z}{|z|} \rho(|z|)$$

where $\rho(t) = e^{I(t)}$ and

$$I(t) := \int_0^t \frac{dr}{rK(r)}.$$

Indeed, setting $z = re^{i\vartheta}$ we have that

$$\frac{\partial f}{\partial r} = \frac{\partial f}{\partial z} \cdot \frac{\partial z}{\partial r} + \frac{\partial f}{\partial \bar{z}} \cdot \frac{\partial \bar{z}}{\partial r} = e^{i\vartheta} \cdot \frac{\partial f}{\partial z} + e^{-i\vartheta} \cdot \frac{\partial f}{\partial \bar{z}}$$

and

$$\frac{\partial f}{\partial \vartheta} = \frac{\partial f}{\partial z} \cdot \frac{\partial z}{\partial \vartheta} + \frac{\partial f}{\partial \bar{z}} \cdot \frac{\partial \bar{z}}{\partial \vartheta} = ire^{i\vartheta} \cdot \frac{\partial f}{\partial z} - ire^{-i\vartheta} \cdot \frac{\partial f}{\partial \bar{z}}$$

and hence

$$\frac{\partial f}{\partial z} = \frac{e^{-i\vartheta}}{2} \left(\frac{\partial f}{\partial r} + \frac{1}{ir} \cdot \frac{\partial f}{\partial \vartheta} \right) = \frac{1}{2} \left(\frac{\rho(r)}{rK(r)} + \frac{\rho(r)}{r} \right) = \frac{\rho(r)}{2r} \cdot \frac{1+K(r)}{K(r)} > 0$$

and

$$\frac{\partial f}{\partial \bar{z}} = \frac{e^{i\vartheta}}{2} \left(\frac{\partial f}{\partial r} - \frac{1}{ir} \cdot \frac{\partial f}{\partial \vartheta} \right) = \frac{e^{2i\vartheta}}{2} \left(\frac{\rho(r)}{rK(r)} - \frac{\rho(r)}{r} \right) = e^{2i\vartheta} \cdot \frac{\rho(r)}{2r} \cdot \frac{1-K(r)}{K(r)},$$

i.e.

$$\lambda(z) = e^{2i\vartheta} \cdot \frac{1-K(r)}{1+K(r)} = -\frac{z}{\bar{z}} \cdot \frac{K(|z|)-1}{K(|z|)+1}$$

and, consequently, $K_\lambda(z) = K(|z|)$.

Acknowledgements. The research of the third author was partially supported by Institute of Mathematics of PAN, Warsaw, Poland, and by Grant F25.1/055 of the State Foundation of Fundamental Investigations of Ukraine.

References

- [Ah₁] AHLFORS L.V., *Lectures on Quasiconformal Mappings*, D. Van Nostrand Company, Inc., Princeton etc., 1966.
- [Ah₂] AHLFORS L.V., *On quasiconformal mappings*, J. Analyse Math. 3 (1954), 1–58.
- [AIM] ASTALA K., IWANIEC T. AND MARTIN G.J., *Elliptic differential equations and quasiconformal mappings in the plane*, Princeton Math. Ser., v. 48, Princeton Univ. Press, Princeton, 2009.
- [Bel] BELINSKII, P. P. *General properties of quasiconformal mappings*, Nauka, Novosibirsk, 1974. (in Russian)
- [Bil] BILUTA, P. A., *Certain extremal problems for mappings which are quasiconformal in the mean*, Sibirsk. Mat. Zh. 6 (1965), 717–726.
- [BJ₁] BRAKALOVA M.A. AND JENKINS J.A., *On solutions of the Beltrami equation*, J. Anal. Math. 76 (1998), 67–92.
- [BJ₂] BRAKALOVA M.A. AND JENKINS J.A., *On solutions of the Beltrami equation. II*, Publ. de l’Inst. Math. 75(89) (2004), 3–8.
- [Bo₁] BOJARSKI B., *Generalized solutions of a system of differential equations of the first order of the elliptic type with discontinuous coefficients*, Mat. Sb. 43(85) (1957), no. 4, 451–503. (in Russian)
- [Bo₂] BOJARSKI B., *Generalized solutions of a system of differential equations of the first order of elliptic type with discontinuous coefficients*, Univ. Print. House, Jyväskylä, 2009.
- [Bo₃] BOJARSKI B., *Primary solutions of general Beltrami equations*, Ann. Acad. Sci. Fenn. Math. 32 (2007), no. 2, 549–557.

- [BGR₁] BOJARSKI B., GUTLYANSKII V. AND RYAZANOV V., *General Beltrami equations and BMO*, Ukrainian Math. Bull. 5 (2008), no. 3, 305–326.
- [BGR₂] BOJARSKI B., GUTLYANSKII V. AND RYAZANOV V., *On the Beltrami equations with two characteristics*, Complex Variables and Elliptic Equations 54 (2009), no. 10, 935–950.
- [Ch] CHEN Z.G., *$\mu(z)$ -homeomorphisms of the plane*, Michigan Math. J. 51 (2003), no. 3, 547–556.
- [Da] DAVID G., *Solutions de l'equation de Beltrami avec $\|\mu\|_\infty = 1$* , Ann. Acad. Sci. Fenn. Ser. AI. Math. AI. 13 (1988), no. 1, 25–70.
- [Go] GOL'BERG, A., *Homeomorphisms with finite mean dilatations*, Complex analysis and dynamical systems II, Contemp. Math., 382 (2005), 177–186.
- [GK] GOL'BERG, A. L. AND KUD'YAVIN, V. S., *Mean coefficients of quasiconformality of a pair of domains*, Ukrain. Mat. Zh. 43 (1991), no. 12, 1709–1712; translation in Ukrainian Math. J. 43 (1991), no. 12, 1594–1597 (1992).
- [GMSV] GUTLYANSKII V., MARTIO O., SUGAWA T. AND VUORINEN M., *On the degenerate Beltrami equation*, Trans. Amer. Math. Soc. 357 (2005), 875–900.
- [GR] GUTLYANSKII V. AND RYAZANOV, V., *Quasiconformal mappings with integral constraints on the M. A. Lavrent'ev characteristic*, Sibirsk. Mat. Zh. 31 (1990), no. 2, 21–36, 223; translation in Siberian Math. J. 31 (1990), no. 2, 205–215.
- [HK] HENCL S. AND KOSKELA P., *Regularity of the inverse of a planar Sobolev homeomorphisms*, Arch. Ration. Mech. Anal. 180 (2006), no. 1.
- [IR] IGNAT'EV A. AND RYAZANOV V., *Finite mean oscillation in the mapping theory*, Ukrainian Math. Bull. 2 (2005), no. 3, 403–424.
- [IM] IWANIEC T. AND MARTIN G., *The Beltrami equation*, Memories of AMS 191 (2008), 1–92.
- [JN] JOHN F. AND NIRENBERG L., *On functions of bounded mean oscillation*, Comm. Pure Appl. Math. 14 (1961), 415–426.
- [Kr] KRUGLIKOV V.I., *The existence and uniqueness of mappings that are quasiconformal in the mean*, p. 123–147. In the book: Metric Questions of the Theory of Functions and Mappings, Kiev, Naukova Dumka, 1973 (in Russian).
- [Kru] KRUSHKAL', S. L., *On mean quasiconformal mappings*, Dokl. Akad. Nauk SSSR 157 (1964), 517–519. (Russian)
- [KK] KRUSHKAL' S.L. AND KÜHNAU R., *Quasiconformal mappings: new methods and applications*, Nauka, Novosibirsk, 1984. (in Russian)
- [Kud] KUD'YAVIN, V. S., *Local structure of plane mappings that are quasiconformal in the mean*, Dokl. Akad. Nauk Ukrain. SSR (1991), no. 3, 10–12. (in Russian)
- [Ku] KÜHNAU R., *Über Extremalprobleme bei im Mittel quasikonformen Abbildungen*, Lecture Notes in Math. 1013 (1983), 113–124. (in German)
- [LV] LEHTO O. AND VIRTANEN K., *Quasiconformal Mappings in the Plane*, Springer, New York etc., 1973.
- [Le] LEHTO O., *Homeomorphisms with a prescribed dilatation*, Lecture Notes in Math. 118 (1968), 58–73.
- [MM] MARTIO O. AND MIKLYUKOV V., *On existence and uniqueness of the degenerate Beltrami equation*, Complex Variables Theory Appl. 49 (2004), 647–656.
- [MMV] MARTIO O., MIKLYUKOV V. AND VUORINEN M., *Some remarks on an existence problem for degenerate elliptic system*, Proc. Amer. Math. Soc. 133 (2005), 1451–1458.

- [MRSY] MARTIO O., RYAZANOV V., SREBRO U. AND YAKUBOV E., *Moduli in Modern Mapping Theory*, Springer, New York, 2009.
- [MP] MAZ'YA V.G. AND POBORCHI S.V., *Differentiable Functions on Bad Domains*, World Scientific, Singapore, 1997.
- [MS] MIKLYUKOV V.M. AND SUVOROV G.D., *On existence and uniqueness of quasiconformal mappings with unbounded characteristics*, In the book: *Investigations in the Theory of Functions of Complex Variables and its Applications*, Kiev, Inst. Math., 1972.
- [Per] PEROVICH M., *Isolated singularity of the mean quasiconformal mappings*, Lect. Notes Math. 743 (1979), 212–214.
- [Pes] PESIN I.N., *Mappings which are quasiconformal in the mean*, Dokl. Akad. Nauk SSSR 187, no. 4 (1969), 740–742. (in Russian)
- [Rya] RYAZANOV V., *On mappings that are quasiconformal in the mean*. Sibirsk. Mat. Zh. 37 (1996), no. 2, 378–388; translation in *Siberian Math. J.* 37 (1996), no. 2, 325–334.
- [RSY₁] *Ryazanov V., Srebro U. and Yakubov E.* Degenerate Beltrami equation and radial Q -homeomorphisms, Reports Dept. Math. Helsinki 369 (2003), 1–34.
- [RSY₂] *Ryazanov V., Srebro U. and Yakubov E.* On ring solutions of Beltrami equation, *J. d'Analyse Math.* 96 (2005), 117–150.
- [RSY₃] *Ryazanov V., Srebro U. and Yakubov E.* The Beltrami equation and ring homeomorphisms, *Ukrainian Math. Bull.* 4 (2007), no. 1, 79–115.
- [RSY₄] *Ryazanov V., Srebro U. and Yakubov E.* Integral conditions in the theory of the Beltrami equations, arXiv: 1001.2821v1 [math.CV] 16 Jan 2010, 1–27.
- [Sa] SAKS S., *Theory of the Integral*, Dover Publ. Inc., New York, 1964.
- [SY] SREBRO U. AND YAKUBOV E., *The Beltrami equation*, Handbook in Complex Analysis: Geometric function theory, Vol. 2, 555–597, Elsevier B. V., 2005.
- [Str] *Strugov Yu.F.* Compactness of the classes of mappings quasiconformal in the mean, Dokl. Akad. Nauk SSSR 243 (1978), no. 4, 859–861.
- [Tu] TUKIA P., *Compactness properties of μ -homeomorphisms*, Ann. Acad. Sci. Fenn. Ser. AI. Math. AI. 16 (1991), no. 1, 47–69.
- [UV] UKHLOV, A. AND VODOPYANOV, S. K., *Mappings associated with weighted Sobolev spaces*, Complex Anal. Dynam. Syst. III, Contemp. Math. 455 (2008), 369–382.
- [Vo] VOLKOVYSHKII, L. I., *Quasiconformal mappings*, L'vov Univ. Press, L'vov, 1954. (in Russian)
- [Zo₁] *Zorich V.A.* Admissible order of growth of the characteristic of quasiconformality in the Lavrent'ev theory, Dokl. Akad. Nauk SSSR 181 (1968).
- [Zo₂] *Zorich V.A.*, Isolated singularities of mappings with bounded distortion, Mat. Sb. 81 (1970), 634–638.
- [Ya] YAKUBOV E., *Solutions of Beltrami's equation with degeneration*, Dokl. Akad. Nauk SSSR 243 (1978), no. 5, 1148–1149.

Bogdan Bojarski, Institute of Mathematics of Polish Academy of Sciences,
 ul. Sniadeckich 8, P.O. Box 21, 00–956 Warsaw, POLAND
 Email: bojarski@impan.gov.pl

Vladimir Gutlyanski, Vladimir Ryazanov, Inst. Appl. Math. Mech., NASU,
74 Roze Luxemburg str., 83114, Donetsk, UKRAINE,
Email: vladimirgut@mail.ru, vlryazanov1@rambler.ru