

# Partition identities arising from involutions

Robin Chapman  
School of Mathematical Sciences  
University of Exeter  
Exeter, EX4 4QE, UK  
rjc@maths.ex.ac.uk

6 February 2002

## Abstract

We give a simple combinatorial proof of three identities of Warnaar. The proofs exploit involutions due to Franklin and Schur.

## 1 Introduction

One of the classical arguments in the combinatorial theory of partitions is Franklin's argument [1] establishing Euler's pentagonal number formula:

$$\prod_{n=1}^{\infty} (1 - q^n) = \sum_{k=-\infty}^{\infty} (-1)^k q^{k(3k+1)/2}. \quad (1)$$

This proceeds by interpreting the left side of (1) as a weighted generating function of partitions into distinct parts:

$$\prod_{n=1}^{\infty} (1 - q^n) = \sum_{\lambda \in \mathcal{D}} (-1)^{n(\lambda)} q^{|\lambda|}$$

Here  $\mathcal{D}$  denotes the set of partitions with distinct parts,  $|\lambda|$  is the number partitioned by  $\lambda$  and  $n(\lambda)$  is the number of parts in  $\lambda$ . Franklin defines an involution  $\sigma$  defined on a "large" subset  $\mathcal{D}' \subseteq \mathcal{D}$  with the property that  $(-1)^{n(\sigma(\lambda))} q^{|\sigma(\lambda)|} = -(-1)^{n(\lambda)} q^{|\lambda|}$ . Thus the sum of  $(-1)^{n(\lambda)} q^{|\lambda|}$  over  $\mathcal{D}'$  vanishes and Euler's formula (1) follows from noting that the sum of  $(-1)^{n(\lambda)} q^{|\lambda|}$  over  $\mathcal{D} - \mathcal{D}'$  is the right side of (1).

Later Schur [3] produced a proof, relying on a more complicated involution, of the Rogers-Ramanujan identities. Schur's involution later formed the basis of an explicit bijective proof due to Garsia and Milne [2] of the Rogers-Ramanujan identities.

In this paper we use Franklin's and Schur's involutions to prove bounded (polynomial rather than power series) versions of Euler's formula and the Rogers-Ramanujan identities.

## 2 Franklin's involution

We adopt the standard  $q$ -series notation: for each integer  $n \geq 0$  define  $(a)_n = \prod_{j=0}^{n-1} (1 - aq^j)$ .

Let  $\lambda = (\lambda_1, \dots, \lambda_k)$  denote a partition, that is, a finite nonincreasing sequence of positive integers,  $|\lambda| = \sum_{j=1}^k \lambda_j$ , the number partitioned by  $\lambda$ , and  $n(\lambda) = k$ , the number of parts in  $\lambda$ . Let  $\mathcal{D}$  denote the set of partitions having distinct parts, that is the set of  $\lambda$  with  $\lambda_1 > \lambda_2 > \dots > \lambda_k$ . For nonempty  $\lambda \in \mathcal{D}$  let  $t(\lambda)$  denote the smallest part of  $\lambda$  and  $s(\lambda)$  be the "slope" of  $\lambda$ , that is, the largest integer  $s$  such that  $\lambda_s = \lambda_1 - s + 1 > 0$ .

For  $j \in \mathbf{Z}$  we define a partition  $\pi_{(j)} \in \mathcal{D}$  as follows:  $\pi_{(0)}$  is the empty partition, for  $j > 0$ ,  $\pi_{(j)} = (2j, 2j - 1, \dots, j + 1)$  and  $\pi_{(-j)} = (2j - 1, 2j - 2, \dots, j)$ . Then  $|\pi_{(j)}| = j(3j + 1)/2$  and  $n(\pi_{(j)}) = |j|$ .

Following Franklin [1] we define an involution  $\sigma$  on the set  $\mathcal{D}' = \mathcal{D} - \{\pi_{(j)} : j \in \mathbf{Z}\}$  as follows:

- if  $t(\lambda) \leq s(\lambda)$  remove the smallest part of  $\lambda$  and add 1 to each of the  $t(\lambda)$  largest parts to yield  $\sigma(\lambda)$ ;
- if  $t(\lambda) > s(\lambda)$  subtract 1 from each of the  $s(\lambda)$  largest parts of  $\lambda$  and create a new smallest part equal to  $s(\lambda)$  to yield  $\sigma(\lambda)$ .

Then  $\sigma$  is an involution on  $\mathcal{D}'$  and  $(-1)^{n(\sigma(\lambda))} q^{|\sigma(\lambda)|} = -(-1)^{n(\lambda)} q^{|\lambda|}$ .

**Theorem 1** *The following identity holds for each integer  $m \geq 0$ :*

$$\sum_{s=0}^{\lfloor m/2 \rfloor} (-1)^s q^{s(2m-s+3)/2} (q^{s+1})_{m-2s} = \sum_{j=\lfloor -m/2 \rfloor}^{\lfloor m/2 \rfloor} (-1)^j q^{j(3j+1)/2}.$$

**Proof** Let  $\mathcal{D}_m$  consist of the partitions in  $\mathcal{D}$  with parts of size at most  $m$ . Then  $\mathcal{D}_m \cap \mathcal{D}'$  is not invariant under  $\sigma$ . Suppose that  $\lambda \in \mathcal{D}_m \cap \mathcal{D}'$  but  $\sigma(\lambda) \notin \mathcal{D}_m \cap \mathcal{D}'$ . In this case  $\lambda_1 = m$  and  $t(\lambda) \leq s(\lambda)$ . Let  $s = s(\lambda)$ . Then  $\lambda$  contains a part  $m - s + 1$  and so  $m - s + 1 \geq s$ . Were equality to hold, then

$\lambda$  would equal  $\pi_{(-s)} \notin \mathcal{D}'$ . Hence  $s \leq m/2$ . Then  $\sigma(\lambda) \in \mathcal{D}_{m,s}$ , for  $s = s(\lambda)$ , where  $\mathcal{D}_{m,s}$  is the set of partitions  $\lambda \in \mathcal{D}$  with largest part  $m+1$ , slope  $s$  and smallest part  $> s$ . Conversely if  $\mu \in \mathcal{D}_{m,s} \cap \mathcal{D}'$ , for some  $s$ , then  $\sigma(\mu) \in \mathcal{D}_m$ . The set  $(\mathcal{D}_m \cup \bigcup_{s=1}^{\lfloor m/2 \rfloor} \mathcal{D}_{m,s}) \cap \mathcal{D}'$  is invariant under  $\sigma$ . It follows that

$$\sum_{\lambda \in \mathcal{D}_m \cup \bigcup_{s=1}^{\lfloor m/2 \rfloor} \mathcal{D}_{m,s}} (-1)^{n(\lambda)} q^{|\lambda|} = \sum_{j: \pi_{(j)} \in \mathcal{D}_m \cup \bigcup_{s=1}^{\lfloor m/2 \rfloor} \mathcal{D}_{m,s}} (-1)^j q^{j(3j+1)/2}. \quad (2)$$

We now examine both sides of (2). The set  $\mathcal{D}_m$  consists of all partitions in  $\mathcal{D}$  with parts from  $\{1, 2, \dots, m\}$ . Hence

$$\sum_{\lambda \in \mathcal{D}_m} (-1)^{n(\lambda)} q^{|\lambda|} = \prod_{j=1}^m (1 - q^j) = (q)_m.$$

The partitions in  $\mathcal{D}_{m,s}$  must contain parts  $m+1, m, m-1, \dots, m+2-s$  and also a subset of  $\{s+1, \dots, m-s\}$ . We have

$$\sum_{\lambda \in \mathcal{D}_{m,s}} (-1)^{n(\lambda)} q^{|\lambda|} = \prod_{j=m+2-s}^{m+1} (-q^j) \times \prod_{i=s+1}^{m-s} (1 - q^i) = (-1)^s q^{s(2m+3-s)/2} (q^{s+1})_{m-2s}.$$

Thus

$$\begin{aligned} \sum_{\lambda \in \mathcal{D}_m \cup \bigcup_{s=1}^{\lfloor m/2 \rfloor} \mathcal{D}_{m,s}} (-1)^{n(\lambda)} q^{|\lambda|} &= (q)_m + \sum_{s=1}^{\lfloor m/2 \rfloor} (-1)^s q^{s(2m+3-s)/2} (q^{s+1})_{m-2s} \\ &= \sum_{s=0}^{\lfloor m/2 \rfloor} (-1)^s q^{s(2m+3-s)/2} (q^{s+1})_{m-2s}. \end{aligned}$$

The partition  $\pi_{(j)}$  lies in  $\mathcal{D}_m$  if and only if  $0 \leq j \leq m/2$  or  $0 \geq j \geq (m-1)/2$ , that is if and only if  $\lfloor -m/2 \rfloor \leq j \leq \lfloor m/2 \rfloor$ . If  $j > 0$  and  $\pi_{(j)} \in \mathcal{D}_{m,s}$ , then  $m+1 = 2j$  and  $s = j$  so that  $2s > m$ . If  $j > 0$  and  $\pi_{(-j)} \in \mathcal{D}_{m,s}$ , then  $m+1 = 2j-1$  and  $s = j$  so again  $2s > m$ . Hence

$$\sum_{j: \pi_{(j)} \in \mathcal{D}_m \cup \bigcup_{s=1}^{\lfloor m/2 \rfloor} \mathcal{D}_{m,s}} (-1)^j q^{j(3j+1)/2} = \sum_{j=\lfloor -m/2 \rfloor}^{\lfloor m/2 \rfloor} (-1)^j q^{j(3j+1)/2}.$$

Equating both sides of (2) gives

$$\sum_{s=0}^{\lfloor m/2 \rfloor} (-1)^s q^{s(2m+3-s)/2} (q^{s+1})_{m-2s} = \sum_{j=\lfloor -m/2 \rfloor}^{\lfloor m/2 \rfloor} (-1)^j q^{j(3j+1)/2}$$

as required.  $\square$

### 3 Schur's involution

Schur [3] produced a proof of the Rogers-Ramanujan identities using an involutive argument akin to Franklin's proof of Euler's formula. Let  $\mathcal{R}$  denote the set of partitions in  $\mathcal{D}$  having parts differing by at least 2. The first Rogers-Ramanujan identity states that

$$\sum_{\mu \in \mathcal{R}} q^{|\mu|} = \prod_{n=1}^{\infty} \frac{1}{(1 - q^{5n-4})(1 - q^{5n-1})}.$$

Using Jacobi's triple product we see that this is equivalent to

$$\sum_{\mu \in \mathcal{R}} q^{|\mu|} = \frac{1}{(q)_{\infty}} \prod_{n=1}^{\infty} (1 - q^{5n-3})(1 - q^{5n-2})(1 - q^{5n}) = \frac{1}{(q)_{\infty}} \sum_{k=-\infty}^{\infty} (-1)^k q^{k(5k+1)/2}$$

and so to

$$\sum_{k=-\infty}^{\infty} (-1)^k q^{k(5k+1)/2} = (q)_{\infty} \sum_{\mu \in \mathcal{R}} q^{|\mu|} = \sum_{\lambda \in \mathcal{D}} \sum_{\mu \in \mathcal{R}} (-1)^{n(\lambda)} q^{|\lambda|+|\mu|}. \quad (3)$$

Hence we define

$$w((\lambda, \mu)) = (-1)^{n(\lambda)} q^{|\lambda|+|\mu|}.$$

for  $(\lambda, \mu) \in \mathcal{D} \times \mathcal{R}$ . Let  $\rho_{(j)} = (2j - 1, 2j - 3, \dots, 1) \in \mathcal{R}$ , and let  $\mathcal{E} = \{(\pi_{(j)}, \rho_{(|j|)}) : j \in \mathbf{Z}\}$ . Note that  $w((\pi_{(j)}, \rho_{(|j|)})) = (-1)^j q^{j(5j+1)/2}$ . Schur defined an involution  $\tau$  on  $(\mathcal{D} \times \mathcal{R}) - \mathcal{E}$  with the property that  $w(\tau(\lambda, \mu)) = -w(\lambda, \mu)$ . The formula (3) is an immediate consequence of the existence of such a  $\tau$ .

We shall apply  $\tau$  to the set of pairs  $(\lambda, \mu) \in \mathcal{D} \times \mathcal{R}$  in which each part of  $\lambda$  and  $\mu$  is at most  $m$ . Let  $\mathcal{R}_m = \mathcal{D}_m \cap \mathcal{R}$ : the set of partitions in  $\mathcal{R}$  having parts of size at most  $m$ . Define

$$e_{m+2}(q) = \sum_{\mu \in \mathcal{R}_m} q^{|\mu|}.$$

The polynomials  $e_{m+2}(q)$  were introduced by Schur and satisfy  $e_2(q) = 1$ ,  $e_3(q) = 1 + q$  and  $e_{m+2}(q) = e_{m+1}(q) + q^m e_m(q)$  for  $m \geq 2$ .

**Theorem 2** *The following identity holds for each integer  $m \geq 0$ :*

$$\sum_{s=0}^{\lfloor m/2 \rfloor} (-1)^s q^{s(4m-3s+5)/2} (q^{s+1})_{m-2s} e_{m-2s+2}(q) = \sum_{j=\lfloor -m/2 \rfloor}^{\lfloor m/2 \rfloor} (-1)^j q^{j(5j+1)/2}.$$

**Proof** We apply Schur's involution  $\tau$  to  $\mathcal{D}_m \times \mathcal{R}_m$  as best we can. For the definition of  $\tau$  we follow the description of Garsia and Milne [2] who used  $\tau$  to construct a bijective proof of the Rogers-Ramanujan identities.

Divide the pairs in  $(\mathcal{D} \times \mathcal{R}) - \mathcal{E}$  into three disjoint classes:

- the class  $\mathcal{T}$  contains those  $(\lambda, \mu)$  with either  $\lambda$  or  $\mu$  empty, and those with  $\lambda_1 - \mu_1 \notin \{0, 1\}$ ,
- the class  $\mathcal{A}$  contains those  $(\lambda, \mu)$  with  $\lambda_1 - \mu_1 = 1$ ,
- the class  $\mathcal{B}$  contains those  $(\lambda, \mu)$  with  $\lambda_1 - \mu_1 = 0$ .

The involution  $\tau$  will preserve  $\mathcal{T}$  and interchange  $\mathcal{A}$  and  $\mathcal{B}$ . It will also negate weights: if  $\tau((\lambda, \mu)) = (\lambda', \mu')$  then  $w((\lambda', \mu')) = -w((\lambda, \mu))$ . For  $(\lambda, \mu) \in \mathcal{T}$ , there is a unique largest part in  $\lambda$  and  $\mu$ ;  $\tau$  simply transfers this part to the other partition. Clearly  $\tau$  is an weight-negating involution on  $\mathcal{T}$ .

We divide each of the class  $\mathcal{A}$  and  $\mathcal{B}$  into three subclasses. For  $(\lambda, \mu) \in \mathcal{A} \cup \mathcal{B}$  we let  $p$  be the smallest part of  $\lambda$ ,  $q$  the slope of  $\lambda$  and  $r$  the *2-slope* of  $\mu$ , the largest integer  $r$  such that  $\mu_r = \mu_1 - 2(r - 1) > 0$ . Then

- the class  $\mathcal{A}_1$  contains those  $(\lambda, \mu) \in \mathcal{A}$  with  $\min(p, q, r) = p$ ,
- the class  $\mathcal{A}_2$  contains those  $(\lambda, \mu) \in \mathcal{A}$  with  $\min(p, q, r) = q < p$ ,
- the class  $\mathcal{A}_3$  contains those  $(\lambda, \mu) \in \mathcal{A}$  with  $\min(p, q, r) = r < \min(p, q)$ ,
- the class  $\mathcal{B}_1$  contains those  $(\lambda, \mu) \in \mathcal{B}$  with  $\min(p, q, r) = p$ ,
- the class  $\mathcal{B}_2$  contains those  $(\lambda, \mu) \in \mathcal{B}$  with  $\min(p, q, r) = r < p$ ,
- the class  $\mathcal{B}_3$  contains those  $(\lambda, \mu) \in \mathcal{B}$  with  $\min(p, q, r) = q < \min(p, r)$ .

The involution  $\tau$  will interchange  $\mathcal{A}_1$  with  $\mathcal{B}_2$ ,  $\mathcal{A}_2$  with  $\mathcal{B}_1$  and  $\mathcal{A}_3$  with  $\mathcal{B}_3$ .

We describe its action on each  $\mathcal{A}_j$ . It is then straightforward to check that  $\tau : \mathcal{A}_1 \rightarrow \mathcal{B}_2$ ,  $\tau : \mathcal{A}_2 \rightarrow \mathcal{B}_1$  and  $\tau : \mathcal{A}_3 \rightarrow \mathcal{B}_3$  are all weight-negating bijections.

Let  $(\lambda, \mu) \in \mathcal{A}_1$ . Then we obtain  $\tau((\lambda, \mu)) = (\lambda', \mu')$  by removing the smallest part  $p$  from  $\lambda$  and adding 1 to the  $p$  largest parts of  $\mu$ .

Let  $(\lambda, \mu) \in \mathcal{A}_2$ . Then  $\tau((\lambda, \mu)) = (\lambda', \mu')$  where  $\lambda' = \sigma(\lambda)$  and  $\sigma$  is the Franklin involution.

Let  $(\lambda, \mu) \in \mathcal{A}_3$ . Then we obtain  $\tau((\lambda, \mu)) = (\lambda', \mu')$  by subtracting 1 from the  $r$  largest parts of  $\mu$ , then moving the largest part of  $\lambda$  to  $\mu$  and finally adding 1 to the  $r$  largest parts of  $\lambda$ . That is  $\lambda' = (\lambda_2 + 1, \lambda_3 + 1, \dots, \lambda_{r+1} + 1, \lambda_{r+2}, \dots)$  and  $\mu' = (\lambda_1, \mu_1 - 1, \mu_2 - 1, \dots, \mu_r - 1, \mu_{r+1}, \dots)$ .

Let  $\mathcal{P}_m = \mathcal{D}_m \times \mathcal{R}_m$ . Then

$$\sum_{(\lambda, \mu) \in \mathcal{P}_m} w((\lambda, \mu)) = (q)_m e_{m+2}(q).$$

For  $1 \leq s \leq m/2$  let  $\mathcal{P}_{m,s}$  denote the set of pairs  $(\lambda, \mu) \in \mathcal{D} \times \mathcal{R}$  where  $\lambda_1 = m+1$ ,  $\mu_1 = m$ ,  $\lambda$  has slope  $s$  and smallest part  $> s$  and  $\mu$  has 2-slope  $\geq s$ . The  $\lambda \in \mathcal{D}$  with  $\lambda_1 = m+1$  having slope  $s$  and smallest part  $> s$  must have the  $s$  parts  $m+1, m, \dots, m-s+2$ , and a subset of  $\{s+1, s+2, \dots, m-s\}$ . It follows that the sum of  $(-1)^{n_\lambda} q^{|\lambda|}$  over these  $\lambda$  is

$$\prod_{j=m-s+2}^{m+1} (-q^j) \times \prod_{i=s+1}^{m-s} (1 - q^i) = (-1)^s q^{s(2m-s+3)/2} (q^{s+1})_{m-2s}.$$

The  $\mu$  in  $\mathcal{R}$  with  $\mu_1 = m$  and having slope at least  $s$  have parts  $m, m-2, \dots, m-s+2$ , together with various distinct parts  $\leq m-s$  differing by at least 2. It follows that the sum of  $q^{|\mu|}$  over these  $\mu$  is

$$q^m q^{m-2} \dots q^{m-2s+2} e_{m-2s+2}(q) = q^{s(m-s+1)} e_{m-2s+2}(q).$$

Hence

$$\begin{aligned} \sum_{(\lambda, \mu) \in \mathcal{P}_{m,s}} w((\lambda, \mu)) &= (-1)^s q^{s(2m-s+3)/2} (q^{s+1})_{m-2s} q^{s(m-s+1)} e_{m-2s+2}(q) \\ &= (-1)^s q^{s(4m-3s+5)/2} (q^{s+1})_{m-2s} e_{m-2s+2}(q). \end{aligned}$$

Let  $\mathcal{Q}_m = \mathcal{P}_m \cup \bigcup_{s=1}^{\lfloor m/2 \rfloor} \mathcal{P}_{m,s}$ . Then

$$\sum_{(\lambda, \mu) \in \mathcal{Q}_m} w((\lambda, \mu)) = \sum_{s=0}^{\lfloor m/2 \rfloor} (-1)^s q^{s(4m-3s+5)/2} (q^{s+1})_{m-2s} e_{m-2s+2}(q).$$

We claim that  $\mathcal{Q}_m - \mathcal{E}$  is closed under  $\tau$ . If  $(\lambda, \mu) \in \mathcal{P}_m$  but  $(\lambda', \mu') = \tau((\lambda, \mu)) \notin \mathcal{P}_m$  then  $(\lambda, \mu) \in \mathcal{B}_1$  and so  $(\lambda', \mu') \in \mathcal{A}_2$ . Then  $\lambda'_1 = m+1$ ,  $\mu'_1 = m$  and if  $s$  is the slope of  $\lambda'$  then all parts of  $\lambda'$  exceed  $s$  while the slope of  $\mu'$  is at least  $s$ . Hence  $\tau((\lambda, \mu)) \in \mathcal{P}_{m,s}$ . On the other hand if  $(\lambda, \mu) \in \mathcal{P}_{m,s} - \mathcal{E}$ , then  $(\lambda, \mu) \in \mathcal{A}_2$  and so  $\tau((\lambda, \mu)) \in \mathcal{P}_m$ . Hence

$$\sum_{(\lambda, \mu) \in \mathcal{Q}_m - \mathcal{E}} w((\lambda, \mu)) = 0.$$

The elements of  $\mathcal{Q}_m \cap \mathcal{E}$  are the  $(\pi_j, \rho_{|j|})$  with  $[-m/2] \leq j \leq [m/2]$ . Hence

$$\sum_{(\lambda, \mu) \in \mathcal{Q}_m} w((\lambda, \mu)) = \sum_{(\lambda, \mu) \in \mathcal{Q}_m \cap \mathcal{E}} w((\lambda, \mu))$$

$$\begin{aligned}
&= \sum_{j=\lfloor -m/2 \rfloor}^{\lfloor m/2 \rfloor} w((\pi_j, \rho_{|j|})) \\
&= \sum_{j=\lfloor -m/2 \rfloor}^{\lfloor m/2 \rfloor} (-1)^j q^{j(5j+1)/2}
\end{aligned}$$

and the theorem follows.  $\square$

The second Rogers-Ramanujan identity states that

$$\sum_{\mu \in \mathcal{R}} q^{|\mu|} = \prod_{n=1}^{\infty} \frac{1}{(1 - q^{5n-3})(1 - q^{5n-2})}$$

where  $\mathcal{R}'$  denotes the set of  $\mu \in \mathcal{R}$  with all parts at least 2. Using the Jacobi triple product, this is equivalent to

$$\sum_{k=-\infty}^{\infty} (-1)^k q^{k(5k+3)/2} = (q)_{\infty} \sum_{\mu \in \mathcal{R}'} q^{|\mu|} = \sum_{\lambda \in \mathcal{D}} \sum_{\mu \in \mathcal{R}'} (-1)^{n(\lambda)} q^{|\lambda|+|\mu|}. \quad (4)$$

There is also a bounded version of (4). To state it we define

$$d_{m+2}(q) = \sum_{\mu \in \mathcal{R}'_m} q^{|\mu|}$$

where  $\mathcal{R}'_m = \mathcal{D}_m \cap \mathcal{R}'$  is the set of partitions in  $\mathcal{R}'$  having parts of size at most  $m$ .

**Theorem 3** *The following identity holds for each integer  $m \geq 0$ :*

$$\sum_{s=0}^{\lfloor m/2 \rfloor} (-1)^s q^{s(4m-3s+5)/2} (q^{s+1})_{m-2s} d_{m-2s+2}(q) = \sum_{j=\lfloor -m/2 \rfloor}^{\lfloor m/2 \rfloor} (-1)^j q^{j(5j+3)/2}.$$

**Proof** This proof follows that of Theorem 2 *mutatis mutandis* so we do not give it in detail. We let  $\rho'_{(j)} = (2j, 2j-2, \dots, 2)$  and let  $\mathcal{E}'$  be the set of pairs  $(\pi_{(j)}, \rho'_{(j)})$  with  $j \geq 0$  and  $(\pi_{(j)}, \rho'_{(-1-j)})$  with  $j < 0$ . The map  $\tau$  is an involution on  $(\mathcal{D} \times \mathcal{R}') - \mathcal{E}'$ . The proof now follows that of Theorem 2 exactly.  $\square$

## 4 Comments

Theorems 2 and 3 appear as the main theorem (Theorem 1.1) in [4]. Warnaar's proof of these results relies on an elaborate formal argument involving Bailey chains. He leaves the formula of 1 as an exercise for the reader. He also remarks that it "seems an extremely challenging problem to find a combinatorial proof of Theorem 1.1". This paper meets that challenge.

## 5 Acknowledgment

I wish to thank David Bressoud and Frank Garvan for alerting me to the reference [2].

## References

- [1] F. Franklin, ‘Sur le développement du produit infini  $(1 - x)(1 - x^2)(1 - x^3)(1 - x^4) \dots$ ’, *C. R. Acad. Sci. Paris*, **92** (1881), 448–450.
- [2] A. M. Garsia & S. C. Milne, A Rogers-Ramanujan bijection, *J. Combin. Theory Ser. A* **31** (1981), 289–339.
- [3] I. J. Schur, Ein Beitrag sur additiven Zahlentheorie und der Theorie der Kettenbrüche, *S.-B. Preuss Akad. Wiss. Phys.-Math. Kl.* (1917), 315–336.
- [4] S. O. Warnaar, Partial-sum analogues of the Rogers-Ramanujan identities, preprint, math.CO/0108193, 2001.