

SCHWARZIAN DERIVATIVES AND UNIFORM LOCAL UNIVALENCE

MARTIN CHUAQUI, PETER DUREN, AND BRAD OSGOOD

Dedicated to Walter Hayman on the occasion of his 80th birthday

ABSTRACT. Quantitative estimates are obtained for the (finite) valence of functions analytic in the unit disk with Schwarzian derivative that is bounded or of slow growth. A harmonic mapping is shown to be uniformly locally univalent with respect to the hyperbolic metric if and only if it has finite Schwarzian norm, thus generalizing a result of B. Schwarz for analytic functions. A numerical bound is obtained for the Schwarzian norms of univalent harmonic mappings.

§1. Finite valence.

Our point of departure is a classical theorem of Nehari [14] that gives a general criterion for univalence of an analytic function in terms of its Schwarzian derivative

$$\mathcal{S}f = (f''/f')' - \frac{1}{2}(f''/f')^2.$$

A positive continuous even function $p(x)$ on the interval $(-1, 1)$ is called a *Nehari function* if $(1 - x^2)^2 p(x)$ is nonincreasing on $[0, 1)$ and no nontrivial solution u of the differential equation $u'' + pu = 0$ has more than one zero in $(-1, 1)$. Nehari's theorem can be stated as follows.

Theorem A. *Let f be analytic and locally univalent in the unit disk \mathbb{D} , and suppose its Schwarzian derivative satisfies*

$$|\mathcal{S}f(z)| \leq 2p(|z|), \quad z \in \mathbb{D}, \quad (1)$$

for some Nehari function $p(x)$. Then f is univalent in \mathbb{D} .

As special cases the theorem includes the criteria $|\mathcal{S}f(z)| \leq 2(1 - |z|^2)^{-2}$ and $|\mathcal{S}f(z)| \leq \pi^2/2$ obtained earlier by Nehari [13], as well as the criterion $|\mathcal{S}f(z)| \leq 4(1 - |z|^2)^{-1}$ stated by Pokornyi [17]. The weaker inequality

$$|\mathcal{S}f(z)| \leq \frac{2(1 + \delta^2)}{(1 - |z|^2)^2}, \quad z \in \mathbb{D},$$

1991 *Mathematics Subject Classification.* Primary 30C99, Secondary 31A05, 30C55.

Key words and phrases. Analytic function, valence, harmonic mapping, Schwarzian derivative, uniform local univalence, Schwarzian norm, minimal surface, harmonic lift.

The authors are supported by Fondecyt Grant # 1030589.

does not imply univalence, but it does imply uniform local univalence in the sense that the hyperbolic distance $d(\alpha, \beta) \geq \pi/\delta$ for any pair of points $\alpha, \beta \in \mathbb{D}$ where $f(\alpha) = f(\beta)$. In a previous paper [4] we gave a streamlined proof of this result, which is due to B. Schwarz [19], and demonstrated the sharpness of the lower bound (see also Minda [12]). Furthermore, we showed that any weaker form $|\mathcal{S}f(z)| \leq Cp(|z|)$ of Nehari's condition (1) still implies that f has finite valence if $(1 - x^2)^2p(x) \rightarrow 0$ as $x \rightarrow 1-$. In particular, if $|\mathcal{S}f(z)| \leq C$ for some constant C and all $z \in \mathbb{D}$, then f has finite valence in the unit disk.

We now derive this last result by a more elegant method, which also provides a quantitative bound for the valence in terms of the constant C . By the *valence* of f we mean $N = \sup_{w \in \mathbb{C}} n(f, w)$, where $n(f, w) \leq \infty$ is the number of points $z \in \mathbb{D}$ for which $f(z) = w$. Here is our theorem.

Theorem 1. *Let f be analytic and locally univalent in the unit disk \mathbb{D} , and suppose its Schwarzian derivative satisfies*

$$|\mathcal{S}f(z)| \leq C, \quad z \in \mathbb{D},$$

for some constant $C > \pi^2/2$. Then $|\alpha - \beta| \geq \sqrt{2/C} \pi$ for any pair of points $\alpha, \beta \in \mathbb{D}$ where $f(\alpha) = f(\beta)$. Consequently, f has finite valence and assumes any given value at most $\left(1 + \frac{\sqrt{2C}}{\pi}\right)^2$ times.

Before embarking on the proof, we recall some standard facts about the Schwarzian derivative. It is Möbius invariant: $\mathcal{S}(T \circ f) = \mathcal{S}f$ for every Möbius transformation

$$T(z) = \frac{az + b}{cz + d}, \quad ad - bc \neq 0.$$

Also, $\mathcal{S}(f \circ T) = ((\mathcal{S}f) \circ T)T'^2$. For any analytic function ψ , the functions f with Schwarzian $\mathcal{S}f = 2\psi$ are precisely those of the form $f = u_1/u_2$, where u_1 and u_2 are linearly independent solutions of the differential equation $u'' + \psi u = 0$. Thus if $\mathcal{S}f = 2\psi$, then $f(\alpha) = f(\beta)$ if and only if some solution of the differential equation $u'' + \psi u = 0$ vanishes at α and β .

We will make use of the following lemma, which is a variant of a lemma in [7].

Lemma 1. *Suppose that $u = u(z)$ is a solution of the differential equation $u'' + \psi u = 0$ for some function ψ analytic in \mathbb{D} . Let $z = z(s)$, $s \in (0, b)$, be an arclength parametrization of a line segment in \mathbb{D} , and suppose that $v(z) = |u(z(s))| > 0$ for s in the interval $(0, b)$. Then*

$$v''(s) + |\psi(z(s))|v(s) \geq 0, \quad 0 < s < b.$$

Proof of lemma. Differentiation of $v^2 = u\bar{u}$ gives

$$v(s)v'(s) = \operatorname{Re}\left\{u'(z(s))z'(s)\overline{u(z(s))}\right\}, \quad 0 < s < b.$$

But $v(s) > 0$ and $|z'(s)| = 1$, so it follows that

$$v(s) |v'(s)| \leq |u'(z(s))| v(s), \quad \text{or} \quad |v'(s)| \leq |u'(z(s))|.$$

Differentiation of vv' gives

$$vv'' + v'^2 = \operatorname{Re}\{u''z'^2\bar{u} + |u'|^2\},$$

since $|z'(s)| = 1$ and $z'(s)$ is constant for the parametrization of a line segment. Introducing the differential equation $u'' = -\psi u$, we conclude that

$$\begin{aligned} vv'' + v'^2 &= |u'|^2 - \operatorname{Re}\{\psi |u|^2 z'^2\} \geq |u'|^2 - |\psi||u|^2 \\ &\geq |v'|^2 - |\psi||u|^2 = v'^2 - |\psi|v^2. \end{aligned}$$

Therefore, $v(v'' + |\psi|v) \geq 0$, and the desired result follows because $v(s) > 0$ on the interval $(0, b)$. \square

Proof of theorem. Under the hypothesis $|\psi(z)| \leq C/2$, where $\mathcal{S}f = 2\psi$, suppose that $f(\alpha) = f(\beta)$ for some pair of distinct points $\alpha, \beta \in \mathbb{D}$. Then some solution of the differential equation $u'' + \psi u = 0$ vanishes at α and β . Without loss of generality, we may suppose that $u(z) \neq 0$ on the open line segment with endpoints α and β . Let $z = z(s)$ be the parametrization of this segment by arclength s , where $z(0) = \alpha$ and $z(b) = \beta$, so that $b = |\alpha - \beta|$. Then by Lemma 1, the function $v(s) = |u(z(s))|$ has the properties $v(0) = v(b) = 0$, $v(s) > 0$, and

$$v''(s) + |\psi(z(s))|v(s) \geq 0, \quad 0 < s < b.$$

We now apply the Sturm comparison theorem (see for instance [1]). Note that $v(s)$ is a real-valued function that satisfies the differential equation $v''(s) + g(s)v(s) = 0$, with

$$g(s) = -v''(s)/v(s) \leq |\psi(z(s))| \leq C/2.$$

On the other hand, the solutions of the differential equation $y'' + (C/2)y = 0$ are sinusoids whose zeros are separated by the distance $\sqrt{2/C}\pi$. By the Sturm comparison theorem,

$$|\alpha - \beta| = b \geq \sqrt{2/C}\pi,$$

as claimed. Note that if $C = \pi^2/2$, then the argument shows that $|\alpha - \beta| \geq 2$, and so we recover Nehari's theorem that f is univalent in \mathbb{D} if $|\mathcal{S}f(z)| \leq \pi^2/2$.

Now for the estimate of valence. Let w be an arbitrary complex number. By what we have already proved, the points in \mathbb{D} where $f(z) = w$ are the centers of disjoint disks of radius $\pi/\sqrt{2C}$. If there are N such points, a comparison of areas shows that

$$N\pi \left(\frac{\pi}{\sqrt{2C}}\right)^2 \leq \pi \left(1 + \frac{\pi}{\sqrt{2C}}\right)^2,$$

which reduces to the stated inequality $N \leq \left(1 + \frac{\sqrt{2C}}{\pi}\right)^2$. \square

The bound on the valence is not sharp. For instance, for $C = \pi^2/2$ it gives $n \leq 4$, whereas Nehari's theorem shows that $n \leq 1$. Nevertheless, the question remains whether the bound is sharp in order of magnitude. Theorem 1 shows that under the condition $|\mathcal{S}f(z)| \leq C$ the sharp bound on the valence is $O(C)$ as $C \rightarrow \infty$. On the other hand, the simple example

$$f(z) = \tan\left(\sqrt{C/2}z\right), \quad \text{for which } \mathcal{S}f(z) = C > \pi^2/2, \quad (2)$$

shows that the valence may increase as fast as \sqrt{C} . Indeed, $f(x) = 0$ for all points $x = \pm k\pi\sqrt{2/C}$ where $k = 0, 1, 2, \dots$, and at least $\frac{\sqrt{2C}}{\pi} - 1$ of these points lie in the unit disk. Thus the bound on the valence cannot be improved to anything better than $O(\sqrt{C})$ as $C \rightarrow \infty$.

§2. Schwarzians of slow growth.

We showed in [4] that for each Nehari function $p(x)$ with $(1-x^2)^2p(x) \rightarrow 0$ as $x \rightarrow 1-$, any condition of the form $|\mathcal{S}f(z)| \leq Cp(|z|)$ implies that f has finite valence in the disk. In the previous section we considered functions with $|\mathcal{S}f(z)| \leq C$ and obtained an explicit estimate for the valence in terms of C . We now take $p(x) = \frac{2}{1-x^2}$, the Nehari function in the univalence criterion of Pokornyi [17], and derive an estimate, in terms of the constant C , for the (finite) valence of functions f with $|\mathcal{S}f(z)| \leq Cp(|z|)$. We will content ourselves with an asymptotic estimate as $C \rightarrow \infty$, although the proof can be adapted to yield an explicit bound.

Theorem 2. *Let f be analytic and locally univalent in \mathbb{D} , and suppose its Schwarzian derivative satisfies*

$$|\mathcal{S}f(z)| \leq \frac{2C}{1-|z|^2}, \quad z \in \mathbb{D}, \quad (3)$$

for a constant $C > 2$. Then f has finite valence $N = N(C) \leq AC \log C$, where A is some absolute constant.

The proof of Theorem 2 will invoke the separation result of Theorem 1. The following geometric lemma will be useful.

Lemma 2. *If n points z_1, z_2, \dots, z_n lie in an annulus $\rho \leq |z| \leq \rho + d \leq 1$ and have the separation property $|z_j - z_k| \geq 2d$ for $j \neq k$, then $n \leq 2\pi/d$.*

Proof of lemma. It will suffice to show that $|\arg\{z_j\} - \arg\{z_k\}| > d$ for $j \neq k$. But if $|\arg\{z_j\} - \arg\{z_k\}| \leq d$ for some $j \neq k$, then by the triangle inequality

$$|z_j - z_k| \leq d + \rho d < 2d,$$

which contradicts the hypothesis. \square

Proof of theorem. In terms of the Nehari function $p(x) = 2/(1 - x^2)$, the hypothesis is that $|\mathcal{S}f(z)| \leq Cp(|z|)$. We claim that f is univalent in the disk $|z| < r_0 = \pi/\sqrt{\pi^2 + 4C}$. Indeed, the function $g(z) = f(r_0z)$ has Schwarzian $\mathcal{S}g(z) = r_0^2 \mathcal{S}f(r_0z)$, and so

$$|\mathcal{S}g(z)| \leq r_0^2 Cp(r_0) = \frac{\pi^2}{2},$$

which implies that g is univalent in \mathbb{D} , by Nehari's theorem. Thus f is univalent in $|z| < r_0$.

We now define the sequence $\{r_k\}$ recursively by the formula

$$r_k - r_{k-1} = d_k = \frac{\pi}{\sqrt{2Cp(r_k)}}, \quad k = 1, 2, \dots \quad (4)$$

If $r_k < 1$, then since $|\mathcal{S}f(z)| \leq Cp(r_k)$ in the disk $|z| \leq r_k$, the Schwarzian of $g(z) = f(r_kz)$ satisfies $|\mathcal{S}g(z)| \leq Cr_k^2 p(r_k)$ in \mathbb{D} . Thus by Theorem 1, if $f(\alpha) = f(\beta)$ for two points α and β in the disk $|z| < r_k$, then

$$|\alpha - \beta| \geq \frac{r_k \sqrt{2} \pi}{\sqrt{Cr_k^2 p(r_k)}} = \frac{\sqrt{2} \pi}{\sqrt{Cp(r_k)}} = 2d_k.$$

An appeal to Lemma 2 now shows that the valence N_k of f in the annulus $r_{k-1} \leq |z| < r_k$ satisfies

$$N_k \leq \frac{2\pi}{d_k} = 2\sqrt{2Cp(r_k)}. \quad (5)$$

Next we make a closer examination of the recurrence relation (4), which we rewrite as

$$x - a = \varepsilon \sqrt{1 - x^2}, \quad \text{where } a = r_{k-1}, \quad x = r_k, \quad \text{and } \varepsilon = \frac{\pi}{2\sqrt{C}}.$$

Squaring and solving the quadratic equation, we find

$$x = \frac{1}{1 + \varepsilon^2} \left(a + \varepsilon \sqrt{1 - a^2 + \varepsilon^2} \right),$$

which leads after further calculation to the formula

$$\frac{1}{x - a} = \frac{\sqrt{1 - a^2 + \varepsilon^2} + \varepsilon a}{\varepsilon(1 - a^2)} = \phi(a), \quad \text{say.}$$

It is important to observe that ϕ is an increasing function on the interval $0 < a < 1$. This be verified by computing its derivative:

$$\varepsilon(1 - a^2)^2 \sqrt{1 - a^2 + \varepsilon^2} \phi'(a) = a(1 - a^2) + 2a\varepsilon^2 + \varepsilon(1 + a^2) \sqrt{1 - a^2 + \varepsilon^2} > 0.$$

Reverting to the original notation, we have $d_k \phi(r_{k-1}) = 1$.

Now let $R = (1 - \frac{1}{4C})^{1/2}$, and observe that $R > R_0$ because $C > 2$. Define the index m by the condition $r_m \leq R < r_{m+1}$. By virtue of (5), the valence of the function f in the disk $|z| \leq r_m$ is bounded by

$$\begin{aligned} 1 + \sum_{k=1}^m N_k &\leq 1 + 2\pi \sum_{k=1}^m \frac{1}{d_k} = 1 + 2\pi \sum_{k=1}^m \phi(r_{k-1})^2 (r_k - r_{k-1}) \\ &\leq 1 + 2\pi \int_0^R \phi(x)^2 dx, \end{aligned}$$

since ϕ is an increasing function. Thus we need to estimate the integral

$$\begin{aligned} \int_0^R \phi(x)^2 dx &= \frac{1}{\varepsilon^2} \int_0^R \frac{(\sqrt{1 + \varepsilon^2 - x^2} + \varepsilon x)^2}{(1 - x^2)^2} dx \\ &= \frac{1}{\varepsilon^2} \int_0^R \frac{dx}{1 - x^2} + \int_0^R \frac{1 + x^2}{(1 - x^2)^2} dx + \frac{2}{\varepsilon} \int_0^R \frac{x \sqrt{1 + \varepsilon^2 - x^2}}{(1 - x^2)^2} dx \\ &= I_1 + I_2 + I_3, \quad \text{say.} \end{aligned}$$

Recall that $R^2 = 1 - \frac{1}{4C}$ and $\varepsilon = \frac{\pi}{2\sqrt{C}}$, so that

$$\begin{aligned} I_1 &\leq \frac{4C}{\pi^2} \int_0^R \frac{dx}{1 - x^2} = \frac{2C}{\pi^2} \log \frac{(1 + R)^2}{1 - R^2} \\ &\leq \frac{2C}{\pi^2} \log(16C) = O(C \log C). \end{aligned}$$

On the other hand,

$$I_2 = O\left(\frac{1}{1 - R^2}\right) = O(C),$$

whereas an integration by parts gives

$$\begin{aligned} I_3 &= \frac{1}{\varepsilon} \left\{ \left[\frac{\sqrt{1 + \varepsilon^2 - x^2}}{1 - x^2} \right]_0^R + \int_0^R \frac{x dx}{(1 - x^2)\sqrt{1 + \varepsilon^2 - x^2}} \right\} \\ &\leq O(C) + \frac{1}{\varepsilon} \int_0^R \frac{x dx}{(1 - x^2)^{3/2}} = O(C). \end{aligned}$$

If $r_m < R$, the same argument that produced the estimate (5) shows that in the annulus $r_m \leq |z| < R$ the valence of f is no greater than $2\sqrt{2C} p(R) = O(C)$.

To complete the proof, we need to estimate the valence of f in the annulus $R \leq |z| < 1$. The radius $R = (1 - \frac{1}{4C})^{1/2}$ is chosen so that $(1 - R^2)^2 C p(R) = \frac{1}{2}$. The radius $R_1 = (1 - \frac{1}{2C})^{1/2}$ has the properties $0 < R_1 < R$ and

$$(1 - R_1^2)^2 C p(R_1) = 2C(1 - R_1^2) = 1.$$

Thus the bound (3) on the Schwarzian derivative of f implies that

$$|\mathcal{S}f(z)| \leq \frac{1}{(1 - |z|^2)^2}, \quad R_1 \leq |z| < 1. \quad (6)$$

Suppose now that $f(\alpha) = f(\beta)$ for two points α and β in the annulus $R \leq |z| < 1$. Then by Nehari's theorem, or rather by its proof, the hyperbolic geodesic joining α and β cannot lie entirely in the annulus $R_1 \leq |z| < 1$. For then the Schwarzian of f would satisfy (6) along such a geodesic. By a well-known technique of Nehari [14], this would imply that a function $g = f \circ \varphi$, where φ is a suitable conformal automorphism of the disk, satisfies $|\mathcal{S}g(x)| \leq (1 - x^2)^{-2}$ on the real interval $-1 < x < 1$ and has the property $g(a) = g(b)$ for a pair of distinct points a and b in that interval. Equivalently, a solution to the associated linear differential equation vanishes at two points of the interval $(-1, 1)$, which is not possible. This shows that f is univalent in each part of the annulus $R \leq |z| < 1$ which lies inside the arch of some hyperbolic geodesic entirely contained in the larger annulus $R_1 \leq |z| < 1$. The conclusion is strongest if we take the hyperbolic geodesic to be tangent to the circle $|z| = R_1$.

The estimate of valence in the annulus $R \leq |z| < 1$ now reduces to a covering problem, namely to estimate the number of curvilinear rectangles required to cover the annulus. Here a *curvilinear rectangle* is understood to mean the intersection of the given annulus with the region inside a hyperbolic geodesic that is tangent to the circle $|z| = R_1$. Observe that the geodesic that is tangent to this circle at the point $z = R_1$ is the image of the imaginary axis under the Möbius automorphism

$$T(z) = \frac{z + R_1}{1 + R_1 z}, \quad z \in \mathbb{D}.$$

In order to locate the two points where this geodesic meets the circle $|z| = R$, we calculate that $|T(iy)| = R$ implies

$$y^2 = \frac{R^2 - R_1^2}{1 - R^2 R_1^2} = \frac{2C}{6C - 1}.$$

Choosing $y > 0$, we find by further calculation that

$$\begin{aligned} \arg\{T(iy)\} &= \tan^{-1} \left(\frac{y}{1 + y^2} \frac{1 - R_1^2}{R_1} \right) \\ &= \tan^{-1} \left(\frac{\sqrt{6C - 1}}{8C - 1} \frac{1}{\sqrt{2C - 1}} \right) \geq \tan^{-1} \left(\frac{\sqrt{3}}{8C} \right) \geq \frac{1}{5C} \end{aligned}$$

for all constants C sufficiently large. Therefore, the annulus $R \leq |z| < 1$ is contained in the union of at most $[5\pi C] + 1$ curvilinear rectangles of the type described, where

$[x]$ denotes the integer part of x . Consequently, the valence of f in this annulus is $O(C)$ as $C \rightarrow \infty$. This concludes the proof of Theorem 2. \square

The example (2) again shows that the estimate of valence in Theorem 2 cannot be improved to $o(\sqrt{C})$. In search of a better lower bound, it is natural to investigate the zeros of solutions of the differential equation

$$y'' + \frac{C}{1-x^2}y = 0 \quad (7)$$

in the interval $(-1, 1)$. The solutions of (7) are easily seen to have the form $y = (1-x^2)u'$, where u is a solution of the Legendre equation

$$(x^2 - 1)u''(x) + 2xu'(x) - Cu(x) = 0. \quad (8)$$

(Compare Kamke [10], 2.240, eq. 14, p. 460.) If $C = n(n+1)$ for $n = 1, 2, \dots$, one solution of (8) is the Legendre polynomial $u = P_n(x)$, which is known to have exactly n simple zeros in the interval $(-1, 1)$. Thus by Rolle's theorem, the derivative $P'_n(x)$, a polynomial of degree $n-1$, has exactly $n-1$ zeros in $(-1, 1)$. In other words, if $C = n(n+1)$, then some solution of (7) has at least $n-1$ zeros in the unit disk. This remains true, by the Sturm comparison theorem, if $n(n+1) < C < (n+1)(n+2)$. The conclusion is that some analytic function whose Schwarzian satisfies (3) has valence (loosely speaking) at least \sqrt{C} , which shows again that the asymptotic estimate of Theorem 2 cannot be improved beyond $N = O(\sqrt{C})$ as $C \rightarrow \infty$.

§3. Uniform local univalence and harmonic mappings.

The *pseudohyperbolic metric* is defined by

$$\rho(\alpha, \beta) = \left| \frac{\alpha - \beta}{1 - \bar{\alpha}\beta} \right|, \quad \alpha, \beta \in \mathbb{D},$$

and is Möbius invariant. More precisely, $\rho(\varphi(\alpha), \varphi(\beta)) = \rho(\alpha, \beta)$ if φ is any Möbius self-mapping of \mathbb{D} . The *pseudohyperbolic disk* with center α and radius r is defined by

$$\Delta(\alpha, r) = \{z \in \mathbb{D} : \rho(z, \alpha) < r\}.$$

It is a true Euclidean disk, but α and r are not the Euclidean center and radius unless $\alpha = 0$. The *hyperbolic metric* is

$$d(\alpha, \beta) = \frac{1}{2} \log \frac{1 + \rho(\alpha, \beta)}{1 - \rho(\alpha, \beta)}.$$

The *Schwarzian norm* of a function f analytic and locally univalent in the unit disk is defined by

$$\|\mathcal{S}f\| = \sup_{z \in \mathbb{D}} (1 - |z|^2)^2 |\mathcal{S}f(z)|.$$

It is Möbius invariant in the sense that $\|\mathcal{S}(f \circ \varphi)\| = \|\mathcal{S}f\|$ for any Möbius self-mapping φ of the unit disk. The previously mentioned result of Nehari [13], a special case of Theorem A, can be rephrased to say that f is univalent in \mathbb{D} if $\|\mathcal{S}f\| \leq 2$. In the converse direction, Kraus [11] showed that $\|\mathcal{S}f\| \leq 6$ whenever f is analytic and univalent in \mathbb{D} . The bound is sharp, since the Koebe function $k(z) = z/(1 - z)^2$ has Schwarzian

$$\mathcal{S}k(z) = -\frac{6}{(1 - z^2)^2}.$$

According to the theorem of B. Schwarz [19], the condition $\|\mathcal{S}f\| < \infty$ implies that f is *uniformly locally univalent* in the hyperbolic metric, or equivalently in the pseudohyperbolic metric. Specifically, this means that for some radius $r > 0$, the function f is univalent in every pseudohyperbolic disk $\Delta(\alpha, r)$. Conversely, if f is uniformly locally univalent, then $\|\mathcal{S}f\| < \infty$. In fact, it is known that $\|\mathcal{S}f\| \leq 6/r^2$. To see this, suppose that f is univalent in every pseudohyperbolic disk $\Delta(\alpha, r)$. For any fixed $\alpha \in \mathbb{D}$, the Möbius transformation

$$\varphi(z) = \frac{rz + \alpha}{1 + \bar{\alpha}rz}, \quad z \in \mathbb{D},$$

maps \mathbb{D} onto $\Delta(\alpha, r)$. Thus $g = f \circ \varphi$ is univalent in \mathbb{D} , and so $\|\mathcal{S}g\| \leq 6$ by Kraus' theorem. In particular, $|\mathcal{S}g(0)| \leq 6$. But

$$\mathcal{S}g(0) = \mathcal{S}(f \circ \varphi)(0) = ((\mathcal{S}f)(\varphi(0)))\varphi'(0)^2 = r^2(1 - |\alpha|^2)^2 \mathcal{S}f(\alpha),$$

and so $r^2(1 - |\alpha|^2)^2 |\mathcal{S}f(\alpha)| \leq 6$. Taking the supremum over all $\alpha \in \mathbb{D}$, we conclude that $\|\mathcal{S}f\| \leq 6/r^2$.

To what extent do these relations generalize to harmonic mappings? A complex-valued harmonic function in a simply connected domain has the canonical representation $f = h + \bar{g}$, unique up to an additive constant, where h and g are analytic functions. By a theorem of H. Lewy (see [9]), the Jacobian $|h'|^2 - |g'|^2$ of a locally univalent harmonic mapping never vanishes. The harmonic mappings with positive Jacobian are said to be *orientation-preserving*. These are harmonic mappings whose *dilatation* $\omega = g'/h'$ is an analytic function with $|\omega(z)| < 1$. An orientation-preserving harmonic mapping lifts to a mapping \tilde{f} onto a minimal surface described by conformal parameters, if and only if $\omega = q^2$, the square of some analytic function q . For such mappings f we have defined [2] the *Schwarzian derivative* by the formula

$$\mathcal{S}f = 2(\sigma_{zz} - \sigma_z^2),$$

where $\sigma = \log(|h'| + |g'|)$ and

$$\sigma_z = \frac{\partial \sigma}{\partial z} = \frac{1}{2} \left(\frac{\partial \sigma}{\partial x} - i \frac{\partial \sigma}{\partial y} \right), \quad z = x + iy.$$

If f is analytic, $\mathcal{S}f$ is the classical Schwarzian. If f is harmonic and φ is analytic, then $f \circ \varphi$ is harmonic and

$$\mathcal{S}(f \circ \varphi) = ((\mathcal{S}f) \circ \varphi)\varphi'^2 + \mathcal{S}\varphi,$$

a generalization of the classical formula for analytic functions f . In particular,

$$\mathcal{S}(f \circ \varphi) = ((\mathcal{S}f) \circ \varphi)\varphi'^2$$

if φ is a Möbius self-mapping of the disk. From this it follows that the Schwarzian norm

$$\|\mathcal{S}f\| = \sup_{z \in \mathbb{D}} (1 - |z|^2)^2 |\mathcal{S}f(z)|.$$

of a harmonic mapping retains the Möbius invariance property $\|\mathcal{S}(f \circ \varphi)\| = \|\mathcal{S}f\|$.

Theorem 3. *Let $f = h + \bar{g}$ be an orientation-preserving harmonic mapping whose dilatation $\omega = g'/h'$ is the square of an analytic function in the unit disk. Then $\|\mathcal{S}f\| < \infty$ if and only if f is uniformly locally univalent.*

The proof will invoke a recent result of Chuaqui and Hernández [6], which we state here for reference.

Theorem B. *Let $f = h + \bar{g}$ be an orientation-preserving harmonic mapping in the unit disk, and suppose that h is univalent and $h(\mathbb{D})$ is convex. Then f is univalent in \mathbb{D} .*

Proof of Theorem B. In the paper [6] this result comes out of a more general argument, but the proof for this special case is so short that we include it here for completeness. If $f(z_1) = f(z_2)$, then $h(z_1) - h(z_2) = \overline{g(z_2)} - \overline{g(z_1)}$. With the notation $w_1 = h(z_1)$ and $w_2 = h(z_2)$, this can be written as

$$\overline{w_1} - \overline{w_2} = \varphi(w_2) - \varphi(w_1), \quad \text{where } \varphi = g \circ h^{-1}.$$

But φ is analytic on the convex domain $h(\mathbb{D})$, so this says that

$$\overline{w_1} - \overline{w_2} = \int_{w_1}^{w_2} \varphi'(w) dw,$$

where the integral is taken over a straight-line segment. However, this is not possible, because $|\varphi'(w)| = |g'(z)/h'(z)| < 1$ by the hypothesis that f is orientation-preserving. \square

We will also need a result that is implicit in work of Sheil-Small [20]. An analytic or harmonic function f is said to be *uniformly locally convex* if there exists a radius $r > 0$ such that f maps every pseudohyperbolic disk $\Delta(\alpha, r)$ univalently onto a convex region.

Theorem C. *Let $f = h + \bar{g}$ be an orientation-preserving harmonic mapping that is uniformly locally univalent in the unit disk. Then its analytic part h is uniformly locally convex.*

Proof of Theorem C. Suppose first that f is univalent in the entire disk \mathbb{D} . Then we may assume without loss of generality that $f \in S_H$, the class of orientation-preserving univalent harmonic mappings of \mathbb{D} for which $h(0) = g(0) = 0$ and $h'(0) = 1$. The analytic part of such a mapping has the power series expansion $h(z) = z + a_2 z^2 + \dots$, and it is a result of Clunie and Sheil-Small that the coefficients a_2 have an absolute bound; in other words, $\lambda = \sup_{f \in S_H} |a_2|$ is finite. It is conjectured that $\lambda = 3$, but the best bound currently known (see [9], p. 97) is approximately $\lambda < 49$. Now if $f \in S_H$, then for each fixed $\zeta \in \mathbb{D}$ the function

$$F(z) = \frac{f\left(\frac{z+\zeta}{1+\bar{\zeta}z}\right) - f(\zeta)}{(1-|\zeta|^2)h'(\zeta)} = H(z) + \overline{G(z)}$$

also belongs to the class S_H , so that $|\frac{1}{2}H''(0)| \leq \lambda$. But a calculation gives

$$H''(0) = (1-|\zeta|^2)\frac{h''(\zeta)}{h'(\zeta)} - 2\bar{\zeta},$$

so we have

$$\left| \frac{\zeta h''(\zeta)}{h'(\zeta)} - \frac{2\rho^2}{1-\rho^2} \right| \leq \frac{2\lambda\rho}{1-\rho^2},$$

which implies that

$$\operatorname{Re} \left\{ 1 + \frac{\zeta h''(\zeta)}{h'(\zeta)} \right\} \geq \frac{1-2\lambda\rho+\rho^2}{1-\rho^2} > 0$$

for $|\zeta| = \rho < \mu = \lambda - \sqrt{\lambda^2 - 1}$. By the familiar analytic criterion for convexity (see [8], p. 42), this shows that $h(z)$ is convex in the disk $|z| < \mu$. If f is assumed to be univalent only in the subdisk $|z| < r$, the preceding result can be adapted to show that h is univalent in the disk $|z| < \mu r$. If f is univalent in the pseudohyperbolic disk $\Delta(\alpha, r)$, then for a suitable Möbius self-mapping φ of \mathbb{D} the composite function $\Phi = f \circ \varphi$ is univalent in $\Delta(0, r)$, and so its analytic part is convex in $\Delta(0, \mu r)$, which implies that the analytic part h of $f = \Phi \circ \varphi^{-1}$ is convex in $\Delta(\alpha, \mu r)$. Since this is true for each $\alpha \in \mathbb{D}$, the conclusion is that h is uniformly locally convex. \square

Proof of Theorem 3. We showed in [4] that $\|\mathcal{S}f\| < \infty$ if and only if $\|\mathcal{S}h\| < \infty$. Therefore, if $\|\mathcal{S}f\| < \infty$, then $\|\mathcal{S}h\| < \infty$, and so h is uniformly locally univalent, by the theorem of B. Schwarz. In other words, h is univalent on every pseudohyperbolic disk $\Delta(\alpha, r)$ for some fixed radius r . Then by the classical result on radius of convexity (see [8], p. 44), h maps every disk $\Delta(\alpha, (2 - \sqrt{3})r)$ to a convex domain.

It now follows from Theorem B that f is univalent in each disk $\Delta(\alpha, (2 - \sqrt{3})r)$. Thus f is uniformly locally univalent.

Conversely, suppose the harmonic mapping f is uniformly locally univalent in \mathbb{D} . Then by Theorem C its analytic part h is uniformly locally convex, hence uniformly locally univalent. Therefore, $\|\mathcal{S}h\| < \infty$ by Kraus' theorem, as discussed at the beginning of this section. It now follows from our result in [4] that $\|\mathcal{S}f\| < \infty$. \square

As a corollary of the proof, we are able to establish a numerical bound on $\|\mathcal{S}f\|$ for univalent harmonic mappings f , analogous to Kraus' bound $\|\mathcal{S}f\| \leq 6$ for analytic univalent functions in the disk. By Möbius invariance we may assume without loss of generality that the harmonic mapping $f = h + \bar{g}$ belongs to the class S_H . Then as shown in the proof of Theorem C, its analytic part $h(z)$ is convex in the disk $|z| < \mu$, where $\mu = \lambda - \sqrt{\lambda^2 - 1}$. Thus the function $H(z) = h(\mu z)$ is convex in \mathbb{D} , so it has Schwarzian norm $\|\mathcal{S}H\| \leq 2$, by a result of Nehari [15]. Since $\|\mathcal{S}H\| = \mu^2 \|\mathcal{S}h\|$, it follows that $\|\mathcal{S}h\| \leq 2/\mu^2$. Consequently, the estimate $\lambda < 49$ shows that $\|\mathcal{S}h\| < 19,204$. On the other hand, a result of Pommerenke [18] implies that

$$\|\mathcal{S}f\| \leq \|\mathcal{S}h\| + 2 \left(1 + \frac{1}{2}\|\mathcal{S}h\|\right)^{1/2} + 7,$$

as we showed in [4]. Inserting the preceding estimate $\|\mathcal{S}h\| < 19,204$, we obtain the absolute bound $\|\mathcal{S}f\| < 19,407$ for all harmonic mappings f that are univalent in \mathbb{D} and have dilatation that is a perfect square.

It is an open problem to determine the sharp bound. We showed in [4] that $\|\mathcal{S}f\| \leq 45$ for all mappings f with dilatation $\omega = q^2$ that are convex in the horizontal direction. We also observed that the horizontal shear of the Koebe function with dilatation $\omega(z) = z^2$ has Schwarzian

$$\mathcal{S}f = -4 \left(\frac{1}{1-z} + \frac{\bar{z}}{1+|z|^2} \right)^2,$$

from which an easy calculation gives $\|\mathcal{S}f\| = 16$. These results are unchanged if the Koebe function is sheared with dilatation $\omega(z) = e^{i\theta} z^2$ for any θ . Therefore, since the Koebe function maximizes the Schwarzian norm for analytic univalent functions, it is reasonable to conjecture that $\|\mathcal{S}f\| \leq 16$ for all univalent harmonic mappings in the disk whose dilatation is a perfect square.

§4. Bounds on valence of harmonic lifts.

Theorems 1 and 2 extend readily to the lifts of harmonic mappings to minimal surfaces. In [3] we obtained the following generalization of Nehari's theorem.

Theorem D. *Let $f = h + \bar{g}$ be a harmonic mapping of the unit disk, with conformal parameter $e^{\sigma(z)} = |h'(z)| + |g'(z)| \neq 0$ and dilatation $g'/h' = q^2$ for some meromorphic function q . Let \tilde{f} denote the Weierstrass–Enneper lift of f to a minimal surface with Gauss curvature $K = K(\tilde{f}(z))$ at the point $\tilde{f}(z)$. Suppose that*

$$|\mathcal{S}f(z)| + e^{2\sigma(z)} |K(\tilde{f}(z))| \leq 2p(|z|), \quad z \in \mathbb{D},$$

for some Nehari function p . Then \tilde{f} is univalent in \mathbb{D} .

The valence estimates for analytic functions in Theorems 1 and 2 have corresponding generalizations to harmonic lifts.

Theorem 1'. *Let $f = h + \bar{g}$ be a harmonic mapping of the unit disk with conformal parameter $e^{\sigma(z)} = |h'(z)| + |g'(z)| \neq 0$ and dilatation $g'/h' = q^2$ for some meromorphic function q , and let \tilde{f} be its lift to a minimal surface with Gauss curvature K . Suppose that*

$$|\mathcal{S}f(z)| + e^{2\sigma(z)}|K(\tilde{f}(z))| \leq C, \quad z \in \mathbb{D},$$

for some constant $C > \pi^2/2$. Then $|\alpha - \beta| \geq \sqrt{2/C}\pi$ for any pair of points $\alpha, \beta \in \mathbb{D}$ where $\tilde{f}(\alpha) = \tilde{f}(\beta)$. Consequently, the lift \tilde{f} has finite valence and meets any given point at most $\left(1 + \frac{\sqrt{2C}}{\pi}\right)^2$ times.

Theorem 2'. *Let a harmonic mapping $f = h + \bar{g}$ be as in Theorem 1' but satisfy the inequality*

$$|\mathcal{S}f(z)| + e^{2\sigma(z)}|K(\tilde{f}(z))| \leq \frac{2C}{1 - |z|^2}, \quad z \in \mathbb{D},$$

for some constant $C > 2$. Then its lift \tilde{f} has finite valence $N = N(C) \leq AC \log C$, where A is some absolute constant.

The proofs of Theorems 1' and 2' reduce ultimately to the same consideration of zeros of solutions to differential equations as in the proofs of Theorems 1 and 2. Here the link with differential equations and the Sturm theory comes from a result of Chuaqui and Gevirtz [5], as developed in our earlier work [3,4]. The details are relatively straightforward and will not be pursued here.

REFERENCES

1. G. Birkhoff and G.-C. Rota, *Ordinary Differential Equations*, 4th Edition, Wiley, New York, 1989.
2. M. Chuaqui, P. Duren, and B. Osgood, *The Schwarzian derivative for harmonic mappings*, J. Analyse Math. **91** (2003), 329–351.
3. M. Chuaqui, P. Duren, and B. Osgood, *Univalence criteria for lifts of harmonic mappings to minimal surfaces*, J. Geom. Analysis, to appear.
4. M. Chuaqui, P. Duren, and B. Osgood, *Schwarzian derivative criteria for valence of analytic and harmonic mappings*, Math. Proc. Cambridge Philos. Soc., to appear.
5. M. Chuaqui and J. Gevirtz, *Simple curves in \mathbb{R}^n and Ahlfors' Schwarzian derivative*, Proc. Amer. Math. Soc. **132** (2004), 223–230.
6. M. Chuaqui and R. Hernández, *Univalent harmonic mappings and linearly connected domains*, J. Math. Anal. Appl., to appear.
7. M. Chuaqui and B. Osgood, *Sharp distortion theorems associated with the Schwarzian derivative*, J. London Math. Soc. **48** (1993), 289–298.
8. P. L. Duren, *Univalent Functions*, Springer-Verlag, New York, 1983.
9. P. Duren, *Harmonic Mappings in the Plane*, Cambridge University Press, Cambridge, U. K., 2004.

10. E. Kamke, *Differentialgleichungen: Lösungsmethoden und Lösungen, Band 1: Gewöhnliche Differentialgleichungen*, 3. Auflage, Becker & Erler, Leipzig, 1944; reprinted by Chelsea Publishing Co., New York, 1948.
11. W. Kraus, *Über den Zusammenhang einiger Charakteristiken eines einfach zusammenhängenden Bereiches mit der Kreisabbildung*, Mitt. Math. Sem. Giessen **21** (1932), 1–28.
12. D. Minda, *The Schwarzian derivative and univalence criteria*, Contemporary Math. **38** (1985), 43–52.
13. Z. Nehari, *The Schwarzian derivative and schlicht functions*, Bull. Amer. Math. Soc. **55** (1949), 545–551.
14. Z. Nehari, *Some criteria of univalence*, Proc. Amer. Math. Soc. **5** (1954), 700–704.
15. Z. Nehari, *A property of convex conformal maps*, J. Analyse Math. **30** (1976), 390–393.
16. Z. Nehari, *Univalence criteria depending on the Schwarzian derivative*, Illinois J. Math. **23** (1979), 345–351.
17. V. V. Pokornyi, *On some sufficient conditions for univalence*, Dokl. Akad. Nauk SSSR **79** (1951), 743–746 (in Russian).
18. Ch. Pommerenke, *Linear-invariante Familien analytischer Funktionen I*, Math. Annalen **155** (1964), 108–154.
19. B. Schwarz, *Complex nonoscillation theorems and criteria of univalence*, Trans. Amer. Math. Soc. **80** (1955), 159–186.
20. T. Sheil-Small, *Constants for planar harmonic mappings*, J. London Math. Soc. **42** (1990), 237–248.

FACULTAD DE MATEMÁTICAS, P. UNIVERSIDAD CATÓLICA DE CHILE, CASILLA 306, SANTIAGO 22, CHILE

E-mail address: `mchuaqui@mat.puc.cl`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MICHIGAN, ANN ARBOR, MICHIGAN 48109–1043

E-mail address: `duren@umich.edu`

DEPARTMENT OF ELECTRICAL ENGINEERING, STANFORD UNIVERSITY, STANFORD, CALIFORNIA 94305

E-mail address: `osgood@ee.stanford.edu`