

DECOMPOSITION THEOREM AND RIESZ BASIS FOR AXISYMMETRIC POTENTIALS IN THE RIGHT HALF-PLANE.

SLAH CHAABI, STEPHANE RIGAT

ABSTRACT. The Weinstein equation with complex coefficients is the equation governing generalized axisymmetric potentials (GASP) which can be written as $L_m[u] = \Delta u + (m/x) \partial_x u = 0$, where $m \in \mathbb{C}$. We generalize results known for $m \in \mathbb{R}$ to $m \in \mathbb{C}$. We give explicit expressions of fundamental solutions for Weinstein operators and their estimates near singularities, then we prove a Green's formula for GASP in the right half-plane \mathbb{H}^+ for $\operatorname{Re} m < 1$. We establish a new decomposition theorem for the GASP in any annular domains for $m \in \mathbb{C}$, which is in fact a generalization of the Bôcher's decomposition theorem. In particular, using bipolar coordinates, we prove for annuli that a family of solutions for GASP equation in terms of associated Legendre functions of first and second kind is complete. For $m \in \mathbb{C}$, we show that this family is even a Riesz basis in some non-concentric circular annulus.

1. INTRODUCTION

In this article, we study the Weinstein differential operator

$$L_m = \Delta + \frac{m}{x} \frac{\partial}{\partial x}$$

with $m \in \mathbb{C}$, well-defined on the right half-plane $\mathbb{H}^+ = \{(x, y) \in \mathbb{R}^2, x > 0\} = \{z \in \mathbb{C}, \operatorname{Re} z > 0\}$. This class of operators is also called operators governing axisymmetric potentials, they have been studied quite intensively in cases $m \in \mathbb{N}$ or $m \in \mathbb{R}$ in [52, 51, 50, 53, 54, 55, 58, 57, 56, 60, 59, 61, 62, 63, 64, 47, 48, 49, 30, 12, 13, 14, 34, 35, 36, 18, 20, 19, 8, 9, 10, 7, 26, 27, 28, 29, 38]. In this paper, we focus exclusively on case $m \in \mathbb{C}$ and some results for integer values of m is recalled. The Weinstein equation is written

$$L_m u = 0. \tag{1.1}$$

Date: February 4, 2014.

2010 Mathematics Subject Classification. 35B07, 33E05, 35J15, 35E05, 42C15.

Key words and phrases. Axially symmetric solutions, Fundamental solutions, Riesz Basis, Elliptic functions and integrals.

Acknowledgements. Both of the authors wish to thank Laurent Baratchart and Alexander Borichev for very useful discussions and remarks on a preliminary version of this paper.

In the sequel, the sense in which the solutions are studied will be specified. We will also look at solutions to the equation in the sense of distributions

$$L_m u = \delta_{(x,y)},$$

where $\delta_{(x,y)}$ denotes the Dirac mass at $(x, y) \in \mathbb{H}^+$.

Here, we deliberately restrict to the case of dimension 2, but many results can be extended directly to the case of higher dimension, *i.e.* for the operators

$$\sum_{k=1}^n \partial_{x_k^2} + \frac{m}{x_n} \partial_{x_n}$$

on the half-space $\mathbb{H}_n^+ = \{x \in \mathbb{R}^n, x_n > 0\}$.

This is Weinstein who was the first to introduce this class of operators in 1948 in [50], he studied the case where $m \in \mathbb{N}^*$. In particular, he obtained the following mean-value formula for axisymmetric potentials which can be extended continuously to $\overline{\mathbb{H}}_+$

$$u(0,0) \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \sin^m \theta d\theta = \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} u(re^{i\theta}) \sin^m \theta d\theta \quad (PM)$$

and he first gave an expression of a fundamental solution in terms of Bessel functions and then in terms of elliptic integrals where the singular point is taken on the y -axis. He also established the link between the axisymmetric potentials for $m \in \mathbb{N}^*$ and the harmonic functions on \mathbb{R}^{m+2} , that we will recall in the proposition 2.4.

In [54, 55, 16], Weinstein and Diaz-Weinstein established the correspondence principle that we will recall between axisymmetric potentials corresponding to m and those corresponding to $2 - m$ (proposition 2.3). They deduced an expression of a fundamental solution (where the singular point is taken on the y -axis) for $m \in \mathbb{R}$ and they made a link between the Weinstein equation and Tricomi equations and their fundamental solutions.

In [35], Huber obtained a Poisson-type formula which generalizes (PM). He also dealt with the extensions of GASP in the whole complex plane. He was also interested in removal properties of singularities for GASP in [36], and he gave an abstract representation formula of nonnegative GASP that BreLOT generalized in [8, 7].

Moreover, Vekua gave means to express fundamental solutions of elliptic equations with analytic coefficients by using the Riemann functions, introduced in the past (see eg [25]) in the real hyperbolic context, he generalized to elliptic equations through the complex operators ∂_z et $\partial_{\bar{z}}$ in [45]. In heuristic words, in the same way that we can say a harmonic function is the real part of a holomorphic function, or the sum of a holomorphic and an anti-holomorphic function, Vekua expressed the fact that solutions of elliptic equations, and therefore especially GASP, are written as a sum of two functional, one applied to an arbitrary holomorphic function and functiun applied to an anti-holomorphic functiun also arbitrary. The functional write

can explicitly in terms of Riemann function, which are obtained by using the hypergeometric functions ([45]) or using fractional derivations ([12]). In [31], Henrici gave a very interesting introduction to the work of Vekua.

More recently, by using the work of Vekua in [41], Savina gave a series representation of fundamental solutions for the operator $\hat{L}u = \Delta u + a\partial_x u + b\partial_y u + cu$ and she studied the convergence of this series. She gave an application of the Helmholtz equation.

In [28], Gilbert studied the non-homogeneous Weinstein equation $m \geq 0$, he gave an integral formula for this class of equations, in particular, an explicit solution is given when the second member depends only of one variable.

Some Dirichlet problems can be found in [38] in [24] in special geometry ("geometry with separable variable").

Even if some results presented in this paper are known for real values of m , we make a totally self-contained presentation with elementary technics not usually used in the previous quoted papers. For instance, usual arguments involving estimates of hypergeometric integrals are replaced by arguments using Lebesgue dominated convergence theorem. The main result is a decomposition theorem for axisymmetric potentials which is new also for real values of m . We obtain a Liouville-type result for the solutions of Weinstein equation on \mathbb{H}^+ , the interesting side of this result is the fact that there is a loss of strict ellipticity of the Weinstein operator on the boundary of \mathbb{H}^+ . An application of the decomposition theorem is given by showing that an explicit family of axisymmetric potentials constructed with the introduction of bipolar coordinates is a Riesz basis in a some annulus.

2. NOTATIONS AND PRELIMINARIES

Throughout the following, $\mathbb{H}^+ = \{(x, y) \in \mathbb{R}^2, x > 0\}$ will denote the right half-plane. All scalar functions will be complex valued. If Ω is an open set of \mathbb{R}^n with $n \in \mathbb{N}^*$, $\mathcal{D}(\Omega)$ will designate the space of C^∞ functions compactly supported on Ω and the support of an arbitrary function f defined on Ω is $\text{supp } f := \overline{\{x \in \Omega, f(x) \neq 0\}}$.

Let K be a compact set of Ω , $\mathcal{D}_K(\Omega)$ is the set of functions $\varphi \in \mathcal{D}(\Omega)$ such that $\text{supp } \varphi \subset K$.

The partial derivatives of a differentiable function u on an open set $\Omega \subset \mathbb{R}^n$ will be denoted $\frac{\partial u}{\partial x_i}$ or $\partial_{x_i} u$, or sometimes u_{x_i} with $i \in \llbracket 1, n \rrbracket$ (for $a < b \in \mathbb{N}$, $\llbracket a, b \rrbracket$ denotes the set of all integers between a and b).

If $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{N}^n$ is a multi-index, we will denote

$$\partial^\alpha := \partial_{x_1}^{\alpha_1} \dots \partial_{x_n}^{\alpha_n} = \frac{\partial^{|\alpha|}}{\partial x_1^{\alpha_1} \dots \partial x_n^{\alpha_n}}$$

with $|\alpha| = \alpha_1 + \dots + \alpha_n$.

It is assumed that the reader is familiar with the terminology of distributions and we refer to [32].

Let L be a linear differential operator on Ω ,

$$L = \sum_{|\alpha| \leq N} a_\alpha \partial^\alpha$$

where $N \in \mathbb{N}$, the previous summation is performed on the multi-indices α of length $|\alpha| \leq N$, a_α are $C^\infty(\Omega)$ functions.

By definition, if T is a distribution, LT will be the distribution : $LT = \sum_{|\alpha| \leq N} a_\alpha \partial^\alpha T$.

L^* will designate the adjoint operator of L in the sense of distributions, namely if T is a distributions,

$$L^*T = \sum_{|\alpha| \leq N} (-1)^{|\alpha|} \partial^\alpha (a_\alpha T).$$

It is noticed that if $f, g \in \mathcal{D}(\Omega)$, we have

$$\langle Lf, g \rangle = \langle f, L^*g \rangle.$$

Let $a \in \Omega$ and L be a differential operator on Ω . A *fundamental solution of L on Ω at $a \in \Omega$* is a distribution T_a such that

$$LT_a = \delta_a,$$

where the previous equality is taken in the sense of distributions on Ω .

This equality can be rewritten

$$\forall \varphi \in \mathcal{D}(\Omega), \quad \varphi(a) = \langle LT_a, \varphi \rangle = \langle T_a, L^*\varphi \rangle.$$

In particular, if $a \in \Omega$ and if T_a is a fundamental solution of L^* at a on Ω and if $g \in \mathcal{D}(\Omega)$ is such that $g = L(\varphi)$ with $\varphi \in \mathcal{D}(\Omega)$, then

$$\forall a \in \Omega, \quad \varphi(a) = \langle T_a, g \rangle.$$

Indeed, we have

$$\forall a \in \Omega, \quad \varphi(a) = \langle \delta_a, \varphi \rangle = \langle L^*T_a, \varphi \rangle = \langle T_a, L\varphi \rangle = \langle T_a, g \rangle.$$

These fundamental solutions is therefore a good tool for solving $L\varphi = g$ on $\mathcal{D}(\Omega)$ if $g \in \mathcal{D}(\Omega)$.

If $m \in \mathbb{N}^*$, the Laplacian in \mathbb{R}^m will be denoted Δ_m , or Δ when $m = 2$. For $m \in \mathbb{C}$, L_m denotes the *Weinstein operator* : $\forall (x, y) \in \mathbb{H}^+$,

$$L_m u(x, y) = \Delta u(x, y) + \frac{m}{x} \frac{\partial u}{\partial x}(x, y), \quad \text{where } u \in C^2(\mathbb{H}^+).$$

The following notation will be sometimes used : if $f(x, y) = (f_1(x, y), f_2(x, y))$ is a C^1 vector function on an open set of \mathbb{R}^2 and \mathbb{C}^2 valued, then

$$\operatorname{div}(f) := \frac{\partial f_1}{\partial x} + \frac{\partial f_2}{\partial y}.$$

Similarly, if $f : \mathbb{R}^2 \rightarrow \mathbb{C}$ is a C^1 scalar function on an open set of \mathbb{R}^2 and \mathbb{C} valued, then

$$\nabla f := \left(\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y} \right).$$

With these notations, the operator L_m can be written as follows : if $u \in C^2(\mathbb{H}^+)$, then

$$L_m u(x, y) = x^{-m} \operatorname{div}(x^m \nabla u)(x, y).$$

It is clear from the Schwarz rule that if u is a function defined on a connected open set of \mathbb{H}^+ such that $\operatorname{div}(\sigma \nabla u) = 0$ where $\sigma : \mathbb{H}^+ \rightarrow \mathbb{R}_*^+$ is a C^1 function, then there is a function v which satisfies the well-known generalized Cauchy-Riemann system of equations :

$$\begin{cases} \frac{\partial v}{\partial x} = -\sigma \frac{\partial u}{\partial y} \\ \frac{\partial v}{\partial y} = \sigma \frac{\partial u}{\partial x} \end{cases}$$

and v satisfies the conjugate equation $\operatorname{div}(\frac{1}{\sigma} \nabla v) = 0$ (see for exemple [6]). This observation justifies the fact that we call L_{-m} with $m \in \mathbb{C}$ the conjugate operator of L_m .

L_m^* denotes adjoint operator of L_m : for all $u \in C^2(\mathbb{H}^+)$ and for all $(x, y) \in \mathbb{H}^+$,

$$L_m^* u(x, y) = \Delta u(x, y) - \frac{\partial}{\partial x} \left(\frac{m u(x, y)}{x} \right) = \Delta u(x, y) - \frac{m}{x} \frac{\partial u}{\partial x}(x, y) + \frac{m}{x^2} u(x, y)$$

This definition is given on \mathbb{H}^+ but it is easily transposed to the case of an open set Ω of \mathbb{H}^+ .

In the case where the functions involved do not depend only of x and y , we will write $L_{m,x,y}$ instead of L_m , which means that the partial derivatives are related to the variables x and y , and all other variables are considered to be fixed.

If $u \in \mathcal{D}(\mathbb{H}^+)$, we define $S_m u \in \mathcal{D}(\mathbb{H}^+)$ by

$$(S_m u)(x, y) = x^{-m} u(x, y).$$

If $u \in \mathcal{D}(\mathbb{H}^+)$, we define $Du \in \mathcal{D}(\mathbb{H}^+)$ by

$$(Du)(x, y) = \frac{\partial u}{\partial x}(x, y).$$

These operators satisfy the following proposition :

Proposition 2.1. S_m conjugates L_m^* and L_m , D conjugates L_{-m}^* and L_m , which means that

$$S_m L_m^* = L_m S_m, \quad L_{-m}^* D = D L_m.$$

Proof. Straightforward computations. □

Remark 2.2. (1) Let $m \in \mathbb{C}$, S_m and $L_m S_m$ are auto-adjoints operators, ie. $S_m = S_m^*$ and $L_m S_m = (L_m S_m)^*$.

- (2) *There is a result, which generalizes the first point of this remark about the conjugation of operators L_m and L_m^* .*

Let $\sigma : \Omega \rightarrow \mathbb{C}$ be a C^1 function which does not vanish, the operator defined on $C^2(\Omega)$ by : for $u \in C^2(\Omega)$,

$$P_\sigma u(x, y) = \frac{1}{\sigma(x, y)} \operatorname{div} (\sigma(x, y) \nabla u(x, y)),$$

where Ω is an open set of \mathbb{R}^2 .

Then

$$P_\sigma^* = \operatorname{div} \left(\sigma \nabla \left(\frac{\cdot}{\sigma} \right) \right).$$

Indeed, if $u, v \in \mathcal{D}(\Omega)$, then we have by using the derivation in the sense of distributions,

$$\begin{aligned} \langle P_\sigma u, v \rangle &= \int_{\Omega} \frac{1}{\sigma(x, y)} \operatorname{div} (\sigma(x, y) \nabla u(x, y)) v(x, y) \, dx dy \\ &= - \int_{\Omega} \sigma \nabla u \cdot \nabla \left(\frac{v}{\sigma} \right) \, dx dy \\ &= \int_{\Omega} u \operatorname{div} \left(\sigma \nabla \left(\frac{v}{\sigma} \right) \right) \\ &= \langle u, P_\sigma^* v \rangle \end{aligned}$$

We define S_σ the operator such that for $u \in C^2(\Omega)$,

$$(S_\sigma u)(x, y) = \frac{1}{\sigma(x, y)} u(x, y).$$

Thus, S_σ conjugates P_σ and P_σ^ , where $P_\sigma^* = \operatorname{div} (\sigma \nabla (\frac{\cdot}{\sigma}))$ because, in an obviously way, we have $S_\sigma P_\sigma^* = P_\sigma S_\sigma$.*

If m is a positive integer, we introduce the operator $T_m : u \mapsto v$ defined as follows :

for a function u defined on an open set Ω of \mathbb{H}^+ , the function v is defined on $\{x \in \mathbb{R}^{m+2}, (\sqrt{x_1^2 + \dots + x_{m+1}^2}, x_{m+2}) \in \Omega\}$ by

$$v(x_1, \dots, x_{m+2}) = u(\sqrt{x_1^2 + \dots + x_{m+1}^2}, x_{m+2}).$$

The two following propositions can be found in Weinstein work ([55]) and will be useful in the sequel (the (short) proofs are just direct computations):

Proposition 2.3. (Weinstein principle [55]) *Let Ω be a relatively compact open set of \mathbb{H}^+ , if $u : \Omega \rightarrow \mathbb{C}$ is C^2 , then for all $m \in \mathbb{C}$,*

$$L_m u = x^{1-m} L_{2-m} [x^{m-1} u].$$

Proposition 2.4. ([50]) *Let Ω be a relatively compact open set of \mathbb{H}^+ . For $u \in C^2(\Omega)$ and if $m \in \mathbb{N}$, then $\Delta_{m+2}(T_m u) = T_m(L_m u)$.*

The two previous propositions will allow us to calculate fundamental solutions for L_m and L_m^* with $m \in \mathbb{N}$ in a first step, and thereafter, for $m \in \mathbb{Z}$. Finally, estimates of these expressions will show that the expressions obtained actually provide fundamental solutions of L_m and L_m^* with $m \in \mathbb{C}$.

3. INTEGRAL EXPRESSIONS OF FUNDAMENTAL SOLUTIONS FOR INTEGER VALUES OF m .

We recall the definition of the Dirac mass in a point : if $(x, y) \in \mathbb{R}^2$, $\delta_{(x,y)}$ is the distribution defined by

$$\forall \varphi \in \mathcal{D}(\mathbb{R}^2), \quad \langle \delta_{(x,y)}, \varphi \rangle = \varphi(x, y).$$

Let m be a positive integer.

Proposition 3.1. (partially in [16, 49, 50]) *Let $m \in \mathbb{N}^*$. For $(x, y) \in \mathbb{H}^+$ and $(\xi, \eta) \in \mathbb{H}^+$,*

$$E_m(x, y, \xi, \eta) = -\frac{\xi^m}{2\pi} \int_{\theta=0}^{\pi} \frac{\sin^{m-1} \theta d\theta}{[(x - \xi)^2 + 4x\xi \sin^2(\frac{\theta}{2}) + (y - \eta)^2]^{m/2}}$$

is a fundamental solution on \mathbb{H}^+ for the operator $L_{m,\xi,\eta}^$ at the fixed point $(x, y) \in \mathbb{H}^+$, which means that in the sense of distributions, we have \mathbb{H}^+ :*

$$L_{m,\xi,\eta}^* E_m(x, y, \xi, \eta) = \delta_{(x,y)}(\xi, \eta).$$

Moreover, if $(\xi, \eta) \in \mathbb{H}^+$ is fixed, then in the sense of distributions on \mathbb{H}^+ , we have

$$L_{m,x,y} E_m(x, y, \xi, \eta) = \delta_{(\xi,\eta)}(x, y),$$

which means that E_m is a fundamental solution on \mathbb{H}^+ of the operator $L_{m,x,y}$ at the fixed point $(\xi, \eta) \in \mathbb{H}^+$.

Proof. Let $m \in \mathbb{N}^*$. We recall that

$$E(x) = -\frac{1}{m \omega_{m+2} \|x\|^m}, \quad x \in \mathbb{R}^{m+2},$$

is a fundamental solution for the Laplacian on \mathbb{R}^{m+2} i. e. that in the sense of distributions, $\Delta_{m+2} E = \delta_0$, where ω_{m+2} is the area of the unit sphere \mathbb{R}^{m+2} . Thus, for all $v \in \mathcal{D}(\mathbb{R}^{m+2})$,

$$v(t_1, \dots, t_{m+2}) = -\frac{1}{m \omega_{m+2}} \int_{\tau \in \mathbb{R}^{m+2}} \Delta_{m+2} v(\tau) \frac{d\tau_1 d\tau_2 \dots d\tau_{m+2}}{((\tau_1 - t_1)^2 + \dots + (\tau_{m+2} - t_{m+2})^2)^{m/2}}$$

where $\tau = (\tau_1, \dots, \tau_{m+2})$.

Applying this relation to the function $v = T_m u$ where $u \in \mathcal{D}(\mathbb{H}^+)$ and due to the proposition 2.4, we have for all $(x, y) \in \mathbb{H}^+$,

$$u(x, y) = -\frac{1}{m \omega_{m+2}} \int_{\mathbb{R}^{m+2}} \frac{(L_m u)(\sqrt{\xi_1^2 + \dots + x^2}, \xi_{m+2}) d\xi_1 \dots d\xi_{m+2}}{((\xi_1 - x)^2 + \xi_2^2 + \dots + \xi_{m+1}^2 + (\xi_{m+2} - y)^2)^{m/2}}$$

We will simplify this integral expression. For this, we will consider the following hyper-spherical coordinates :

$$\begin{aligned}\xi_1 &= \xi \cos \theta_1 \\ \xi_2 &= \xi \sin \theta_1 \cos \theta_2 \\ &\vdots \\ \xi_{m-1} &= \xi \sin \theta_1 \cdots \sin \theta_{m-2} \cos \theta_{m-1} \\ \xi_m &= \xi \sin \theta_1 \cdots \sin \theta_{m-1} \cos \theta_m \\ \xi_{m+1} &= \xi \sin \theta_1 \cdots \sin \theta_m\end{aligned}$$

where $\xi^2 = \xi_1^2 + \cdots + \xi_{m+1}^2$, $\theta_m \in]-\pi, \pi[$ and $\theta_1, \dots, \theta_{m-1} \in]0, \pi[$. The absolute value of the determinant of the Jacobian matrix defined by this system of coordinates is

$$\xi^m \sin \theta_{m-1} \sin^2 \theta_{m-2} \cdots \sin^{m-1} \theta_1$$

Then we have for all $(x, y) \in \mathbb{H}^+$,

$$u(x, y) = \int_{\eta=-\infty}^{\infty} \int_{\xi=0}^{\infty} L_m(u)(\xi, \eta) E_m(x, y, \xi, \eta) d\xi d\eta \quad (3.1)$$

with

$$E_m(x, y, \xi, \eta) = -\frac{\xi^m}{m \omega_{m+2}} \int_{\theta_m=-\pi}^{\pi} \int_{\theta_1, \dots, \theta_{m-1}=0}^{\pi} \frac{\sin \theta_{m-1} \sin^2 \theta_{m-2} \cdots \sin^{m-1} \theta_1 d\theta_1 \cdots d\theta_m}{(\xi^2 - 2x\xi \cos \theta_1 + x^2 + (y - \eta)^2)^{m/2}}$$

Since $I := \int_{\theta_m=-\pi}^{\pi} \int_{\theta_2, \dots, \theta_{m-1}=0}^{\pi} \sin \theta_{m-1} \sin^2 \theta_{m-2} \cdots \sin^{m-2} \theta_2 d\theta_2 \cdots d\theta_{m-1} d\theta_m$ is the area of the unit sphere on \mathbb{R}^m because

$$\omega_m = \int_{\mathbb{S}_m} 1 d\sigma = \int_{\theta_{m-1}=-\pi}^{\pi} \int_{\theta_1, \dots, \theta_{m-2}=0}^{\pi} \sin \theta_{m-2} \sin^2 \theta_{m-3} \cdots \sin^{m-2} \theta_1 d\theta_2 \cdots d\theta_{m-1},$$

E_m can be written as :

$$E_m(x, y, \xi, \eta) = -\frac{\omega_m \xi^m}{m \omega_{m+2}} \int_{\theta=0}^{\pi} \frac{\sin^{m-1} \theta d\theta}{(\xi^2 - 2x\xi \cos \theta + x^2 + (y - \eta)^2)^{m/2}}.$$

Using the fact that $\omega_m = \frac{2\pi^{\frac{m}{2}}}{\Gamma(\frac{m}{2})}$, we get

$$E_m(x, y, \xi, \eta) = -\frac{\xi^m}{2\pi} \int_{\theta=0}^{\pi} \frac{\sin^{m-1} \theta d\theta}{((x - \xi)^2 + 4x\xi \sin^2(\frac{\theta}{2}) + (y - \eta)^2)^{m/2}}$$

and we have proved moreover thanks to (3.1) that

$$L_{m, \xi, \eta}^* E_m(x, y\xi, \eta) = \delta_{(x, y)}(\xi, \eta).$$

Moreover, since for all $(x, y) \in \mathbb{H}^+$ and for all $(\xi, \eta) \in \mathbb{H}^+$, we have

$$E_m(x, y, \xi, \eta) = \left(\frac{x}{\xi}\right)^{-m} E_m(\xi, \eta, x, y)$$

and thanks to the Proposition 2.1, S_m conjugates L_m^* and L_m , we have in the sense of distributions

$$L_{m,x,y}E_m(x,y,\xi,\eta) = L_{m,x,y}\left(\left(\frac{x}{\xi}\right)^{-m}E_m(\xi,\eta,x,y)\right) = \left(\frac{x}{\xi}\right)^{-m}L_{m,x,y}^*E_m(\xi,\eta,x,y),$$

then

$$L_{m,x,y}E_m(x,y,\xi,\eta) = \left(\frac{x}{\xi}\right)^{-m}\delta_{(\xi,\eta)}(x,y) = \delta_{(\xi,\eta)},$$

and this completes the proof. \square

For $m \in \mathbb{Z} \setminus \mathbb{N}$, the previous proposition and the Weinstein principle gives us the following proposition :

Proposition 3.2. (partially in [16, 49, 50]) *Let $m \in \mathbb{Z} \setminus \mathbb{N}^*$. For $(x,y) \in \mathbb{H}^+$ and $(\xi,\eta) \in \mathbb{H}^+$,*

$$\begin{aligned} E_m(x,y,\xi,\eta) &= \left(\frac{\xi}{x}\right)^{m-1}E_{2-m}(x,y,\xi,\eta) \\ &= -\frac{\xi x^{1-m}}{2\pi} \int_{\theta=0}^{\pi} \frac{\sin^{1-m}\theta d\theta}{[(x-\xi)^2 + 4x\xi \sin^2(\frac{\theta}{2}) + (y-\eta)^2]^{1-\frac{m}{2}}} \end{aligned}$$

is a fundamental solution on \mathbb{H}^+ for the operator $L_{m,\xi,\eta}^$ at the fixed point $(x,y) \in \mathbb{H}^+$ and it is also a fundamental solution on \mathbb{H}^+ of the operator $L_{m,x,y}$ at the fixed point $(\xi,\eta) \in \mathbb{H}^+$.*

Proof. We have for all $m \in \mathbb{N}^*$, $u \in \mathcal{D}(\mathbb{H}^+)$ and $(x,y) \in \mathbb{H}^+$,

$$u(x,y) = \int_{(\xi,\eta) \in \mathbb{H}^+} (L_m u) E_m(x,y,\xi,\eta) d\xi d\eta,$$

and by the Weinstein principle (proposition 2.3), we have

$$u(x,y) = \int_{\mathbb{H}^+} \xi^{1-m} L_{2-m}(\xi^{m-1}u) E_m(x,y,\xi,\eta) d\xi d\eta.$$

Denoting $v(x,y) = x^{m-1}u(x,y)$, we obtain

$$x^{1-m}v(x,y) = \int_{\mathbb{H}^+} \xi^{1-m} (L_{2-m}v) E_m(x,y,\xi,\eta) d\xi d\eta,$$

then, for all $m' \in \mathbb{Z} \setminus \mathbb{N}^*$, $v \in \mathcal{D}(\mathbb{H}^+)$ and $(x,y) \in \mathbb{H}^+$, putting $m = 2 - m'$, we have

$$v(x,y) = \int_{\mathbb{H}^+} (L_{m'}v) \left(\frac{\xi}{x}\right)^{m'-1} E_{2-m'}(x,y,\xi,\eta) d\xi d\eta.$$

The proof of the second point is totally similar. \square

4. FUNDAMENTAL SOLUTIONS FOR THE WEINSTEIN EQUATION WITH COMPLEX COEFFICIENTS

In this section, we will generalize the result obtained in the previous section for $m \in \mathbb{Z}$ to $m \in \mathbb{C}$.

More precisely, if $\operatorname{Re} m \geq 1$, then

$$E_m = -\frac{\xi^m}{2\pi} \int_{\theta=0}^{\pi} \frac{\sin^{m-1} \theta d\theta}{[(x-\xi)^2 + 4x\xi \sin^2(\frac{\theta}{2}) + (y-\eta)^2]^{m/2}}$$

is suitable, and if $\operatorname{Re} m < 1$, then

$$E_m = -\frac{\xi x^{1-m}}{2\pi} \int_{\theta=0}^{\pi} \frac{\sin^{1-m} \theta d\theta}{[(x-\xi)^2 + 4x\xi \sin^2(\frac{\theta}{2}) + (y-\eta)^2]^{1-\frac{m}{2}}}$$

is suitable.

In the sequel, E_m will always designate the corresponding formula (depending of $\operatorname{Re} m \geq 1$ or $\operatorname{Re} m < 1$).

Proposition 4.1. *For $m \in \mathbb{C}$ and $(\xi, \eta) \in \mathbb{H}^+$ fixed, we have*

$$\forall (x, y) \in \mathbb{H}^+ \setminus \{(\xi, \eta)\} \quad L_{m,x,y} E_m(x, y, \xi, \eta) = 0.$$

and for $(x, y) \in \mathbb{H}^+$ fixed, we have

$$\forall (\xi, \eta) \in \mathbb{H}^+ \setminus \{(x, y)\} \quad L_{m,\xi,\eta}^* E_m(x, y, \xi, \eta) = 0.$$

Proof. For convenience in the calculations, it should be denoted

$$f_m(x, y, \xi, \eta, \theta) = \frac{1}{[(x-\xi)^2 + 4x\xi \sin^2(\frac{\theta}{2}) + (y-\eta)^2]^{\frac{m}{2}}}.$$

To prove the first equality of the proposition, it suffices to show that

$$\int_{\theta=0}^{\pi} L_{m,x,y} f_m(x, y, \xi, \eta, \theta) \sin^{m-1} \theta d\theta = 0.$$

Let's compute the derivatives of the function f_m :

$$\partial_x f_m = \frac{-m}{2} \frac{2(x-\xi) + 4\xi \sin^2 \frac{\theta}{2}}{[(x-\xi)^2 + 4x\xi \sin^2(\frac{\theta}{2}) + (y-\eta)^2]^{\frac{m}{2}+1}} \quad (= -m(x-\xi \cos \theta) f_{m+2})$$

and

$$\begin{aligned} \partial_{xx} f_m &= \frac{-m}{[(x-\xi)^2 + 4x\xi \sin^2(\frac{\theta}{2}) + (y-\eta)^2]^{\frac{m}{2}+1}} + \\ &\quad + \frac{m}{2} \left(\frac{m}{2} + 1 \right) \frac{(2(x-\xi) + 4\xi \sin^2 \frac{\theta}{2})^2}{[(x-\xi)^2 + 4x\xi \sin^2(\frac{\theta}{2}) + (y-\eta)^2]^{\frac{m}{2}+2}} \end{aligned}$$

and

$$\partial_{yy} f_m = \frac{-m}{[(x-\xi)^2 + 4x\xi \sin^2(\frac{\theta}{2}) + (y-\eta)^2]^{\frac{m}{2}+1}} +$$

$$+\frac{m}{2}\left(\frac{m}{2}+1\right)\frac{(2(y-\eta))^2}{[(x-\xi)^2+4x\xi\sin^2\left(\frac{\theta}{2}\right)+(y-\eta)^2]^{\frac{m}{2}+2}}$$

We then have

$$\begin{aligned}\Delta f_m &= \frac{-2m}{[(x-\xi)^2+4x\xi\sin^2\left(\frac{\theta}{2}\right)+(y-\eta)^2]^{\frac{m}{2}+1}}+ \\ &+\frac{m}{2}\left(\frac{m}{2}+1\right)\frac{(2(x-\xi)+4\xi\sin^2\frac{\theta}{2})^2+(2(y-\eta))^2}{[(x-\xi)^2+4x\xi\sin^2\left(\frac{\theta}{2}\right)+(y-\eta)^2]^{\frac{m}{2}+2}}.\end{aligned}$$

However

$$\left(2(x-\xi)+4\xi\sin^2\frac{\theta}{2}\right)^2+(2(y-\eta))^2=4\left[(x-\xi)^2+4x\xi\sin^2\left(\frac{\theta}{2}\right)+(y-\eta)^2\right]-4\xi^2\sin^2\theta$$

then

$$\begin{aligned}\Delta f_m &= \frac{m^2}{[(x-\xi)^2+4x\xi\sin^2\left(\frac{\theta}{2}\right)+(y-\eta)^2]^{\frac{m}{2}+1}} \\ &\quad -\frac{m(m+2)\xi^2\sin^2\theta}{[(x-\xi)^2+4x\xi\sin^2\left(\frac{\theta}{2}\right)+(y-\eta)^2]^{\frac{m}{2}+2}}.\end{aligned}$$

Noting that

$$\frac{\partial f_{m+2}}{\partial\theta}=-\frac{(m+2)x\xi\sin\theta}{[(x-\xi)^2+4x\xi\sin^2\left(\frac{\theta}{2}\right)+(y-\eta)^2]^{\frac{m}{2}+2}},$$

we have

$$\Delta f_m=m^2 f_{m+2}+m\frac{\xi}{x}\sin\theta\frac{\partial f_{m+2}}{\partial\theta}$$

and by integration by parts, we have :

$$\begin{aligned}\int_{\theta=0}^{\pi}\Delta f_m\sin^{m-1}\theta d\theta &=m^2\int_{\theta=0}^{\pi}f_{m+2}\sin^{m-1}\theta d\theta+m\frac{\xi}{x}\int_{\theta=0}^{\pi}\frac{\partial f_{m+2}}{\partial\theta}\sin^m\theta d\theta \\ &=\frac{m}{x}\int_{\theta=0}^{\pi}m(x-\xi\cos\theta)f_{m+2}\sin^{m-1}\theta d\theta \\ &=-\frac{m}{x}\int_{\theta=0}^{\pi}\partial_x f_m\sin^{m-1}\theta d\theta,\end{aligned}$$

and the result is deduced in the case $\operatorname{Re} m \geq 1$. The proof is totally similar if $\operatorname{Re} m < 1$. The second equality of the proposition can be deduced immediately of the fact that S_m conjugates L_m^* and L_m (see proposition 2.1). \square

In the sequel, we will denote

$$d^2=(x-\xi)^2+(y-\eta)^2 \text{ and } k=\frac{4x\xi}{d^2}.$$

The following proposition gives the behavior of these functions near their singularity. And it will be useful to show that they are indeed fundamental solutions for $m \in \mathbb{C}$ and not only for integer values of m . In particular, we show that the behavior of the fundamental solutions is close to the behavior

of fundamental solutions for the Laplacian. This fact is well known for elliptic operators. But we emphasize here that in the proof of this proposition, the estimates of elliptic integrals are totally elementary estimates (using the dominated convergence theorem) and here we do not use estimates arising from classical estimates of hypergeometric functions. From those integral expressions, we deduce the following estimations :

Proposition 4.2. *Let $m \in \mathbb{C}$. For $(x, y) \in \mathbb{H}^+$ fixed,*

$$E_m(x, y, \xi, \eta) \underset{(\xi, \eta) \rightarrow (x, y)}{\sim} \frac{1}{2\pi} \ln \sqrt{(x - \xi)^2 + (y - \eta)^2}$$

Proof. We start with $\operatorname{Re} m \geq 1$.

In this case, we have :

$$\begin{aligned} E_m(x, y, \xi, \eta) &= -\frac{\xi^m}{2\pi} \int_{\theta=0}^{\pi} \frac{\sin^{m-1} \theta d\theta}{[(x - \xi)^2 + 4x\xi \sin^2(\frac{\theta}{2}) + (y - \eta)^2]^{m/2}} \\ &= -\frac{1}{2\pi} \left(\frac{\xi}{d}\right)^m \int_{\theta=0}^{\pi} \frac{\sin^{m-1} \theta d\theta}{(1 + k \sin^2 \frac{\theta}{2})^{m/2}}. \end{aligned}$$

Note that when $d \rightarrow 0$, $k \rightarrow +\infty$.

We have the following proposition :

Proposition 4.3. *When $k \rightarrow +\infty$ and $m \in \mathbb{C}$,*

$$\int_{\theta=0}^{\pi} \frac{\sin^{m-1} \theta d\theta}{(1 + k \sin^2 \frac{\theta}{2})^{m/2}} \underset{k \rightarrow +\infty}{\sim} \frac{2^{m-1}}{k^{m/2}} \ln k.$$

Proof. Putting $u = \sin \frac{\theta}{2}$, this integral is equal to

$$2^m \int_0^1 \frac{u^{m-1} (1 - u^2)^{\frac{m-2}{2}} du}{(1 + ku^2)^{m/2}} = \frac{2^m}{k^{m/2}} \int_0^1 \frac{u^{m-1} (1 - u^2)^{\frac{m-2}{2}} du}{(\frac{1}{k} + u^2)^{m/2}}.$$

However

$$\int_0^1 \frac{u^{m-1} (1 - u^2)^{\frac{m-2}{2}} du}{(\frac{1}{k} + u^2)^{m/2}} - \int_0^1 \frac{u^{m-1} du}{(\frac{1}{k} + u^2)^{m/2}} = - \int_0^1 \frac{u^{m-1}}{(\frac{1}{k} + u^2)^{m/2}} (1 - (1 - u^2)^{\frac{m-2}{2}}) du$$

and by monotone convergence, we obtain

$$\xrightarrow{k \rightarrow +\infty} - \int_0^1 \frac{u^{m-1}}{(u^2)^{m/2}} (1 - (1 - u^2)^{\frac{m-2}{2}}) du = - \int_0^1 \frac{1 - (1 - u^2)^{\frac{m-2}{2}}}{u} du$$

The change of variable $u = \frac{1}{\sqrt{k}} \operatorname{sh} t$ gives us

$$\int_0^1 \frac{u^{m-1} du}{(\frac{1}{k} + u^2)^{m/2}} = \int_0^{\operatorname{argsh} \sqrt{k}} \operatorname{th}^{m-1} t dt$$

Since $\operatorname{th}^{m-1} t$ tends to 1 when $t \rightarrow +\infty$, we deduce that when $k \rightarrow +\infty$

$$\int_0^{\operatorname{argsh} \sqrt{k}} \operatorname{th}^{m-1} dt \underset{k \rightarrow +\infty}{\sim} \int_0^{\operatorname{argsh} \sqrt{k}} dt = \operatorname{argsh} \sqrt{k} \underset{k \rightarrow +\infty}{\sim} \frac{1}{2} \ln k.$$

The proof is completed. \square

Due to Proposition 4.3, we have

$$E_m(x, y, \xi, \eta) \underset{d \rightarrow 0^+}{\sim} -\frac{1}{2\pi} \left(\frac{x}{d}\right)^m \frac{2^{m-1}}{k^{m/2}} \ln k \underset{d \rightarrow 0^+}{\sim} \frac{1}{2\pi} \ln d.$$

The case $\operatorname{Re} m < 1$ is analogous. \square

Now, we can prove the main result of this section,

Theorem 4.4. *Let $m \in \mathbb{C}$. For $(x, y) \in \mathbb{H}^+$ and $(\xi, \eta) \in \mathbb{H}^+$,*

$$E_m(x, y, \xi, \eta) = -\frac{\xi^m}{2\pi} \int_{\theta=0}^{\pi} \frac{\sin^{m-1} \theta d\theta}{[(x-\xi)^2 + 4x\xi \sin^2(\frac{\theta}{2}) + (y-\eta)^2]^{m/2}} \quad \text{if } \operatorname{Re} m \geq 1$$

$$\begin{aligned} \text{and } E_m(x, y, \xi, \eta) &= \left(\frac{\xi}{x}\right)^{m-1} E_{2-m}(x, y, \xi, \eta) \\ &= -\frac{\xi x^{1-m}}{2\pi} \int_{\theta=0}^{\pi} \frac{\sin^{1-m} \theta d\theta}{[(x-\xi)^2 + 4x\xi \sin^2(\frac{\theta}{2}) + (y-\eta)^2]^{1-\frac{m}{2}}} \quad \text{if } \operatorname{Re} m < 1 \end{aligned}$$

is a fundamental solution on \mathbb{H}^+ for $L_{m,\xi,\eta}^*$ at the fixed point $(x, y) \in \mathbb{H}^+$, which means that in the sense of distributions on \mathbb{H}^+ :

$$L_{m,\xi,\eta}^* E_m(x, y, \xi, \eta) = \delta_{(x,y)}(\xi, \eta).$$

Moreover, if $(\xi, \eta) \in \mathbb{H}^+$ is fixed, then in the sense of distributions on \mathbb{H}^+ :

$$L_{m,x,y} E_m(x, y, \xi, \eta) = \delta_{(\xi,\eta)}(x, y),$$

which means that E_m is a fundamental solution on \mathbb{H}^+ of $L_{m,x,y}$ at the fixed point $(\xi, \eta) \in \mathbb{H}^+$.

Proof. Let $m \in \mathbb{C}$ and $u \in \mathcal{D}(\mathbb{H}^+)$. Let $(x, y) \in \mathbb{H}^+$ and $\varepsilon > 0$ such that $D((x, y), \varepsilon) \subset \mathbb{H}^+$ where $D((x, y), \varepsilon)$ is the disk of center (x, y) and of radius ε .

We put

$$\begin{aligned} I_\varepsilon &:= \int_{\mathbb{H}^+ \setminus D((x,y),\varepsilon)} L_m(u)(\xi, \eta) E_m(x, y, \xi, \eta) d\xi d\eta = \\ &= \int_{\mathbb{H}^+ \setminus D((x,y),\varepsilon)} (L_m(u)(\xi, \eta) E_m(x, y, \xi, \eta) - u(\xi, \eta) L_m^*(E_m)(x, y, \xi, \eta)) d\xi d\eta \end{aligned}$$

because $L_m^*(E_m) = 0$ on $\mathbb{H}^+ \setminus D((x, y), \varepsilon)$. An elementary calculation gives us

$$\begin{aligned} L_m(u)E_m - uL_m^*(E) &= \partial_\xi \left((\partial_\xi u) E_m - u(\partial_\xi E_m) + \frac{m}{\xi} u E_m \right) \\ &\quad + \partial_\eta \left((\partial_\eta u) E_m - u(\partial_\eta E_m) \right). \end{aligned}$$

We will recall the Green formula in the framework that will be useful to us here.

Recall. Let Ω be an open set of \mathbb{R}^2 whose boundary is piecewise C^1 -differentiable.

By denoting \vec{n} the outer unit normal vector to $\partial\Omega$ and ds the arc length element on $\partial\Omega$ (positively oriented), if $X = (X_1, X_2) : \bar{\Omega} \rightarrow \mathbb{C}^2$ is a C^1 vector field, then

$$\int_{\Omega} \operatorname{div} X(x, y) dx dy = \int_{\partial\Omega} X(x, y) \cdot \vec{n}(x, y) ds$$

With this reminder applied to the open set $\Omega = U \setminus D((x, y), \varepsilon)$ where U is a regular open set of \mathbb{H}^+ containing the support of u , we have

$$I_{\varepsilon} = - \int_{\substack{t \in [0, 2\pi] \\ (\xi, \eta) = (x, y) + \varepsilon(\cos t, \sin t)}} \left(\left((\partial_{\xi} u) E_m - u(\partial_{\xi} E_m) + \frac{m}{\xi} u E_m \right) \cos t + \right. \\ \left. + ((\partial_{\eta} u) E_m - u(\partial_{\eta} E_m)) \sin t \right) \varepsilon dt$$

Proposition 4.2 shows that

$$\int_{\substack{t \in [0, 2\pi] \\ (\xi, \eta) = (x, y) + \varepsilon(\cos t, \sin t)}} \left[\left[(\partial_{\xi} u) + \frac{m}{\xi} u \right] \cos t + (\partial_{\eta} u) \sin t \right] E_m \varepsilon dt \xrightarrow{\varepsilon \rightarrow 0^+} 0$$

because $\lim_{\varepsilon \rightarrow 0} \varepsilon \ln \varepsilon = 0$. Then, if we want to prove that $\lim_{\varepsilon \rightarrow 0} I_{\varepsilon}$ exists, we have to prove the existence of

$$\lim_{\varepsilon \rightarrow 0} \int_{\substack{t \in [0, 2\pi] \\ (\xi, \eta) = (x, y) + \varepsilon(\cos t, \sin t)}} u \left((\partial_{\xi} E_m) \cos t + (\partial_{\eta} E_m) \sin t \right) \varepsilon dt,$$

and this limit will be equal to the limit of I_{ε} .

Now, we assume that $\operatorname{Re} m \geq 1$.

We denote J_{ε} the integral in the previous expression. A computation gives

$$J_{\varepsilon} = - \underbrace{\frac{m}{2\pi} \int_{\substack{t \in [0, 2\pi] \\ (\xi, \eta) = (x, y) + \varepsilon(\cos t, \sin t)}} u \frac{\xi^{m-1}}{\varepsilon^m} \int_0^{\pi} \frac{\sin^{m-1} \theta d\theta}{(1 + k \sin^2 \frac{\theta}{2})^{m/2}} \varepsilon \cos t dt}_{J_{\varepsilon,1}} + \\ + \underbrace{\frac{m}{2\pi} \int_{\substack{t \in [0, 2\pi] \\ (\xi, \eta) = (x, y) + \varepsilon(\cos t, \sin t)}} u \frac{\xi^m}{\varepsilon^{m+2}} \int_0^{\pi} \frac{\sin^{m-1} \theta d\theta}{(1 + k \sin^2 \frac{\theta}{2})^{m/2+1}} \varepsilon^2 dt}_{J_{\varepsilon,2}} + \\ + \underbrace{\frac{m}{2\pi} \int_{\substack{t \in [0, 2\pi] \\ (\xi, \eta) = (x, y) + \varepsilon(\cos t, \sin t)}} u \frac{\xi^m}{\varepsilon^{m+2}} \int_0^{\pi} \frac{2x \sin^2 \frac{\theta}{2} \sin^{m-1} \theta d\theta}{(1 + k \sin^2 \frac{\theta}{2})^{m/2+1}} \varepsilon \cos t dt}_{J_{\varepsilon,3}}$$

where $k = \frac{4x\xi}{\varepsilon^2}$.

We have the following propositions :

Proposition 4.5. *When $k \rightarrow +\infty$ and $m \in \mathbb{C}$*

$$\int_{\theta=0}^{\pi} \frac{\sin^2 \frac{\theta}{2} \sin^{m-1} \theta d\theta}{(1 + k \sin^2 \frac{\theta}{2})^{m/2+1}} \underset{k \rightarrow +\infty}{\sim} \frac{2^{m-1}}{k^{\frac{m}{2}+1}} \ln k.$$

Proof. We put $u = \sin \frac{\theta}{2}$, this integral is equal to

$$2^m \int_0^1 \frac{u^{m+1}(1-u^2)^{\frac{m-2}{2}} du}{(1+ku^2)^{m/2+1}} = \frac{2^m}{k^{m/2+1}} \int_0^1 \frac{u^{m+1}(1-u^2)^{\frac{m-2}{2}} du}{(\frac{1}{k} + u^2)^{m/2+1}}.$$

However

$$\begin{aligned} & \int_0^1 \frac{u^{m+1}(1-u^2)^{\frac{m-2}{2}} du}{(\frac{1}{k} + u^2)^{m/2+1}} - \int_0^1 \frac{u^{m+1} du}{(\frac{1}{k} + u^2)^{m/2+1}} = - \int_0^1 \frac{u^{m+1}}{(\frac{1}{k} + u^2)^{m/2+1}} (1 - (1-u^2)^{\frac{m-2}{2}}) du \\ & \xrightarrow{k \rightarrow +\infty} - \int_0^1 \frac{u^{m+1}}{(u^2)^{m/2+1}} (1 - (1-u^2)^{\frac{m-2}{2}}) du = - \int_0^1 \frac{1 - (1-u^2)^{\frac{m-2}{2}}}{u} du. \end{aligned}$$

The change of variable $u = \frac{1}{\sqrt{k}} \text{sh } t$ gives us

$$\int_0^1 \frac{u^{m+1} du}{(\frac{1}{k} + u^2)^{m/2+1}} = \int_0^{\text{argsh } \sqrt{k}} \text{th}^{m+1} t dt$$

Since $\text{th}^{m+1} t$ tends to 1 when $t \rightarrow +\infty$, it follows that when $k \rightarrow +\infty$

$$\int_0^{\text{argsh } \sqrt{k}} \text{th}^{m+1} t dt \underset{k \rightarrow +\infty}{\sim} \int_0^{\text{argsh } \sqrt{k}} dt = \text{argsh } \sqrt{k} \underset{k \rightarrow +\infty}{\sim} \frac{1}{2} \ln k.$$

The proposition is well proven.. □

Proposition 4.6. *When $k \rightarrow +\infty$ and $m \in \mathbb{C}$*

$$\int_{\theta=0}^{\pi} \frac{\sin^{m-1} \theta d\theta}{(1 + k \sin^2 \frac{\theta}{2})^{m/2+1}} \underset{k \rightarrow +\infty}{\sim} \frac{2^m}{mk^{\frac{m}{2}}}$$

Proof. Putting as previously $u = \sin \frac{\theta}{2}$, this integral is equal to

$$2^m \int_0^1 \frac{u^{m-1}(1-u^2)^{\frac{m-2}{2}} du}{(1+ku^2)^{m/2+1}} = \frac{2^m}{k^{m/2+1}} \int_0^1 \frac{u^{m-1}(1-u^2)^{\frac{m-2}{2}} du}{(\frac{1}{k} + u^2)^{m/2+1}}.$$

However

$$\int_0^1 \frac{u^{m-1}(1-u^2)^{\frac{m-2}{2}} du}{(\frac{1}{k} + u^2)^{m/2+1}} - \int_0^1 \frac{u^{m-1} du}{(\frac{1}{k} + u^2)^{m/2+1}} = - \int_0^1 \frac{u^{m-1}}{(\frac{1}{k} + u^2)^{m/2+1}} (1 - (1-u^2)^{\frac{m-2}{2}}) du$$

We first estimate the right hand side of this equality :

$$\int_0^1 \frac{u^{m-1}}{(\frac{1}{k} + u^2)^{m/2+1}} (1 - (1-u^2)^{\frac{m-2}{2}}) du - \int_0^1 \frac{u^{m-1}}{(\frac{1}{k} + u^2)^{m/2+1}} \left(\frac{m-2}{2} u^2 \right) du$$

$$\begin{aligned}
&= \int_0^1 \frac{u^{m-1}}{\left(\frac{1}{k} + u^2\right)^{m/2+1}} \left(1 - \frac{m-2}{2}u^2 - (1-u^2)^{\frac{m-2}{2}}\right) du \\
&\xrightarrow{k \rightarrow +\infty} \int_0^1 \frac{u^{m-1}}{(u^2)^{m/2+1}} \left(1 - \frac{m-2}{2}u^2 - (1-u^2)^{\frac{m-2}{2}}\right) du \\
&= \int_0^1 \frac{1 - \frac{m-2}{2}u^2 - (1-u^2)^{\frac{m-2}{2}}}{u^3} du. \tag{*}
\end{aligned}$$

As seen in the proof of Proposition 4.5, we have

$$\frac{m-2}{2} \int_0^1 \frac{u^{m+1}}{\left(\frac{1}{k} + u^2\right)^{\frac{m}{2}+1}} du \underset{k \rightarrow +\infty}{\sim} \frac{m-2}{4} \ln k. \tag{**}$$

Through (*) and (**), one obtains :

$$\int_0^1 \frac{u^{m-1}}{\left(\frac{1}{k} + u^2\right)^{m/2+1}} (1 - (1-u^2)^{\frac{m-2}{2}}) du \underset{k \rightarrow +\infty}{\sim} \frac{m-2}{4} \ln k.$$

The change of variable $u = \frac{1}{\sqrt{k}} \operatorname{sh} t$ gives us

$$\int_0^1 \frac{u^{m-1} du}{\left(\frac{1}{k} + u^2\right)^{m/2+1}} = k \int_0^{\operatorname{argsh} \sqrt{k}} \frac{\operatorname{th}^{m-1} t}{\operatorname{ch}^2 t} dt = \frac{k}{m} \operatorname{th}^m (\operatorname{argsh} \sqrt{k}).$$

It follows that when $k \rightarrow +\infty$,

$$\int_0^1 \frac{u^{m-1} du}{\left(\frac{1}{k} + u^2\right)^{m/2+1}} \underset{k \rightarrow +\infty}{\sim} \frac{k}{m}.$$

We thus obtain

$$\int_0^1 \frac{u^{m-1} (1-u^2)^{\frac{m-2}{2}} du}{\left(\frac{1}{k} + u^2\right)^{m/2+1}} \underset{k \rightarrow +\infty}{\sim} \frac{k}{m}.$$

And

$$\int_{\theta=0}^{\pi} \frac{\sin^{m-1} \theta d\theta}{\left(1 + k \sin^2 \frac{\theta}{2}\right)^{m/2+1}} \underset{k \rightarrow +\infty}{\sim} \frac{2^m}{mk^{\frac{m}{2}}}$$

and this completes the proof. \square

Let us return to the proof of Theorem 4.4.

The Proposition 4.3 shows that

$$\begin{aligned}
&J_{\varepsilon,1} \underset{\varepsilon \rightarrow 0+}{\sim} -\frac{m}{2\pi} \int_{(\xi,\eta)=(x,y)+\varepsilon(\cos t, \sin t)}^{t \in [0,2\pi]} u \frac{x^{m-1}}{\varepsilon^m} \frac{2^{m-1}}{k^{m/2}} (\ln k) \varepsilon \cos t dt \\
&\underset{\varepsilon \rightarrow 0+}{\sim} +\frac{m}{2\pi x} \varepsilon \ln \varepsilon \left(\int_{(\xi,\eta)=(x,y)+\varepsilon(\cos t, \sin t)}^{t \in [0,2\pi]} u(x + \varepsilon \cos t, y + \varepsilon \sin t) \cos t dt \right)
\end{aligned}$$

which tends to 0.

The Proposition 4.5 shows that

$$J_{\varepsilon,3} \underset{\varepsilon \rightarrow 0+}{\sim} \frac{m}{2\pi} \int_{(\xi,\eta)=(x,y)+\varepsilon(\cos t, \sin t)}^{t \in [0,2\pi]} u \frac{x^m}{\varepsilon^{m+2}} (2x) \frac{2^{m-1}}{k^{m/2+1}} (\ln k) \varepsilon \cos t dt$$

$$\underset{\varepsilon \rightarrow 0^+}{\sim} -\frac{m}{4\pi x} \varepsilon \ln \varepsilon \left(\int_{\substack{t \in [0, 2\pi] \\ (\xi, \eta) = (x, y) + \varepsilon(\cos t, \sin t)}} u(x + \varepsilon \cos t, y + \varepsilon \sin t) \cos t dt \right)$$

which tends to 0.

Finally, the proposition 4.6 shows that

$$\begin{aligned} J_{\varepsilon, 2} &\underset{\varepsilon \rightarrow 0^+}{\sim} \frac{m}{2\pi} \int_{\substack{t \in [0, 2\pi] \\ (\xi, \eta) = (x, y) + \varepsilon(\cos t, \sin t)}} u \frac{x^m}{\varepsilon^{m+2}} \frac{2^m}{mk^{m/2}} \varepsilon^2 dt \\ &\underset{\varepsilon \rightarrow 0^+}{\sim} \frac{1}{2\pi} \int_{\substack{t \in [0, 2\pi] \\ (\xi, \eta) = (x, y) + \varepsilon(\cos t, \sin t)}} u(x + \varepsilon \cos t, y + \varepsilon \sin t) dt \xrightarrow{\varepsilon \rightarrow 0^+} u(x, y). \end{aligned}$$

So we have proved that for all $m \in \mathbb{C}$ such that $\operatorname{Re} m > 0$,

$$\begin{aligned} \lim_{\varepsilon \rightarrow 0^+} \int_{\mathbb{H}^+ \setminus D((x, y), \varepsilon)} L_m(u)(\xi, \eta) E_m(x, y, \xi, \eta) d\xi d\eta &= \\ &= \int_{\mathbb{H}^+} L_m(u)(\xi, \eta) E_m(x, y, \xi, \eta) d\xi d\eta = u(x, y) \end{aligned}$$

therefore that E_m is indeed a fundamental solution of L_m^* for all $m \in \mathbb{C}$ with $\operatorname{Re} m > 0$.

Proof for $m \in \mathbb{C}$ with $\operatorname{Re} m \leq 1$ is completely similar.

We also have the dual assertions for fundamental solutions of L_m for all $m \in \mathbb{C}$ thanks to Proposition 2.1. \square

The following proposition is roughly a consequence of the previous theorem.

Proposition 4.7. *Let $m \in \mathbb{C}$ and let Ω be a relatively compact open set of \mathbb{H}^+ whose boundary is piecewise C^1 -differentiable.*

Then, for $(x, y) \in \Omega$ and $u \in C^2(\overline{\Omega})$, by denoting \vec{n} the outer unit normal vector to $\partial\Omega$ and ds the arc length element on $\partial\Omega$ (positively oriented), we have

$$\begin{aligned} u(x, y) &= \int_{\Omega} L_m(u) E_m d\xi d\eta \\ &\quad - \int_{\partial\Omega} \left[(\partial_\xi u) E_m - u(\partial_\xi E_m) + \frac{m}{\xi} u E_m, (\partial_\eta u) E_m - u(\partial_\eta E_m) \right] \cdot \vec{n} ds \end{aligned}$$

where $u := u(\xi, \eta)$ and $E_m := E_m(x, y, \xi, \eta)$.

Proof. Indeed, if $u \in C^2(\overline{\Omega})$, we have for $(x, y) \in \Omega$ and $\varepsilon > 0$ such that $\overline{D((x, y), \varepsilon)} \subset \Omega$:

$$\int_{\Omega \setminus D((x, y), \varepsilon)} L_m(u) E_m d\xi d\eta = \int_{\Omega \setminus D((x, y), \varepsilon)} (L_m(u) E_m - L_m^*(E_m) u) d\xi d\eta.$$

Thanks to the Green formula previously recalled, this last integral is equal to

$$\int_{\partial\Omega} \left[(\partial_\xi u) E_m - u(\partial_\xi E_m) + \frac{m}{\xi} u E_m, (\partial_\eta u) E_m - u(\partial_\eta E_m) \right] \cdot \vec{n} ds$$

$$- \int_{\substack{t \in [0, 2\pi] \\ (\xi, \eta) = (x, y) + \varepsilon(\cos t, \sin t)}} \left(\left((\partial_\xi u) E_m - u(\partial_\xi E_m) + \frac{m}{\xi} u E_m \right) \cos t + \right. \\ \left. + ((\partial_\eta u) E_m - u(\partial_\eta E_m)) \sin t \right) \varepsilon dt,$$

and from what we saw in the previous proof, this last expression tends to

$$\int_{\partial\Omega} \left[(\partial_\xi u) E_m - u(\partial_\xi E_m) + \frac{m}{\xi} u E_m, (\partial_\eta u) E_m - u(\partial_\eta E_m) \right] \cdot \vec{n} ds + u(x, y),$$

when $\varepsilon \rightarrow 0$.

The integrability nature of E_m near (x, y) shows that

$$\lim_{\varepsilon \rightarrow 0} \int_{\Omega \setminus D((x, y), \varepsilon)} L_m(u) E_m d\xi d\eta = \int_{\Omega} L_m(u) E_m d\xi d\eta,$$

and the proof of the proposition is complete. \square

5. LIOUVILLE-TYPE RESULT AND DECOMPOSITION THEOREM FOR THE AXISYMMETRIC POTENTIALS

In the previous section, we have just seen that there are two different expressions of the fundamental solutions depending on the values of m . For the rest, each of the expressions have different behaviors according to the value of m . We will look at the two cases separately : $\operatorname{Re} m < 1$ and $\operatorname{Re} m \geq 1$.

More specifically, we need fundamental solutions which vanish at the boundary of \mathbb{H}^+ , that is to say zero on the y -axis and zero to infinity.

For $\operatorname{Re} m < 1$, the formula

$$E_m(x, y, \xi, \eta) = -\frac{\xi x^{1-m}}{2\pi} \int_{\theta=0}^{\pi} \frac{\sin^{1-m} \theta d\theta}{[(x - \xi)^2 + 4x\xi \sin^2(\frac{\theta}{2}) + (y - \eta)^2]^{1-\frac{m}{2}}}$$

shows that E_m satisfies this property ($E_m(x, y, \cdot, \cdot)$ tends to 0 when $x \rightarrow 0+$ and $\|(x, y)\| \rightarrow +\infty$).

For $\operatorname{Re} m \geq 1$, the expression

$$E_m(x, y, \xi, \eta) = -\frac{\xi^m}{2\pi} \int_{\theta=0}^{\pi} \frac{\sin^{m-1} \theta d\theta}{[(x - \xi)^2 + 4x\xi \sin^2(\frac{\theta}{2}) + (y - \eta)^2]^{m/2}}$$

no longer satisfies this property. Contrariwise,

$$E_m(x, y, \xi, \eta) - E_m(-x, y, \xi, \eta)$$

is also a fundamental solution on \mathbb{H}^+ , and satisfies this property.

Then, we will put

- For $\operatorname{Re} m < 1$:

$$F_m(x, y, \xi, \eta) = E_m(x, y, \xi, \eta)$$

- For $\operatorname{Re} m \geq 1$:

$$F_m(x, y, \xi, \eta) = E_m(x, y, \xi, \eta) - E_m(-x, y, \xi, \eta).$$

We will need the following definition of convergence to the boundary of \mathbb{H}^+ .

Definition. Let $u : \mathbb{H}^+ \rightarrow \mathbb{R}$ be a function defined on \mathbb{H}^+ . We write

$$\lim_{\partial\mathbb{H}^+} u = 0$$

if and only if

$$\forall \varepsilon > 0, \quad \exists N \in \mathbb{N}, \quad \forall n \geq N, \quad \forall (x, y) \in \mathbb{H}^+, \\ x \leq \frac{1}{n} \text{ or } \|(x, y)\| \geq n \quad \implies \quad |u(x, y)| \leq \varepsilon.$$

In other words, this amounts to considering that the boundary $\partial\mathbb{H}^+$ of \mathbb{H}^+ consists of y -axis points and points at infinity and to say that the concept of punctual convergence to the boundary of \mathbb{H}^+ is a uniform convergence. Indeed, we do not need the uniform convergence. More precisely, we have the following proposition :

Proposition 5.1. *Let $u : \mathbb{H}^+ \rightarrow \mathbb{C}$. We have*

$$\lim_{\partial\mathbb{H}^+} u = 0$$

if and only if

$$\lim_{\|(x,y)\| \rightarrow +\infty} u(x, y) = 0 \quad \text{and} \quad \forall y \in \mathbb{R}, \quad \lim_{(0,y)} u = 0.$$

Proof. The direct implication is easy. Conversely, we assume

$$\lim_{\|(x,y)\| \rightarrow +\infty} u(x, y) = 0 \quad \text{and} \quad \forall y \in \mathbb{R}, \quad \lim_{(0,y)} u = 0$$

and we have to show $\lim_{\partial\mathbb{H}^+} u = 0$.

Let $\varepsilon > 0$. There is $A > 0$ such that for all $(\xi, \eta) \in \mathbb{H}^+$,

$$\sqrt{\xi^2 + \eta^2} \geq A \quad \implies \quad |u(\xi, \eta)| \leq \varepsilon.$$

Similarly, for all $y \in \mathbb{R}$, there is $\alpha_y \in]0, 1[$ such that for all $(\xi, \eta) \in \mathbb{H}^+$

$$\sqrt{\xi^2 + (\eta - y)^2} < \alpha_y \quad \implies \quad |u(\xi, \eta)| \leq \varepsilon.$$

The interval $[-A, A]$ is compact.

By the Lebesgue covering lemma, there is $\alpha > 0$ such that for all $y' \in [-A, A]$, the ball $B(y', \alpha)$ is included in one of the balls $B(y, \alpha_y)$ with $y \in [-A, A]$.

In particular, if $(\xi, \eta) \in \mathbb{H}^+$ is such that $0 < \xi < \alpha$, then $|u(\xi, \eta)| \leq \varepsilon$. This completes the proof. \square


The following proposition is a Liouville-type result for the axisymmetric potentials in the right half-plane and this result is not immediate because there is the loss of strict ellipticity of the Weinstein operator on the y -axis. In [4] (see Theorem 7.1), we can found an interesting result on the description of a class of non-strictly elliptic equations with unbounded coefficients.

Proposition 5.2. *Let $u \in C^2(\mathbb{H}^+)$ such that $L_m u = 0$ and $\lim_{\partial\mathbb{H}^+} u = 0$. Then $u \equiv 0$ on \mathbb{H}^+ .*

Proof. For $(\xi, \eta) \in \mathbb{H}^+$ and $N \in \mathbb{N}^*$, we define

$$\phi_N(\xi, \eta) = \theta_1(N\xi)\theta_2\left(\frac{\xi}{N}\right)\theta_2\left(\frac{\eta}{N}\right)$$

where θ_1 and θ_2 are smooth functions on \mathbb{R} , valued on $[0, 1]$ and such that $\theta_1(t) = 1$ for $t \geq 1$, $\theta_1(t) = 0$ for $t \leq \frac{1}{2}$, $\theta_2(t) = 1$ for $t \in [-\frac{1}{2}, \frac{1}{2}]$ and $\theta_2(t) = 0$ for $t \in \mathbb{R} \setminus]-\frac{1}{2}, \frac{1}{2}[$. We assume that all derivatives of θ_1 and θ_2 vanish at $\{-1, -\frac{1}{2}, \frac{1}{2}, 1\}$.



If $u \in C^2(\mathbb{H}^+)$ satisfies $L_m u = 0$, then $u\phi_N \in C^2(\mathbb{H}^+)$ and is compactly supported on \mathbb{H}^+ . Throughout the following, we fix $(x, y) \in \mathbb{H}^+$. For N sufficiently large, thanks to Proposition 4.7 (true if E_m is replaced by F_m), we have


$$u(x, y) = u(x, y)\phi_N(x, y) = \int_{\mathbb{H}^+} L_m(u\phi_N)F_m d\xi d\eta$$

(because the function $L_m(u\phi_N)$ is identically zero in a neighborhood of the singularity of F_m), thus

$$\begin{aligned} u(x, y) &= \int_{\mathbb{H}^+} [L_m(u)\phi_N + uL_m(\phi_N) + 2\nabla u \cdot \nabla \phi_N]F_m d\xi d\eta \\ &= \int_{\mathbb{H}^+} u[L_m(\phi_N)F_m - 2\operatorname{div}(F_m \nabla \phi_N)]d\xi d\eta \\ &= \int_{D_1 \cup \dots \cup D_8} u[L_m(\phi_N)F_m - 2\operatorname{div}(F_m \nabla \phi_N)]d\xi d\eta \\ &= - \int_{D_1 \cup \dots \cup D_8} u[L_{-m}(\phi_N)F_m + 2\nabla F_m \cdot \nabla \phi_N]d\xi d\eta \end{aligned}$$

where D_1, \dots, D_8 are the following domains (which depend of N):

$$\begin{aligned} D_1 &= \left[\frac{1}{2N}, \frac{1}{N}\right] \times \left[-\frac{N}{2}, \frac{N}{2}\right], & D_2 &= \left[\frac{1}{N}, \frac{N}{2}\right] \times \left[\frac{N}{2}, N\right], \\ D_3 &= \left[\frac{N}{2}, N\right] \times \left[-\frac{N}{2}, \frac{N}{2}\right], & D_4 &= \left[\frac{1}{N}, \frac{N}{2}\right] \times \left[-N, -\frac{N}{2}\right], \\ D_5 &= \left[\frac{1}{2N}, \frac{1}{N}\right] \times \left[\frac{N}{2}, N\right], & D_6 &= \left[\frac{N}{2}, N\right] \times \left[\frac{N}{2}, N\right], \\ D_7 &= \left[\frac{N}{2}, N\right] \times \left[-N, -\frac{N}{2}\right] & \text{and } D_8 &= \left[\frac{1}{2N}, \frac{1}{N}\right] \times \left[-N, -\frac{N}{2}\right]. \end{aligned}$$


 Figure : Domains D_i

Since $\lim_{\partial\mathbb{H}^+} u = 0$, then

$$u_N := \sup_{(\xi, \eta) \in D_1 \cup \dots \cup D_8} |u(\xi, \eta)| \xrightarrow{N \rightarrow +\infty} 0.$$

We will estimate each integrals supported on D_1, \dots, D_8 . For this, we need the following lemmas which will give us estimates of each terms when N tends to infinity. We recall that, if $(u_N)_N$ and $(v_N)_N$ are complex sequences, $u_N = \mathcal{O}(v_N)$ means that there exists a constant M such that, for every N sufficiently large, $|u_N| \leq M|v_N|$; $u_N = o(v_N)$ means that for every $\varepsilon > 0$, for every N sufficiently large, $|u_N| \leq \varepsilon|v_N|$.

Lemma 5.3. *On D_1 , we have*

$$\sup \left| \frac{\partial \phi_N}{\partial \xi} \right| = \mathcal{O}(N) \quad \text{and} \quad \sup \left| \frac{\partial \phi_N}{\partial \eta} \right| = 0.$$

On $D_2 \cup D_4$, we have

$$\sup \left| \frac{\partial \phi_N}{\partial \xi} \right| = 0 \quad \text{and} \quad \sup \left| \frac{\partial \phi_N}{\partial \eta} \right| = \mathcal{O}\left(\frac{1}{N}\right).$$

On D_3 , we have

$$\sup \left| \frac{\partial \phi_N}{\partial \xi} \right| = \mathcal{O}\left(\frac{1}{N}\right) \quad \text{and} \quad \sup \left| \frac{\partial \phi_N}{\partial \eta} \right| = 0.$$

On $D_5 \cup D_8$, we have

$$\sup \left| \frac{\partial \phi_N}{\partial \xi} \right| = \mathcal{O}(N) \quad \text{and} \quad \sup \left| \frac{\partial \phi_N}{\partial \eta} \right| = \mathcal{O}\left(\frac{1}{N}\right).$$

On $D_6 \cup D_7$, we have

$$\sup \left| \frac{\partial \phi_N}{\partial \xi} \right| = \mathcal{O}\left(\frac{1}{N}\right) \quad \text{and} \quad \sup \left| \frac{\partial \phi_N}{\partial \eta} \right| = \mathcal{O}\left(\frac{1}{N}\right).$$

On $D_1 \cup D_5 \cup D_8$, we have

$$\sup |L_{-m}(\phi_N)| = \mathcal{O}(N^2).$$

On $D_2 \cup D_3 \cup D_4 \cup D_6 \cup D_7$, we have

$$\sup |L_{-m}(\phi_N)| = \mathcal{O}\left(\frac{1}{N^2}\right).$$

Proof. *For $(\xi, \eta) \in D_1$, $\phi_N(\xi, \eta) = \theta_1(N\xi)$ and thus

$$\begin{aligned} \frac{\partial \phi_N}{\partial \xi}(\xi, \eta) &= N\theta'_1(N\xi) \quad , \quad \frac{\partial \phi_N}{\partial \eta}(\xi, \eta) = 0, \\ L_{-m}\phi_N(\xi, \eta) &= N^2\theta''_1(N\xi) - \frac{mN}{\xi}\theta'_1(N\xi), \end{aligned}$$

which give us

$$\sup_{D_1} \left| \frac{\partial \phi_N}{\partial \xi} \right| = \mathcal{O}(N), \quad \sup_{D_1} \left| \frac{\partial \phi_N}{\partial \eta} \right| = 0, \quad \sup_{D_1} |L_{-m}(\phi_N)| = \mathcal{O}(N^2)$$

since the derivatives of θ_1 are bounded and for $(\xi, \eta) \in D_1$, one gets $\xi \geq \frac{1}{2N}$.

*For $(\xi, \eta) \in D_2$, $\phi_N(\xi, \eta) = \theta_2\left(\frac{\eta}{N}\right)$ and thus

$$\begin{aligned} \frac{\partial \phi_N}{\partial \xi}(\xi, \eta) &= 0 \quad , \quad \frac{\partial \phi_N}{\partial \eta}(\xi, \eta) = \frac{1}{N}\theta'_2\left(\frac{\eta}{N}\right), \\ L_{-m}\phi_N(\xi, \eta) &= \frac{1}{N^2}\theta''_2\left(\frac{\eta}{N}\right), \end{aligned}$$

which give us

$$\sup_{D_2} \left| \frac{\partial \phi_N}{\partial \xi} \right| = 0, \quad \sup_{D_2} \left| \frac{\partial \phi_N}{\partial \eta} \right| = \mathcal{O}\left(\frac{1}{N}\right), \quad \sup_{D_2} |L_{-m}(\phi_N)| = \mathcal{O}\left(\frac{1}{N^2}\right)$$

*So does same with D_4 .

* For $(\xi, \eta) \in D_3$, $\phi_N(\xi, \eta) = \theta_2\left(\frac{\xi}{N}\right)$ and thus

$$\begin{aligned} \frac{\partial \phi_N}{\partial \xi}(\xi, \eta) &= \frac{1}{N}\theta'_2\left(\frac{\xi}{N}\right) \quad , \quad \frac{\partial \phi_N}{\partial \eta}(\xi, \eta) = 0, \\ L_{-m}\phi_N(\xi, \eta) &= \frac{1}{N^2}\theta''_2\left(\frac{\xi}{N}\right) - \frac{1}{N}\frac{m}{\xi}\theta'_2\left(\frac{\xi}{N}\right), \end{aligned}$$

which give us

$$\sup_{D_3} \left| \frac{\partial \phi_N}{\partial \xi} \right| = \mathcal{O}\left(\frac{1}{N}\right), \quad \sup_{D_3} \left| \frac{\partial \phi_N}{\partial \eta} \right| = 0, \quad \sup_{D_3} |L_{-m}(\phi_N)| = \mathcal{O}\left(\frac{1}{N^2}\right)$$

* For $(\xi, \eta) \in D_5$, $\phi_N(\xi, \eta) = \theta_1(N\xi)\theta_2\left(\frac{\eta}{N}\right)$ and thus

$$\frac{\partial \phi_N}{\partial \xi}(\xi, \eta) = N\theta'_1(N\xi)\theta_2\left(\frac{\eta}{N}\right) \quad , \quad \frac{\partial \phi_N}{\partial \eta}(\xi, \eta) = \frac{1}{N}\theta_1(N\xi)\theta'_2\left(\frac{\eta}{N}\right),$$

$$L_{-m}\phi_N(\xi, \eta) = N^2\theta_1''(N\xi)\theta_2\left(\frac{\eta}{N}\right) + \frac{1}{N^2}\theta_1(N\xi)\theta_2''\left(\frac{\eta}{N}\right) - \frac{m}{\xi}N\theta_1'(N\xi)\theta_2\left(\frac{\eta}{N}\right)$$

which give us

$$\sup_{D_5} \left| \frac{\partial\phi_N}{\partial\xi} \right| = \mathcal{O}(N), \quad \sup_{D_5} \left| \frac{\partial\phi_N}{\partial\eta} \right| = \mathcal{O}\left(\frac{1}{N}\right), \quad \sup_{D_5} |L_{-m}(\phi_N)| = \mathcal{O}(N^2).$$

* So does same with D_8 .

*For $(\xi, \eta) \in D_6$, $\phi_N(\xi, \eta) = \theta_2\left(\frac{\xi}{N}\right)\theta_2\left(\frac{\eta}{N}\right)$ and thus

$$\frac{\partial\phi_N}{\partial\xi}(\xi, \eta) = \frac{1}{N}\theta_2'\left(\frac{\xi}{N}\right)\theta_2\left(\frac{\eta}{N}\right), \quad \frac{\partial\phi_N}{\partial\eta}(\xi, \eta) = \frac{1}{N}\theta_2\left(\frac{\xi}{N}\right)\theta_2'\left(\frac{\eta}{N}\right),$$

$$L_{-m}\phi_N(\xi, \eta) = \frac{1}{N^2}\theta_2''\left(\frac{\xi}{N}\right)\theta_2\left(\frac{\eta}{N}\right) + \frac{1}{N^2}\theta_2\left(\frac{\xi}{N}\right)\theta_2''\left(\frac{\eta}{N}\right) - \frac{m}{N\xi}\theta_2'\left(\frac{\xi}{N}\right)\theta_2\left(\frac{\eta}{N}\right)$$

which give us

$$\sup_{D_6} \left| \frac{\partial\phi_N}{\partial\xi} \right| = \mathcal{O}\left(\frac{1}{N}\right), \quad \sup_{D_6} \left| \frac{\partial\phi_N}{\partial\eta} \right| = \mathcal{O}\left(\frac{1}{N}\right), \quad \sup_{D_6} |L_{-m}(\phi_N)| = \mathcal{O}\left(\frac{1}{N^2}\right).$$

* So does same with D_7 . Hence the lemma resulting. \square

We now estimate the following quantities for $i \in \{1, \dots, 8\}$:

$$\int_{D_i} |F_m| d\xi d\eta, \quad \int_{D_i} |\partial_\xi F_m| d\xi d\eta \quad \text{et} \quad \int_{D_i} |\partial_\eta F_m| d\xi d\eta.$$

Lemma 5.4. *For $\text{Re } m < 1$, we have :*

- for $i = 1$:

$$\int_{D_i} |F_m| d\xi d\eta = \mathcal{O}\left(\frac{1}{N^2}\right), \quad \int_{D_i} \left| \frac{\partial F_m}{\partial\xi} \right| d\xi d\eta = \mathcal{O}\left(\frac{1}{N}\right).$$

- for $i = 2, 4$:

$$\int_{D_i} |F_m| d\xi d\eta = \mathcal{O}(N^2), \quad \int_{D_i} \left| \frac{\partial F_m}{\partial\eta} \right| d\xi d\eta = \mathcal{O}(N).$$

- for $i = 3$:

$$\int_{D_i} |F_m| d\xi d\eta = \mathcal{O}(N^2), \quad \int_{D_i} \left| \frac{\partial F_m}{\partial\xi} \right| d\xi d\eta = \mathcal{O}(N).$$

- for $i = 5, 8$:

$$\int_{D_i} |F_m| d\xi d\eta = \mathcal{O}\left(\frac{1}{N^2}\right), \quad \int_{D_i} \left| \frac{\partial F_m}{\partial\xi} \right| d\xi d\eta = \mathcal{O}\left(\frac{1}{N}\right),$$

$$\int_{D_i} \left| \frac{\partial F_m}{\partial\eta} \right| d\xi d\eta = \mathcal{O}\left(\frac{1}{N^2}\right).$$

- for $i = 6, 7$:

$$\int_{D_i} |F_m| d\xi d\eta = \mathcal{O}(N^2), \quad \int_{D_i} \left| \frac{\partial F_m}{\partial \xi} \right| d\xi d\eta = \mathcal{O}(N),$$

$$\int_{D_i} \left| \frac{\partial F_m}{\partial \eta} \right| d\xi d\eta = \mathcal{O}(N).$$

Proof. For $\text{Re } m < 1$, we have

$$F_m(\xi, \eta) = -\frac{\xi x^{1-m}}{2\pi} \int_{\theta=0}^{\pi} \frac{\sin^{1-m} \theta d\theta}{[(x-\xi)^2 + 4x\xi \sin^2(\frac{\theta}{2}) + (y-\eta)^2]^{1-\frac{m}{2}}}.$$

therefore there is a constant C_1 such that for all $(\xi, \eta) \in \mathbb{H}^+$, we have

$$|F_m(\xi, \eta)| \leq \frac{C_1 \xi}{[(x-\xi)^2 + (\eta-y)^2]^{1-\frac{\text{Re } m}{2}}}. \quad (5.1)$$

Similarly, we have

$$\frac{\partial F_m}{\partial \xi} = \frac{F_m}{\xi} - \frac{\xi x^{1-m}}{2\pi} (m-2) \int_{\theta=0}^{\pi} \frac{[(\xi-x) + 2x \sin^2(\frac{\theta}{2})] \sin^{1-m} \theta d\theta}{[(x-\xi)^2 + 4x\xi \sin^2(\frac{\theta}{2}) + (y-\eta)^2]^{2-\frac{m}{2}}},$$

and as before, as

$$\forall \theta \in [0, \pi], \quad \left| \frac{[(\xi-x) + 2x \sin^2(\frac{\theta}{2})] \sin^{1-m} \theta}{[(x-\xi)^2 + 4x\xi \sin^2(\frac{\theta}{2}) + (y-\eta)^2]^{2-\frac{m}{2}}} \right| \leq \frac{|(\xi-x) + 2x \sin^2(\frac{\theta}{2})|}{((x-\xi)^2 + (\eta-y)^2)^{2-\frac{\text{Re } m}{2}}}$$

$$= \frac{|\xi - x \cos \theta|}{((x-\xi)^2 + (\eta-y)^2)^{2-\frac{\text{Re } m}{2}}} \leq \frac{\xi + x}{((x-\xi)^2 + (\eta-y)^2)^{2-\frac{\text{Re } m}{2}}},$$

there exists a constant C_2 such that for all N large enough and for all $(\xi, \eta) \in \mathbb{H}^+$, we have

$$\left| \frac{\partial F_m}{\partial \xi} \right| \leq C_2 \left[\frac{1}{[(x-\xi)^2 + (\eta-y)^2]^{1-\frac{\text{Re } m}{2}}} + \frac{\xi(x+\xi)}{((x-\xi)^2 + (\eta-y)^2)^{2-\frac{\text{Re } m}{2}}} \right]. \quad (5.2)$$

Finally, as

$$\frac{\partial F_m}{\partial \eta} = (2-m)(\eta-y) \frac{\xi x^{1-m}}{2\pi} \int_{\theta=0}^{\pi} \frac{\sin^{1-m} \theta}{[(x-\xi)^2 + 4x\xi \sin^2(\frac{\theta}{2}) + (y-\eta)^2]^{2-\frac{m}{2}}},$$

there exists a constant C_3 such that for all N large enough and for all $(\xi, \eta) \in \mathbb{H}^+$, we have

$$\left| \frac{\partial F_m}{\partial \eta} \right| \leq \frac{C_3 \xi}{|\eta-y|^{3-\text{Re } m}}. \quad (5.3)$$

Using these inequalities, we estimate integrals of these functions on the domains D_i .

On D_1 : Inequality (5.1) give us

$$\begin{aligned} \int_{D_1} |F_m| d\xi d\eta &= \mathcal{O}(1) \int_{\xi=\frac{1}{2N}}^{\xi=\frac{1}{N}} \int_{\eta=-\frac{N}{2}}^{\eta=\frac{N}{2}} \frac{\xi d\xi d\eta}{[(x-\xi)^2 + (\eta-y)^2]^{1-\frac{\operatorname{Re} m}{2}}} \\ &= \mathcal{O}(1/N^2) \int_{\eta=-\frac{N}{2}}^{\eta=\frac{N}{2}} \frac{d\eta}{[(x-\frac{1}{N})^2 + (\eta-y)^2]^{1-\frac{\operatorname{Re} m}{2}}} = \mathcal{O}(1/N^2). \end{aligned}$$

Then, thanks to (5.2), we have

$$\begin{aligned} \int_{D_1} \left| \frac{\partial F_m}{\partial \xi} \right| d\xi d\eta &= \mathcal{O}(1) \int_{\xi=\frac{1}{2N}}^{\xi=\frac{1}{N}} \int_{\eta=-\frac{N}{2}}^{\eta=\frac{N}{2}} \left[\frac{1}{[(x-\xi)^2 + (\eta-y)^2]^{1-\frac{\operatorname{Re} m}{2}}} \right. \\ &\quad \left. + \frac{\xi(x+\xi)}{((x-\xi)^2 + (\eta-y)^2)^{2-\frac{\operatorname{Re} m}{2}}} \right] d\xi d\eta \\ &= \mathcal{O}(1/N) \int_{\eta=-\frac{N}{2}}^{\eta=\frac{N}{2}} \frac{d\eta}{[(x-\frac{1}{N})^2 + (\eta-y)^2]^{1-\frac{\operatorname{Re} m}{2}}} + \mathcal{O}(1/N^2) = \mathcal{O}(1/N). \end{aligned}$$

On D_2 : due to inequality (5.1), we have

$$\begin{aligned} \int_{D_2} |F_m| d\xi d\eta &= \mathcal{O}(1) \int_{\xi=\frac{1}{N}}^{\xi=\frac{N}{2}} \int_{\eta=\frac{N}{2}}^{\eta=N} \frac{\xi d\xi d\eta}{[(x-\xi)^2 + (\eta-y)^2]^{1-\frac{\operatorname{Re} m}{2}}} \\ &= \mathcal{O}(1) \int_{\xi=\frac{1}{N}}^{\xi=\frac{N}{2}} \int_{\eta=\frac{N}{2}}^{\eta=N} \frac{\xi}{|\eta-y|^{2-\operatorname{Re} m}} d\xi d\eta = \mathcal{O}(N^2) \int_{\eta=\frac{N}{2}}^{\eta=N} \frac{d\eta}{|\eta-y|^{2-\operatorname{Re} m}} \\ &= \mathcal{O}(N^2) \left[\frac{1}{(N-y)^{1-\operatorname{Re} m}} - \frac{1}{(\frac{N}{2}-y)^{1-\operatorname{Re} m}} \right] = \mathcal{O}(N^{\operatorname{Re} m+1}). \end{aligned}$$

Then, thanks to (5.3), we have

$$\begin{aligned} \int_{D_2} \left| \frac{\partial F_m}{\partial \eta} \right| d\xi d\eta &= \mathcal{O}(1) \int_{\xi=\frac{1}{N}}^{\xi=\frac{N}{2}} \int_{\eta=\frac{N}{2}}^{\eta=N} \frac{\xi}{|\eta-y|^{3-\operatorname{Re} m}} d\xi d\eta \\ &= \mathcal{O}(N^2) \int_{\eta=\frac{N}{2}}^{\eta=N} \frac{d\eta}{|\eta-y|^{3-\operatorname{Re} m}} = \mathcal{O}(N^{\operatorname{Re} m}) \end{aligned}$$

On D_3 : due to inequality (5.1), we have

$$\begin{aligned} \int_{D_3} |F_m| d\xi d\eta &= \mathcal{O}(1) \int_{\xi=\frac{N}{2}}^{\xi=N} \int_{\eta=-\frac{N}{2}}^{\eta=\frac{N}{2}} \frac{\xi d\xi d\eta}{[(x-\xi)^2 + (\eta-y)^2]^{1-\frac{\operatorname{Re} m}{2}}} \\ &= \mathcal{O}(1) \int_{\xi=\frac{N}{2}}^{\xi=N} \int_{\eta=-\frac{N}{2}}^{\eta=\frac{N}{2}} \frac{\xi d\xi d\eta}{[(x-\frac{N}{2})^2 + (\eta-y)^2]^{1-\frac{\operatorname{Re} m}{2}}} \end{aligned}$$

$$\begin{aligned}
&= \mathcal{O}(N^2) \int_{\eta=-\frac{N}{2}}^{\frac{N}{2}} \frac{d\eta}{\left[(x - \frac{N}{2})^2 + (\eta - y)^2\right]^{1 - \frac{\operatorname{Re} m}{2}}} \\
&= \mathcal{O}(N^2) \int_{\eta=-\frac{N}{2}}^{\frac{N}{2}} \frac{d\eta}{[1 + (\eta - y)^2]^{1 - \frac{\operatorname{Re} m}{2}}} = \mathcal{O}(N^2).
\end{aligned}$$

Then, thanks to (5.2), we have

$$\begin{aligned}
\int_{D_3} \left| \frac{\partial F_m}{\partial \xi} \right| d\xi d\eta &= \mathcal{O}(1) \int_{\xi=\frac{N}{2}}^N \int_{\eta=-\frac{N}{2}}^{\frac{N}{2}} \left[\frac{1}{\left[(x - \xi)^2 + (\eta - y)^2\right]^{1 - \frac{\operatorname{Re} m}{2}}} \right. \\
&\quad \left. + \frac{\xi(x + \xi)}{\left((x - \xi)^2 + (\eta - y)^2\right)^{2 - \frac{\operatorname{Re} m}{2}}} \right] d\xi d\eta \\
&= \mathcal{O}(N) \int_{\eta=-\frac{N}{2}}^{\frac{N}{2}} \frac{d\eta}{\left[(x - \frac{N}{2})^2 + (\eta - y)^2\right]^{1 - \frac{\operatorname{Re} m}{2}}} + \mathcal{O}(N^3) \int_{\eta=-\frac{N}{2}}^{\frac{N}{2}} \frac{d\eta}{\left[(x - \frac{N}{2})^2 + (\eta - y)^2\right]^{2 - \frac{\operatorname{Re} m}{2}}} \\
&= \mathcal{O}(N) + \mathcal{O}(N^3) \int_{\eta=-\frac{N}{2}}^{\frac{N}{2}} \frac{d\eta}{\left(x - \frac{N}{2}\right)^{4 - \operatorname{Re} m}} \\
&= \mathcal{O}(N) + \mathcal{O}(N^{\operatorname{Re} m}) = \mathcal{O}(N).
\end{aligned}$$

On D_4 : this case is analogous to the case D_2 .

Sur D_5 : due to inequality (5.1), we have

$$\begin{aligned}
\int_{D_5} |F_m| d\xi d\eta &= \mathcal{O}(1) \int_{\xi=\frac{1}{2N}}^{\frac{1}{N}} \int_{\eta=\frac{N}{2}}^N \frac{\xi d\xi d\eta}{\left[(x - \xi)^2 + (\eta - y)^2\right]^{1 - \frac{\operatorname{Re} m}{2}}} \\
&= \mathcal{O}(1/N^2) \int_{\eta=\frac{N}{2}}^N \frac{d\eta}{(\eta - y)^{2 - \operatorname{Re} m}} \\
&= \mathcal{O}(1/N^2) \left[\frac{1}{(N - y)^{1 - \operatorname{Re} m}} - \frac{1}{\left(\frac{N}{2} - y\right)^{1 - \operatorname{Re} m}} \right] = \mathcal{O}(1/N^{3 - \operatorname{Re} m}).
\end{aligned}$$

Then, thanks to (5.2), we have

$$\begin{aligned}
\int_{D_5} \left| \frac{\partial F_m}{\partial \xi} \right| d\xi d\eta &= \mathcal{O}(1) \int_{\xi=\frac{1}{2N}}^{\frac{1}{N}} \int_{\eta=\frac{N}{2}}^N \left[\frac{1}{\left[(x - \xi)^2 + (\eta - y)^2\right]^{1 - \frac{\operatorname{Re} m}{2}}} \right. \\
&\quad \left. + \frac{\xi(x + \xi)}{\left((x - \xi)^2 + (\eta - y)^2\right)^{2 - \frac{\operatorname{Re} m}{2}}} \right] d\xi d\eta
\end{aligned}$$

$$\begin{aligned}
 &= \mathcal{O}(1) \int_{\xi=\frac{1}{2N}}^{\xi=\frac{1}{N}} \int_{\eta=\frac{N}{2}}^{\eta=N} \left[\frac{1}{\left[\left(x - \frac{1}{N}\right)^2 + (\eta - y)^2 \right]^{1 - \frac{\operatorname{Re} m}{2}}} \right. \\
 &\quad \left. + \frac{\xi(x + \xi)}{\left(\left(x - \frac{1}{N}\right)^2 + (\eta - y)^2 \right)^{2 - \frac{\operatorname{Re} m}{2}}} \right] d\xi d\eta \\
 &= \mathcal{O}\left(\frac{1}{N}\right)
 \end{aligned}$$

The estimate (5.3) gives

$$\int_{D_5} \left| \frac{\partial F_m}{\partial \eta} \right| d\xi d\eta = \mathcal{O}(1) \int_{\xi=\frac{1}{2N}}^{\xi=\frac{1}{N}} \int_{\eta=\frac{N}{2}}^{\eta=N} \frac{\xi d\xi d\eta}{|\eta - y|^{3 - \operatorname{Re} m}} = \mathcal{O}\left(\frac{1}{N^2}\right)$$

On D_6 : due to (5.1), we have

$$\begin{aligned}
 \int_{D_6} |F_m| d\xi d\eta &= \mathcal{O}(1) \int_{\xi=\frac{N}{2}}^{\xi=N} \int_{\eta=\frac{N}{2}}^{\eta=N} \frac{\xi d\xi d\eta}{\left[(x - \xi)^2 + (\eta - y)^2 \right]^{1 - \frac{\operatorname{Re} m}{2}}} \\
 &= \mathcal{O}(N^2) \int_{\eta=\frac{N}{2}}^{\eta=N} \frac{d\eta}{(\eta - y)^{2 - \operatorname{Re} m}} \\
 &= \mathcal{O}(N^2) \left[\frac{1}{(N - y)^{1 - \operatorname{Re} m}} - \frac{1}{\left(\frac{N}{2} - y\right)^{1 - \operatorname{Re} m}} \right] = \mathcal{O}(N^{1 + \operatorname{Re} m}).
 \end{aligned}$$

Then, thanks to (5.2), we have

$$\begin{aligned}
 \int_{D_6} \left| \frac{\partial F_m}{\partial \xi} \right| d\xi d\eta &= \mathcal{O}(1) \int_{\xi=\frac{N}{2}}^{\xi=N} \int_{\eta=\frac{N}{2}}^{\eta=N} \left[\frac{1}{\left[(x - \xi)^2 + (\eta - y)^2 \right]^{1 - \frac{\operatorname{Re} m}{2}}} \right. \\
 &\quad \left. + \frac{\xi(x + \xi)}{\left((x - \xi)^2 + (\eta - y)^2 \right)^{2 - \frac{\operatorname{Re} m}{2}}} \right] d\xi d\eta \\
 &= \mathcal{O}(1) \int_{\xi=\frac{N}{2}}^{\xi=N} \int_{\eta=\frac{N}{2}}^{\eta=N} \left[\frac{1}{(\eta - y)^{2 - \operatorname{Re} m}} + \frac{\xi(x + \xi)}{(\eta - y)^{4 - \operatorname{Re} m}} \right] d\xi d\eta \\
 &= \mathcal{O}(N) + \mathcal{O}(N^3) \int_{\eta=\frac{N}{2}}^{\eta=N} \frac{d\eta}{(\eta - y)^{4 - \operatorname{Re} m}} = \mathcal{O}(N) + \mathcal{O}(N^{\operatorname{Re} m}) = \mathcal{O}(N).
 \end{aligned}$$

The estimate (5.3) gives

$$\begin{aligned}
 \int_{D_6} \left| \frac{\partial F_m}{\partial \eta} \right| d\xi d\eta &= \mathcal{O}(1) \int_{\xi=\frac{N}{2}}^{\xi=N} \int_{\eta=\frac{N}{2}}^{\eta=N} \frac{\xi d\xi d\eta}{|\eta - y|^{3 - \operatorname{Re} m}} = \mathcal{O}(N^2) \int_{\eta=\frac{N}{2}}^{\eta=N} \frac{d\eta}{|\eta - y|^{3 - \operatorname{Re} m}} \\
 &= \mathcal{O}(N^{\operatorname{Re} m}).
 \end{aligned}$$

Sur D_7 : this case is analogous to the case D_6 .

Sur D_8 : this case is analogous to the case D_5 . \square

Lemma 5.5. *For $\operatorname{Re} m \geq 1$, all estimations obtained on the Lemma 5.4 are true.*

Proof. For $\operatorname{Re} m \geq 1$, we have

$$F_m(x, y, \xi, \eta) = -\frac{\xi^m}{2\pi} \int_{\theta=0}^{\pi} \sin^{m-1} \theta \left(\frac{1}{[(x-\xi)^2 + 4x\xi \sin^2 \frac{\theta}{2} + (y-\eta)^2]^{m/2}} - \frac{1}{[(x+\xi)^2 - 4x\xi \sin^2 \frac{\theta}{2} + (y-\eta)^2]^{m/2}} \right) d\theta.$$

Since for all $(\xi, \eta) \in \mathbb{H}^+$, we have

$$\left| \left[(x+\xi)^2 - 4x\xi \sin^2 \frac{\theta}{2} + (y-\eta)^2 \right]^{m/2} \right| = \left| [x^2 + \xi^2 + 2x\xi \cos \theta + (y-\eta)^2]^{m/2} \right|,$$

then for all $(\xi, \eta) \in \mathbb{H}^+$,

$$\left| \left[(x+\xi)^2 - 4x\xi \sin^2 \frac{\theta}{2} + (y-\eta)^2 \right]^{m/2} \right| \geq ((x-\xi)^2 + (y-\eta)^2)^{\frac{\operatorname{Re} m}{2}} \quad (5.4)$$

and there is a constant C'_1 such that for all $(\xi, \eta) \in \mathbb{H}^+$, we have

$$|F_m| \leq \frac{C'_1 \xi^{\operatorname{Re} m}}{((x-\xi)^2 + (y-\eta)^2)^{\frac{\operatorname{Re} m}{2}}}. \quad (5.5)$$

This inequality does not suffice to estimate integrals supported on D_1 . We can improve inequality (5.5) as follows :

We rewrite F_m as

$$F_m(x, y, \xi, \eta) = -\frac{\xi^m}{2\pi} \int_{\theta=0}^{\pi} \sin^{m-1} \theta K_m(x, y, \xi, \eta, \theta) d\theta$$

where

$$K_m(x, y, \xi, \eta, \theta) = \frac{1}{[(x-\xi)^2 + 4x\xi \sin^2 \frac{\theta}{2} + (y-\eta)^2]^{m/2}} - \frac{1}{[(x+\xi)^2 - 4x\xi \sin^2 \frac{\theta}{2} + (y-\eta)^2]^{m/2}}.$$

For $(x, y) \in \mathbb{H}^+$ fixed, $\theta \in [0, \pi]$ fixed and $\eta \in \mathbb{R}$ fixed, we define the function g_m on $[-1/N, 1/N]$ with $1/N < x$ by

$$g_m(\xi) = \frac{1}{[(x-\xi)^2 + 4x\xi \sin^2 \frac{\theta}{2} + (y-\eta)^2]^{m/2}}.$$

This function is well defined because

$$(x-\xi)^2 + 4x\xi \sin^2 \frac{\theta}{2} + (y-\eta)^2 = x^2 + \xi^2 - 2x\xi \cos \theta + (y-\eta)^2 \geq (x-|\xi|)^2 + (y-\eta)^2$$

and this last term is larger than $(x - 1/N)^2 > 0$.

We have

$$K_m(x, y, \xi, \eta, \theta) = g_m(\xi) - g_m(-\xi)$$

thus

$$|K_m(x, y, \xi, \eta, \theta)| \leq 2\xi \sup_{[-\xi, \xi]} |g'_m| \leq 2|m|\xi \frac{|\xi - x| + 2x}{[(x - \xi)^2 + (y - \eta)^2]^{1 + \frac{1}{2}\text{Re } m}},$$

which implies that there exists a constant c'_1 such that

$$\forall (\xi, \eta) \in D_1, \quad |F_m| \leq c'_1 \frac{\xi^{\text{Re } m + 1}}{[(x - \xi)^2 + (\eta - y)^2]^{1 + \frac{1}{2}\text{Re } m}}. \quad (5.6)$$

Similarly, we have

$$\begin{aligned} \frac{\partial F_m}{\partial \xi} = \frac{m F_m}{\xi} + \frac{m \xi^m}{2\pi} \int_{\theta=0}^{\pi} \sin^{m-1} \theta \left(\frac{(\xi - x) + 2x \sin^2 \frac{\theta}{2}}{[(x - \xi)^2 + 4x\xi \sin^2 \frac{\theta}{2} + (y - \eta)^2]^{\frac{m}{2} + 1}} \right. \\ \left. - \frac{(\xi + x) - 2x \sin^2 \frac{\theta}{2}}{[(x + \xi)^2 - 4x\xi \sin^2 \frac{\theta}{2} + (y - \eta)^2]^{\frac{m}{2} + 1}} \right) d\theta. \end{aligned} \quad (5.7)$$

and as before,

$$\begin{aligned} \forall \theta \in [0, \pi], \quad \left| \frac{[(\xi - x) + 2x \sin^2 \frac{\theta}{2}] \sin^{m-1} \theta}{[(x - \xi)^2 + 4x\xi \sin^2 \frac{\theta}{2} + (y - \eta)^2]^{\frac{m}{2} + 1}} \right| &\leq \frac{|(\xi - x) + 2x \sin^2 \frac{\theta}{2}|}{[(x - \xi)^2 + (y - \eta)^2]^{\frac{\text{Re } m}{2} + 1}} \\ &= \frac{|\xi - x \cos \theta|}{[(x - \xi)^2 + (y - \eta)^2]^{\frac{\text{Re } m}{2} + 1}} \leq \frac{\xi + x}{[(x - \xi)^2 + (y - \eta)^2]^{\frac{\text{Re } m}{2} + 1}} \end{aligned}$$

and thanks to (5.4) :

$$\begin{aligned} \forall \theta \in [0, \pi], \quad \left| \frac{[(\xi + x) - 2x \sin^2 \frac{\theta}{2}] \sin^{m-1} \theta}{[(x + \xi)^2 - 4x\xi \sin^2 \frac{\theta}{2} + (y - \eta)^2]^{\frac{m}{2} + 1}} \right| &\leq \frac{|(\xi + x) - 2x \sin^2 \frac{\theta}{2}|}{[(x - \xi)^2 + (y - \eta)^2]^{\frac{\text{Re } m}{2} + 1}} \\ &= \frac{|\xi + x \cos \theta|}{[(x - \xi)^2 + (y - \eta)^2]^{\frac{\text{Re } m}{2} + 1}} \leq \frac{\xi + x}{[(x - \xi)^2 + (y - \eta)^2]^{\frac{\text{Re } m}{2} + 1}}. \end{aligned}$$

Those estimations, the formula (5.7) and the inequality (5.5) show that there is a constant C'_2 such that large enough N and for all $(\xi, \eta) \in \mathbb{H}^+$, we have

$$\left| \frac{\partial F_m}{\partial \xi} \right| \leq C'_2 \left(\frac{\xi^{\text{Re } m - 1}}{[(x - \xi)^2 + (y - \eta)^2]^{\frac{\text{Re } m}{2}}} + \frac{\xi^{\text{Re } m} (\xi + x)}{[(x - \xi)^2 + (y - \eta)^2]^{\frac{\text{Re } m}{2} + 1}} \right). \quad (5.8)$$

We can improve this inequality on D_1 , for this, we need to use the inequality (5.6) instead of (5.5) and we obtain that there is two constants C''_2 and C'''_2 (which do not depend of N) such that for all $(\xi, \eta) \in D_1$

$$\left| \frac{\partial F_m}{\partial \xi} \right| \leq C''_2 \left(\frac{\xi^{\text{Re } m}}{[(x - \xi)^2 + (y - \eta)^2]^{1 + \frac{\text{Re } m}{2}}} + \frac{\xi^{\text{Re } m} (\xi + x)}{[(x - \xi)^2 + (y - \eta)^2]^{\frac{\text{Re } m}{2} + 1}} \right)$$

$$\leq C_2''' \frac{\xi^{\operatorname{Re} m}}{[(x - \xi)^2 + (y - \eta)^2]^{1 + \frac{\operatorname{Re} m}{2}}} \quad (5.9)$$

Finally,

$$\frac{\partial F_m}{\partial \eta} = \frac{m(\eta - y)\xi^m}{2\pi} \int_{\theta=0}^{\pi} \sin^{m-1} \theta \left(\frac{1}{[(x - \xi)^2 + 4x\xi \sin^2 \frac{\theta}{2} + (y - \eta)^2]^{\frac{m}{2}+1}} - \frac{1}{[(x + \xi)^2 - 4x\xi \sin^2 \frac{\theta}{2} + (y - \eta)^2]^{\frac{m}{2}+1}} \right) d\theta.$$

Similarly, there is a constant C_3' such that for all large enough N and for all $(\xi, \eta) \in \mathbb{H}^+$, we have

$$\left| \frac{\partial F_m}{\partial \eta} \right| \leq C_3' \frac{|\eta - y| \xi^{\operatorname{Re} m}}{((x - \xi)^2 + (y - \eta)^2)^{\frac{\operatorname{Re} m}{2}+1}}. \quad (5.10)$$

Thanks to those inequalities, we will estimate the integrals of those functions on each domain D_i .

On D_1 : due to (5.6), we have

$$\begin{aligned} \int_{D_1} |F_m| d\xi d\eta &= \mathcal{O}(1) \int_{\xi=\frac{1}{2N}}^{\xi=\frac{1}{N}} \int_{\eta=-\frac{N}{2}}^{\eta=\frac{N}{2}} \frac{\xi^{\operatorname{Re} m+1} d\xi d\eta}{[(x - \xi)^2 + (\eta - y)^2]^{1 + \frac{1}{2}\operatorname{Re} m}} \\ &= \mathcal{O}(N) \int_{\xi=\frac{1}{2N}}^{\xi=\frac{1}{N}} \xi^{\operatorname{Re} m+1} d\xi = \mathcal{O}(N) \left[\left(\frac{1}{N} \right)^{\operatorname{Re} m+2} - \left(\frac{1}{2N} \right)^{\operatorname{Re} m+2} \right] \\ &= \mathcal{O}(1/N^{\operatorname{Re} m+1}). \end{aligned}$$

Then thanks to (5.9), we have

$$\begin{aligned} \int_{D_1} \left| \frac{\partial F_m}{\partial \xi} \right| d\xi d\eta &= \mathcal{O}(1) \int_{\xi=\frac{1}{2N}}^{\xi=\frac{1}{N}} \int_{\eta=-\frac{N}{2}}^{\eta=\frac{N}{2}} \frac{\xi^{\operatorname{Re} m} d\xi d\eta}{[(x - \xi)^2 + (\eta - y)^2]^{1 + \frac{1}{2}\operatorname{Re} m}} \\ &= \mathcal{O}(N) \int_{\xi=\frac{1}{2N}}^{\xi=\frac{1}{N}} \xi^{\operatorname{Re} m} d\xi = \mathcal{O}(1/N^{\operatorname{Re} m}). \end{aligned}$$

On D_2 : due to (5.5), we have

$$\begin{aligned} \int_{D_2} |F_m| d\xi d\eta &= \mathcal{O}(1) \int_{\xi=\frac{1}{N}}^{\frac{N}{2}} \int_{\eta=\frac{N}{2}}^N \frac{\xi^{\operatorname{Re} m} d\xi d\eta}{((x - \xi)^2 + (y - \eta)^2)^{\frac{\operatorname{Re} m}{2}}} \\ &= \mathcal{O}(1) \int_{\xi=\frac{1}{N}}^{\frac{N}{2}} \int_{\eta=\frac{N}{2}}^N \frac{\xi^{\operatorname{Re} m} d\xi d\eta}{|y - \frac{N}{2}|^{\operatorname{Re} m}} = \mathcal{O}(N^2), \end{aligned}$$

because we integrate a bounded function (independently of N) on a domain with measure controlled by $\mathcal{O}(N^2)$.

Then, the inequality (5.10) implies

$$\begin{aligned} \int_{D_2} \left| \frac{\partial F_m}{\partial \eta} \right| d\xi d\eta &= \mathcal{O}(1) \int_{\xi=\frac{1}{N}}^{\frac{N}{2}} \int_{\eta=\frac{N}{2}}^N \frac{|\eta - y| \xi^{\operatorname{Re} m} d\xi d\eta}{((x - \xi)^2 + (y - \eta)^2)^{\frac{\operatorname{Re} m}{2} + 1}} \\ &= \mathcal{O}(1) \int_{\xi=\frac{1}{N}}^{\frac{N}{2}} \int_{\eta=\frac{N}{2}}^N \frac{\xi^{\operatorname{Re} m} d\xi d\eta}{|y - \eta|^{\operatorname{Re} m + 1}} = \mathcal{O}(1) \int_{\xi=\frac{1}{N}}^{\frac{N}{2}} \int_{\eta=\frac{N}{2}}^N \frac{N^{\operatorname{Re} m} d\xi d\eta}{|\frac{N}{2} - y|^{\operatorname{Re} m + 1}} = \mathcal{O}(N). \end{aligned}$$

On D_3 : due to (5.5), we have

$$\begin{aligned} \int_{D_3} |F_m| d\xi d\eta &= \mathcal{O}(1) \int_{\xi=\frac{N}{2}}^N \int_{\eta=-\frac{N}{2}}^{\frac{N}{2}} \frac{\xi^{\operatorname{Re} m} d\xi d\eta}{((x - \xi)^2 + (y - \eta)^2)^{\frac{\operatorname{Re} m}{2}}} \\ &= \mathcal{O}(1) \int_{\xi=\frac{N}{2}}^N \int_{\eta=-\frac{N}{2}}^{\frac{N}{2}} \frac{\xi^{\operatorname{Re} m} d\xi d\eta}{((x - \frac{N}{2})^2 + (y - \eta)^2)^{\frac{\operatorname{Re} m}{2}}} \\ &= \mathcal{O}(N^{\operatorname{Re} m + 1}) \int_{\eta=-\frac{N}{2}}^{\frac{N}{2}} \frac{d\eta}{((x - \frac{N}{2})^2 + (y - \eta)^2)^{\frac{\operatorname{Re} m}{2}}} = \mathcal{O}(N^2). \end{aligned}$$

Then, thanks to (5.8), we have

$$\begin{aligned} \int_{D_3} \left| \frac{\partial F_m}{\partial \xi} \right| d\xi d\eta &= \mathcal{O}(1) \int_{\xi=\frac{N}{2}}^N \int_{\eta=-\frac{N}{2}}^{\frac{N}{2}} \left(\frac{\xi^{\operatorname{Re} m - 1}}{[(x - \xi)^2 + (y - \eta)^2]^{\frac{\operatorname{Re} m}{2}}} \right. \\ &\quad \left. + \frac{\xi^{\operatorname{Re} m} (\xi + x)}{[(x - \xi)^2 + (y - \eta)^2]^{\frac{\operatorname{Re} m}{2} + 1}} \right) d\xi d\eta \\ &= \mathcal{O}(N^{\operatorname{Re} m}) \int_{\eta=-\frac{N}{2}}^{\frac{N}{2}} \frac{d\eta}{[(x - \frac{N}{2})^2 + (y - \eta)^2]^{\frac{\operatorname{Re} m}{2}}} \\ &\quad + \mathcal{O}(N^{\operatorname{Re} m + 2}) \int_{\eta=-\frac{N}{2}}^{\frac{N}{2}} \frac{d\eta}{[(x - \frac{N}{2})^2 + (y - \eta)^2]^{\frac{\operatorname{Re} m}{2} + 1}} \\ &= \mathcal{O}(N) + \mathcal{O}(N) = \mathcal{O}(N). \end{aligned}$$

On D_4 : this case is analogous to the case D_2 .

On D_5 : due to (5.5), we have

$$\begin{aligned} \int_{D_5} |F_m| d\xi d\eta &= \mathcal{O}(1) \int_{\xi=\frac{1}{2N}}^{\frac{1}{N}} \int_{\eta=\frac{N}{2}}^N \frac{\xi^{\operatorname{Re} m} d\xi d\eta}{((x - \xi)^2 + (y - \eta)^2)^{\frac{\operatorname{Re} m}{2}}} \\ &= \mathcal{O}(1/N^{\operatorname{Re} m + 1}) \int_{\eta=\frac{N}{2}}^N \frac{d\eta}{|y - \frac{N}{2}|^{\operatorname{Re} m}} = \mathcal{O}(1/N^{2\operatorname{Re} m}). \end{aligned}$$

Then, thanks to (5.8), we have

$$\int_{D_5} \left| \frac{\partial F_m}{\partial \xi} \right| d\xi d\eta = \mathcal{O}(1) \int_{\xi=\frac{1}{2N}}^{\frac{1}{N}} \int_{\eta=\frac{N}{2}}^N \left(\frac{\xi^{\operatorname{Re} m - 1}}{[(x - \xi)^2 + (y - \eta)^2]^{\frac{\operatorname{Re} m}{2}}} \right)$$

$$\begin{aligned}
& + \frac{\xi^{\operatorname{Re} m}(\xi + x)}{[(x - \xi)^2 + (y - \eta)^2]^{\frac{\operatorname{Re} m}{2} + 1}} \Big) d\xi d\eta \\
& = \mathcal{O}(1) \int_{\xi=\frac{1}{2N}}^{\xi=\frac{1}{N}} \int_{\eta=\frac{N}{2}}^{\eta=N} \left(\frac{\xi^{\operatorname{Re} m - 1}}{|y - \eta|^{\operatorname{Re} m}} + \frac{\xi^{\operatorname{Re} m}(\xi + x)}{|y - \eta|^{\operatorname{Re} m + 2}} \right) d\xi d\eta \\
& = \mathcal{O}(1/N^{2\operatorname{Re} m - 1}).
\end{aligned}$$

With the inequality (5.10), we have

$$\begin{aligned}
\int_{D_5} \left| \frac{\partial F_m}{\partial \eta} \right| d\xi d\eta & = \mathcal{O}(1) \int_{\xi=\frac{1}{2N}}^{\xi=\frac{1}{N}} \int_{\eta=\frac{N}{2}}^{\eta=N} \frac{|\eta - y| \xi^{\operatorname{Re} m} d\xi d\eta}{((x - \xi)^2 + (y - \eta)^2)^{\frac{\operatorname{Re} m}{2} + 1}} \\
& = \mathcal{O}(1) \int_{\xi=\frac{1}{2N}}^{\xi=\frac{1}{N}} \int_{\eta=\frac{N}{2}}^{\eta=N} \frac{\xi^{\operatorname{Re} m} d\xi d\eta}{|y - \eta|^{\operatorname{Re} m + 1}} = \mathcal{O}(1/N^{2\operatorname{Re} m + 1})
\end{aligned}$$

On D_6 : due to (5.5), we have

$$\begin{aligned}
\int_{D_6} |F_m| d\xi d\eta & = \mathcal{O}(1) \int_{\xi=\frac{N}{2}}^{\xi=N} \int_{\eta=\frac{N}{2}}^{\eta=N} \frac{\xi^{\operatorname{Re} m} d\xi d\eta}{((x - \xi)^2 + (y - \eta)^2)^{\frac{\operatorname{Re} m}{2}}} \\
& = \mathcal{O}(N^{\operatorname{Re} m + 1}) \int_{\eta=\frac{N}{2}}^{\eta=N} \frac{d\eta}{(\frac{N}{2} - y)^{\operatorname{Re} m}} = \mathcal{O}(N^2).
\end{aligned}$$

Then, thanks to (5.8), we obtain

$$\begin{aligned}
\int_{D_6} \left| \frac{\partial F_m}{\partial \xi} \right| d\xi d\eta & = \mathcal{O}(1) \int_{\xi=\frac{N}{2}}^{\xi=N} \int_{\eta=\frac{N}{2}}^{\eta=N} \left(\frac{\xi^{\operatorname{Re} m - 1}}{[(x - \xi)^2 + (y - \eta)^2]^{\frac{\operatorname{Re} m}{2}}} \right. \\
& \quad \left. + \frac{\xi^{\operatorname{Re} m}(\xi + x)}{[(x - \xi)^2 + (y - \eta)^2]^{\frac{\operatorname{Re} m}{2} + 1}} \right) d\xi d\eta \\
& = \mathcal{O}(1) \int_{\xi=\frac{N}{2}}^{\xi=N} \int_{\eta=\frac{N}{2}}^{\eta=N} \left(\frac{\xi^{\operatorname{Re} m - 1}}{|y - \eta|^{\operatorname{Re} m}} + \frac{\xi^{\operatorname{Re} m}(\xi + x)}{|y - \eta|^{\operatorname{Re} m + 2}} \right) d\xi d\eta \\
& = \mathcal{O}(N) + \mathcal{O}(N^{\operatorname{Re} m + 2}) \int_{\eta=\frac{N}{2}}^{\eta=N} \frac{d\eta}{|y - \eta|^{\operatorname{Re} m + 2}} = \mathcal{O}(N).
\end{aligned}$$

Finally, the inequality (5.10) implies

$$\begin{aligned}
\int_{D_6} \left| \frac{\partial F_m}{\partial \eta} \right| d\xi d\eta & = \mathcal{O}(1) \int_{\xi=\frac{N}{2}}^{\xi=N} \int_{\eta=\frac{N}{2}}^{\eta=N} \frac{|\eta - y| \xi^{\operatorname{Re} m} d\xi d\eta}{((x - \xi)^2 + (y - \eta)^2)^{\frac{\operatorname{Re} m}{2} + 1}} \\
& = \mathcal{O}(N^{\operatorname{Re} m + 1}) \int_{\eta=\frac{N}{2}}^{\eta=N} \frac{d\eta}{|y - \eta|^{\operatorname{Re} m + 1}} = \mathcal{O}(N).
\end{aligned}$$

On D_7 : this case is analogous to the case D_6 .

On D_8 : this case is analogous to the case D_5 .

□

In the following table, we summarize the results obtained on the previous lemmas :

i	$\sup_{D_i} L_{-m}\phi_N $	$\int_{D_i} F_m d\xi d\eta$	$(\partial_\xi \phi_N , \partial_\eta \phi_N)$	$\int_{D_i} \partial_\xi F_m $	$\int_{D_i} \partial_\eta F_m $
1	$\mathcal{O}(N^2)$	$\mathcal{O}(1/N^2)$	$(\mathcal{O}(N), 0)$	$\mathcal{O}(\frac{1}{N})$	\times
2	$\mathcal{O}(1/N^2)$	$\mathcal{O}(N^2)$	$(0, \mathcal{O}(\frac{1}{N}))$	\times	$\mathcal{O}(N)$
3	$\mathcal{O}(1/N^2)$	$\mathcal{O}(N^2)$	$(\mathcal{O}(\frac{1}{N}), 0)$	$\mathcal{O}(N)$	\times
4	$\mathcal{O}(1/N^2)$	$\mathcal{O}(N^2)$	$(0, \mathcal{O}(\frac{1}{N}))$	\times	$\mathcal{O}(N)$
5	$\mathcal{O}(N^2)$	$\mathcal{O}(1/N^2)$	$(\mathcal{O}(N), \mathcal{O}(\frac{1}{N}))$	$\mathcal{O}(\frac{1}{N})$	$\mathcal{O}(\frac{1}{N^2})$
6	$\mathcal{O}(1/N^2)$	$\mathcal{O}(N^2)$	$(\mathcal{O}(\frac{1}{N}), \mathcal{O}(\frac{1}{N}))$	$\mathcal{O}(N)$	$\mathcal{O}(N)$
7	$\mathcal{O}(1/N^2)$	$\mathcal{O}(N^2)$	$(\mathcal{O}(\frac{1}{N}), \mathcal{O}(\frac{1}{N}))$	$\mathcal{O}(N)$	$\mathcal{O}(N)$
8	$\mathcal{O}(N^2)$	$\mathcal{O}(1/N^2)$	$(\mathcal{O}(N), \mathcal{O}(\frac{1}{N}))$	$\mathcal{O}(\frac{1}{N})$	$\mathcal{O}(\frac{1}{N^2})$

We can easily check that for each $i \in \{1, \dots, 8\}$, the quantities

$$\sup_{D_i} |L_{-m}\phi_N| \int_{D_i} |F_m|, \quad \sup_{D_i} |\partial_\xi \phi_N| \int_{D_i} |\partial_\xi F_m| \quad \text{and} \quad \sup_{D_i} |\partial_\eta \phi_N| \int_{D_i} |\partial_\eta F_m|$$

stay bounded. Therefore,

$$u(x, y) = o(1)$$

when $N \rightarrow +\infty$. Thus

$$u \equiv 0$$

and this completes the proof of the Proposition 5.2. \square

Lemma 5.6. *Let $u \in \mathcal{D}(\mathbb{H}^+)$ and let $(x, y) \in \mathbb{H}^+$, we define*

$$U(x, y) = \int_{\mathbb{H}^+} u(\xi, \eta) F_m(x, y, \xi, \eta) d\xi d\eta,$$

then $\lim_{\|(x,y)\| \rightarrow +\infty} U = 0$, and for all $y \in \mathbb{R}$, $\lim_{(0,y)} U = 0$.

Moreover, $U \in C^\infty(\mathbb{H}^+ \setminus \text{supp } u)$ and for all $(x, y) \notin \text{supp } u$, we have $L_{m,x,y} U(x, y) = 0$.

Proof. When (ξ, η) is fixed, and since

$$F_m(x, y, \xi, \eta) = -\frac{\xi x^{1-m}}{2\pi} \int_{\theta=0}^{\pi} \frac{\sin^{1-m} \theta d\theta}{[(x-\xi)^2 + 4x\xi \sin^2(\frac{\theta}{2}) + (y-\eta)^2]^{1-\frac{m}{2}}}$$

for $\text{Re } m < 1$, then $F_m(x, y, \xi, \eta) \xrightarrow{\|(x,y)\| \rightarrow +\infty} 0$ and the first result of the lemma is shown.

Similarly, if $\text{Re } m \geq 1$,

$$F_m(x, y, \xi, \eta) = -\frac{\xi^m}{2\pi} \int_{\theta=0}^{\pi} \sin^{m-1} \theta \left[\frac{1}{[(x-\xi)^2 + 4x\xi \sin^2(\frac{\theta}{2}) + (y-\eta)^2]^{\frac{m}{2}}} \right]$$

$$\left. - \frac{1}{[(x + \xi)^2 - 4x\xi \sin^2\left(\frac{\theta}{2}\right) + (y - \eta)^2]^{\frac{m}{2}}} \right] d\theta$$

then $F_m(x, y, \xi, \eta) \xrightarrow{\|(x, y)\| \rightarrow +\infty} 0$ and the first result of the lemma is shown.

For the second result, it suffices to see, for $\operatorname{Re} m < 1$, that

$$F_m(x, y, \xi, \eta) \underset{(x, y) \rightarrow (0, y')}{\sim} - \frac{\xi x^{1-m}}{2\pi[\xi^2 + (y' - \eta)^2]^{1-m/2}} \int_0^\pi \sin^{1-m} \theta d\theta$$

which implies the desired result.

Now, we assume that $\operatorname{Re} m \geq 1$. Let (ξ, η) be fixed in the support of u , which is a compact set of \mathbb{H}^+ . In particular, there exist $M > 0$ and $\alpha > 0$ which do not depend of u such that $\|(\xi, \eta)\| \leq M$ et $\xi \geq 2\alpha$. Let y be in \mathbb{R} . By denotting for $x \in [-\alpha, \alpha]$,

$$f_m(x) = \frac{1}{[(x - \xi)^2 + 4x\xi \sin^2\left(\frac{\theta}{2}\right) + (y - \eta)^2]^{\frac{m}{2}}},$$

By the mean value inequality, for $x > 0$ near 0, we have

$$|f_m(x) - f_m(0)| \leq x \sup_{[0, \alpha]} |f'_m|$$

and

$$|f_m(-x) - f_m(0)| \leq x \sup_{[-\alpha, 0]} |f'_m|.$$

then

$$|f_m(x) - f_m(-x)| \leq 2x \sup_{[-\alpha, \alpha]} |f'_m| \leq 2x|m| \frac{3M + \alpha}{\alpha^{\operatorname{Re} m + 2}}.$$

In particular,

$$\sup_{\substack{(\xi, \eta) \in \operatorname{supp} u \\ y \in \mathbb{R}}} |F_m(x, y)| = \mathcal{O}(x)$$

when $x \rightarrow 0+$. The second result is proved.

The last result can be deduced of the fact that if $(x, y) \neq (\xi, \eta)$ are both in \mathbb{H}^+ , then

$$L_{m, x, y} F_m(x, y, \xi, \eta) = 0.$$

□

Remark 5.7. *If $U \in \mathcal{D}(\mathbb{H}^+)$, then $L_{m, x, y} U = u$, but this identity is not necessary true if $U \notin \mathcal{D}(\mathbb{H}^+)$. In particular, we can not say that in the Lemma 5.6, we have $L_m U = u$.*

Now, we will prove a decomposition theorem for axisymmetric potentials, it is interesting to compare the following theorem to known result in [5, Theorem 2 section 4] (the fundamental difference is that in this work, the conductivity is not extended in all domain by reflection through the boundary $\partial\Omega$).

Theorem 5.8. *Let $m \in \mathbb{C}$. Let Ω be an open set of \mathbb{H}^+ and let K be a compact set of Ω . If $u \in C^2(\Omega \setminus K)$ satisfies $L_m u = 0$ in $\Omega \setminus K$, then u has a unique decomposition as follows :*

$$u = v + w,$$

where $v \in C^2(\Omega)$ satisfies $L_m v = 0$ in Ω and $w \in C^2(\mathbb{H}^+ \setminus K)$ satisfies $L_m w = 0$ in $\mathbb{H}^+ \setminus K$ with $\lim_{\partial\mathbb{H}^+} w = 0$.

Proof. For $E \subset \mathbb{C}$ and $\rho > 0$, we define $E_\rho = \{x \in \mathbb{C}, d(x, E) < \rho\}$ (E_ρ is a neighborhood of E).

At first, we assume that Ω is a relatively compact open set of \mathbb{H}^+ . We choose ρ as small as K_ρ and $(\partial\Omega)_\rho$ are disjoint. There is a function $\varphi_\rho \in \mathcal{D}(\mathbb{H}^+)$ compactly supported on $\Omega \setminus K$ such that $\varphi_\rho \equiv 1$ in a neighborhood of $\Omega \setminus (K_\rho \cup (\partial\Omega)_\rho)$.




Figure : $\varphi_\rho \equiv 1$ on the gray domain

For $z = x + iy \in \Omega \setminus (K_\rho \cup (\partial\Omega)_\rho)$, we denote

$$F_z(\zeta) := F_m(x, y, \xi, \eta) \quad \text{and} \quad L_\zeta := L_{m, \xi, \eta} \quad \text{for } \zeta = \xi + i\eta,$$

Thanks to Proposition 4.7, we have

$$\begin{aligned} u(z) &= u\varphi_\rho(z) = \int_{\Omega_\rho} F_z(\zeta) L_\zeta(u\varphi_\rho)(\zeta) d\xi d\eta \\ &= \int_{(\partial\Omega)_\rho} F_z(\zeta) L_\zeta(u\varphi_\rho)(\zeta) d\xi d\eta + \int_{K_\rho} F_z(\zeta) L_\zeta(u\varphi_\rho)(\zeta) d\xi d\eta \\ &= v_\rho(z) + w_\rho(z). \end{aligned}$$

Then, the last result of Lemma 5.6 shows us that v_ρ satisfies $L_m v_\rho = 0$ on $\Omega \setminus (\partial\Omega)_\rho$ and w_ρ satisfies $L_m w_\rho = 0$ on $\mathbb{H}^+ \setminus K_\rho$. We also have $\lim_{\partial\mathbb{H}^+} w_\rho = 0$.

Now, we assume that $\sigma < \rho$. As previously, we obtain the decomposition $u = v_\sigma + w_\sigma$ on $\Omega \setminus (K_\sigma \cup (\partial\Omega)_\sigma)$. We claim that $v_\rho = v_\sigma$ on $\Omega \setminus (\partial\Omega)_\rho$ and $w_\rho = w_\sigma$ on $\mathbb{H}^+ \setminus K_\rho$. To see this, note that if $z \in \Omega \setminus (K_\rho \cup (\partial\Omega)_\rho)$, then $v_\rho(z) + w_\rho(z) = v_\sigma(z) + w_\sigma(z)$.

We will designate by (1) the Weinstein equation $L_m u = 0$. Thus $w_\rho - w_\sigma$ satisfies (1) on $\mathbb{H}^+ \setminus K_\rho$, which is equal to $v_\sigma - v_\rho$ on $\Omega \setminus (K_\rho \cup (\partial\Omega)_\rho)$, therefore $v_\sigma - v_\rho$ extends to a solution of (1) on $\Omega \setminus (\partial\Omega)_\rho$.

Finally, $w_\rho - w_\sigma$ extends to a solution of (1) on \mathbb{H}^+ , and $\lim_{\partial\mathbb{H}^+} w_\rho - w_\sigma = 0$.

Due to Proposition 5.2, we have

$$w_\rho = w_\sigma,$$

and then $v_\rho = v_\sigma$.

For $z \in \Omega$, we can define $v(z) = v_\rho(z)$ for ρ as small as $z \in \Omega \setminus (\partial\Omega)_\rho$. Similarly, for $z \in \mathbb{H}^+ \setminus K$, we put $w(z) = w_\rho(z)$ for small ρ . We have proved the desired decomposition $u = v + w$.

Now, assume that Ω is an arbitrary domain of \mathbb{H}^+ and let u be a solution of $L_m u = 0$ on $\Omega \setminus K$. We choose $a \in \mathbb{H}^+$ and R large enough so that $K \subset D(a, R)$ and $D(a, R)$ be a relatively compact set of \mathbb{H}^+ . Let $\omega = \Omega \cap D(a, R)$. Note that K is a compact set of ω which is a relatively compact open set of \mathbb{H}^+ and u satisfies (1) on $\omega \setminus K$. Applying the results demonstrated for relatively compact open sets, we obtain

$$u(z) = \tilde{v}(z) + \tilde{w}(z)$$

for $z \in \omega \setminus K$, where \tilde{v} satisfies (1) on ω and \tilde{w} satisfies (1) on $\mathbb{H}^+ \setminus K$ with $\lim_{\partial\mathbb{H}^+} \tilde{w} = 0$. Note that $V = u - \tilde{w}$ satisfies (1) on $\Omega \setminus K$ and V can be extended into a solution of (1) in a neighborhood of K because $V = \tilde{v}$ on ω . The sum $u = V + \tilde{w}$ provides us a desired decomposition of u .

As before, if we have another decomposition $u = v + w$ with $v \in C^2(\Omega)$, $L_m v = 0$ and with $w \in C^2(\mathbb{H}^+ \setminus K)$, $L_m w = 0$ and $\lim_{\partial\mathbb{H}^+} w = 0$, then we have $V - v = w - \tilde{w}$ on $\Omega \setminus K$. The function $w - \tilde{w}$ can be extended on \mathbb{H}^+ into a solution of $L_m(w - \tilde{w}) = 0$ on \mathbb{H}^+ with $\lim_{\partial\mathbb{H}^+}(w - \tilde{w}) = 0$. Thanks to Proposition 5.2, we obtain $w = \tilde{w}$, then $V = v$, which completes the proof of the decomposition theorem. \square

The following proposition is a Poisson formula for the axisymmetric potentials in \mathbb{H}^+ :

Proposition 5.9. *Let $m \in \mathbb{C}$ be such that $\operatorname{Re} m < 1$ and $u : \mathbb{R} \rightarrow \mathbb{R}$ be a continuous and bounded function. Then there is a unique axisymmetric potential $U \in C^2(\mathbb{H}^+)$ such that $\lim_{\|(x,y)\| \rightarrow +\infty} U(x,y) = 0$ and for all $y \in \mathbb{R}$,*

$$\lim_{(0,y)} U = u(y).$$

Moreover, we have for all $(x,y) \in \mathbb{H}^+$,

$$U(x,y) = C_m x^{1-m} \int_{\eta=-\infty}^{\infty} \frac{u(\eta) d\eta}{(x^2 + (y-\eta)^2)^{1-\frac{m}{2}}}$$

where $C_m = \frac{1-m}{2\pi} \int_{\theta=0}^{\pi} \sin^{1-m} \theta d\theta = \frac{1}{2^m \pi} \frac{\Gamma^2(1-\frac{m}{2})}{\Gamma(1-m)}$.

Proof. We define $f(x, y) = \frac{x^{1-m}}{(x^2+(y-\eta)^2)^{1-\frac{m}{2}}}$. To show that U is a solution of $L_m U = 0$, it suffices to prove that $L_m f = 0$ by differentiation under the integral sign. We have

$$\partial_x f = \frac{(1-m)x^{-m}}{(x^2+(y-\eta)^2)^{1-\frac{m}{2}}} - \frac{(2-m)x^{2-m}}{(x^2+(y-\eta)^2)^{2-\frac{m}{2}}}$$

and

$$\partial_{xx} f = -\frac{m(1-m)x^{-m-1}}{(x^2+(y-\eta)^2)^{1-\frac{m}{2}}} - \frac{(2-m)(3-2m)x^{1-m}}{(x^2+(y-\eta)^2)^{2-\frac{m}{2}}} + \frac{(2-m)(4-m)x^{3-m}}{(x^2+(y-\eta)^2)^{3-\frac{m}{2}}}$$

and

$$\partial_{yy} f = -\frac{(2-m)x^{1-m}}{(x^2+(y-\eta)^2)^{2-\frac{m}{2}}} + \frac{(2-m)(4-m)(y-\eta)^2 x^{1-m}}{(x^2+(y-\eta)^2)^{3-\frac{m}{2}}}.$$

Then,

$$\Delta f = \frac{m(2-m)x^{1-m}}{(x^2+(y-\eta)^2)^{2-\frac{m}{2}}} - \frac{m(1-m)x^{-m-1}}{(x^2+(y-\eta)^2)^{1-\frac{m}{2}}}$$

and we deduce that $L_m f(x, y) = 0$.

We have

$$U(x, y) = C_m x^{1-m} \int_{\eta=-\infty}^{\infty} \frac{u(\eta) d\eta}{(x^2+(y-\eta)^2)^{1-\frac{m}{2}}} = \frac{C_m}{x} \int_{\eta=-\infty}^{\infty} \frac{u(\eta) d\eta}{(1+(\frac{y-\eta}{x})^2)^{1-\frac{m}{2}}}$$

By a change of variable $t = \frac{y-\eta}{x}$, we obtain

$$U(x, y) = C_m \int_{t=-\infty}^{\infty} \frac{u(y-tx) dt}{(1+t^2)^{1-\frac{m}{2}}}$$

Thanks to the dominated convergence theorem, it suffices to show that

$$C_m \int_{t=-\infty}^{\infty} \frac{dt}{(1+t^2)^{1-\frac{m}{2}}} = \frac{1-m}{2\pi} \int_{\theta=0}^{\pi} \sin^{1-m} \theta d\theta \int_{t=-\infty}^{\infty} \frac{dt}{(1+t^2)^{1-\frac{m}{2}}} = 1.$$

To see this, according [1] (page 258), note that

$$\int_{t=-\infty}^{\infty} \frac{dt}{(1+t^2)^{1-\frac{m}{2}}} = B\left(\frac{1}{2}, \frac{1-m}{2}\right) = \frac{\Gamma(1/2)\Gamma(\frac{1-m}{2})}{\Gamma(1-\frac{m}{2})}$$

where B is the Euler beta function and

$$\frac{1-m}{2\pi} \int_{\theta=0}^{\pi} \sin^{1-m} \theta d\theta = \frac{1-m}{2\pi} 2^{1-m} B\left(1-\frac{m}{2}, 1-\frac{m}{2}\right) = \frac{1-m}{2\pi} 2^{1-m} \frac{\Gamma^2(1-\frac{m}{2})}{\Gamma(2-m)}.$$

Then, using the duplication formula for the Γ function,

$$\Gamma(2z) = \pi^{-1/2} 2^{2z-1} \Gamma(z) \Gamma\left(z + \frac{1}{2}\right)$$

and the recurrence formula $\Gamma(z + 1) = z\Gamma(z)$, we obtain the desired result,

$$\frac{\Gamma(1/2)\Gamma\left(\frac{1-m}{2}\right)}{\Gamma\left(1-\frac{m}{2}\right)} \frac{1-m}{2\pi} 2^{1-m} \frac{\Gamma^2\left(1-\frac{m}{2}\right)}{\Gamma(2-m)} = 1.$$

The uniqueness follows from the proposition 5.2. So, we proved the proposition. \square

Remark 5.10. *We could ask ourselves the question of the existence of a such reproducing formula if $\operatorname{Re} m \geq 1$. In fact, if $m \in \mathbb{N}^*$ and if $u \in C^2(\overline{\mathbb{H}^+})$ satisfies $L_m(u) = 0$ on \mathbb{H}^+ , then the function v defined on \mathbb{R}^{m+2} by*

$$v(x_1, \dots, x_{m+2}) = u(0, x_{m+2})$$

and

$$v(x_1, \dots, x_{m+2}) = u\left(\sqrt{x_1^2 + \dots + x_{m+1}^2}, x_{m+2}\right)$$

is harmonic on $(\mathbb{R}^{m+1})_* \times \mathbb{R}$. In particular, if $m \geq 2$, the Proposition 18 in [15], page 310 shows that v can be extended to a harmonic function on \mathbb{R}^{m+2} , which tends to 0 at infinity. We then deduce that the function v is identically zero, then $u \equiv 0$, demonstrating that the problem to find a solution of $L_m(u) = 0$ with u vanishing at infinity and that the values of u are known on the y -axis is a problem that does not make sense. In this case, no solution to the Dirichlet problem is a consequence of loss of the ellipticity of the equation $L_m u = 0$ at the boundary of \mathbb{H}^+ . Therefore, we do not deal with the case $\operatorname{Re} m \geq 1$.

6. FOURIER-LEGENDRE DECOMPOSITION

First, we will introduce a specific system of coordinates named bipolar coordinates (τ, θ) (see [37]) and numerical applications on extremal bounded problems using this system of coordinates have been realized in [21, 22, 23]. Let $\alpha > 0$. We suppose that there is a positive charge at $A = (-\alpha, 0)$ and a negative charge at $B = (\alpha, 0)$ (the absolute values of the two charges are identical). The potential generated by this charges at a point M is $\ln\left(\frac{MA}{MB}\right)$ (modulo a multiplicative constant).




Figure : Bipolar coordinates

By definition, the bipolar coordinates are

$$\tau := \ln \frac{MA}{MB} \quad \text{and} \quad \theta = \widehat{AMB}.$$

The bipolar coordinates are linked to the Cartesian coordinates by the following formulas :

$$x = \frac{\alpha \operatorname{sh} \tau}{\operatorname{ch} \tau - \cos \theta}, \quad y = \frac{\alpha \sin \theta}{\operatorname{ch} \tau - \cos \theta}.$$

Let $R > 0$ and $a = \sqrt{R^2 + \alpha^2}$, the disk of center $(a, 0)$ and of radius R is defined in terms of bipolar coordinates by

$$\tau \geq \tau_0 = \ln \left(\frac{a}{R} + \sqrt{\frac{a^2}{R^2} - 1} \right) = \operatorname{argch} \frac{a}{R}.$$

Moreover, the right half-plane is

$$\mathbb{H}^+ = \{(\tau, \theta) : \tau \in]0 + \infty], \theta \in [0, 2\pi[\}.$$

The level lines $\tau = \tau_0$ are circles of center $(\alpha \operatorname{coth} \tau_0, 0)$ and radii $\alpha / \operatorname{sh} \tau_0$. This implies that for all τ_0, τ_1 such that $0 < \tau_0 < \tau_1$, the set $\{(\tau, \theta), \tau \geq \tau_0\}$ is a closed disk and the set $\{(\tau, \theta), 0 < \tau < \tau_1\}$ is the complement on \mathbb{H}^+ of the closed disk $\{\tau \geq \tau_1\}$.




Figure : Level lines (with $\alpha = 1$)

The following theorem is known for $m = -1$ by physicists ([3, 44, 42, 43, 33, 39]). We extend this result to complex values of m :

Theorem 6.1. *Let u be a solution of $L_m u = 0$ in an open set of \mathbb{H}^+ and putting*

$$v_m(\tau, \theta) = \operatorname{sh}^{\frac{m-1}{2}} \tau (\operatorname{ch} \tau - \cos \theta)^{-m/2} u(\tau, \theta)$$

where by definition,

$$\operatorname{sh}^{\frac{m-1}{2}} \tau (\operatorname{ch} \tau - \cos \theta)^{-m/2} = \exp \left(\frac{m-1}{2} \ln \operatorname{sh} \tau - \frac{m}{2} \ln (\operatorname{ch} \tau - \cos \theta) \right)$$

then

$$\frac{\partial^2 v_m}{\partial \tau^2} + \frac{\partial^2 v_m}{\partial \theta^2} + \operatorname{coth} \tau \frac{\partial v_m}{\partial \tau} + \left(\frac{1}{4} - \frac{(m-1)^2}{4 \operatorname{sh}^2 \tau} \right) v_m = 0.$$

Proof. We have

$$\frac{\partial u}{\partial \tau} = \alpha \left[\frac{1 - \operatorname{ch} \tau \cos \theta}{(\operatorname{ch} \tau - \cos \theta)^2} \frac{\partial u}{\partial x} - \frac{\operatorname{sh} \tau \sin \theta}{(\operatorname{ch} \tau - \cos \theta)^2} \frac{\partial u}{\partial y} \right]$$

and

$$\frac{\partial u}{\partial \theta} = \alpha \left[\frac{-\operatorname{sh} \tau \sin \theta}{(\operatorname{ch} \tau - \cos \theta)^2} \frac{\partial u}{\partial x} + \frac{\operatorname{ch} \tau \cos \theta - 1}{(\operatorname{ch} \tau - \cos \theta)^2} \frac{\partial u}{\partial y} \right].$$

Thus, we obtain

$$\frac{\partial u}{\partial x} = \frac{1}{\alpha} \left((1 - \operatorname{ch} \tau \cos \theta) \frac{\partial u}{\partial \tau} - \operatorname{sh} \tau \sin \theta \frac{\partial u}{\partial \theta} \right),$$

and

$$\begin{aligned} \frac{\partial^2 u}{\partial \tau^2} &= \frac{\alpha^2}{(\operatorname{ch} \tau - \cos \theta)^4} \left[(1 - \operatorname{ch} \tau \cos \theta)^2 \frac{\partial^2 u}{\partial x^2} + \operatorname{sh}^2 \tau \sin^2 \theta \frac{\partial^2 u}{\partial y^2} \right. \\ &\quad \left. - 2(1 - \operatorname{ch} \tau \cos \theta) \operatorname{sh} \tau \sin \theta \frac{\partial^2 u}{\partial x \partial y} \right] \\ &+ \frac{\alpha}{(\operatorname{ch} \tau - \cos \theta)^3} \left[\operatorname{sh} \tau (\cos^2 \theta + \operatorname{ch} \tau \cos \theta - 2) \frac{\partial u}{\partial x} + \sin \theta (\operatorname{ch}^2 \tau - 2 + \cos \theta \operatorname{ch} \tau) \frac{\partial u}{\partial y} \right] \end{aligned}$$

and

$$\begin{aligned} \frac{\partial^2 u}{\partial \theta^2} &= \frac{\alpha^2}{(\operatorname{ch} \tau - \cos \theta)^4} \left[\operatorname{sh}^2 \tau \sin^2 \theta \frac{\partial^2 u}{\partial x^2} + (\operatorname{ch} \tau \cos \theta - 1) \frac{\partial^2 u}{\partial y^2} \right. \\ &\quad \left. + 2(1 - \operatorname{ch} \tau \cos \theta) \operatorname{sh} \tau \sin \theta \frac{\partial^2 u}{\partial x \partial y} \right] \\ &+ \frac{\alpha}{(\operatorname{ch} \tau - \cos \theta)^3} \left[\operatorname{sh} \tau (2 - \cos^2 \theta - \cos \theta \operatorname{ch} \tau) \frac{\partial u}{\partial x} + \sin \theta (2 - \operatorname{ch}^2 \tau - \operatorname{ch} \tau \cos \theta) \frac{\partial u}{\partial y} \right] \end{aligned}$$

In particular, we have

$$\frac{\partial^2 u}{\partial \tau^2} + \frac{\partial^2 u}{\partial \theta^2} = \frac{\alpha^2}{(\operatorname{ch} \tau - \cos \theta)^2} \left[\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} \right].$$

Therefore, we obtain

$$L_{m,x,y} u = \left(\frac{\operatorname{ch} \tau - \cos \theta}{\alpha} \right)^2 \left(\frac{\partial^2 u}{\partial \tau^2} + \frac{\partial^2 u}{\partial \theta^2} + \frac{m(1 - \operatorname{ch} \tau \cos \theta)}{\operatorname{sh} \tau (\operatorname{ch} \tau - \cos \theta)} \frac{\partial u}{\partial \tau} - \frac{m \sin \theta}{\operatorname{ch} \tau - \cos \theta} \frac{\partial u}{\partial \theta} \right).$$

We put

$$u(\tau, \theta) = \frac{(\operatorname{ch} \tau - \cos \theta)^{m/2}}{\operatorname{sh}^{\frac{m-1}{2}} \tau} v_m(\tau, \theta)$$

and we calculate $L_{m,x,y} u$ in terms of $F(\tau, \theta)$. Denoting

$$r_m(\tau, \theta) = \frac{(\operatorname{ch} \tau - \cos \theta)^{m/2}}{\operatorname{sh}^{\frac{m-1}{2}} \tau},$$

we have

$$\frac{\partial r_m}{\partial \theta} = \frac{m}{2} \frac{\sin \theta}{\operatorname{ch} \tau - \cos \theta} r_m$$

and

$$\frac{\partial^2 r_m}{\partial \theta^2} = \frac{m}{4(\operatorname{ch} \tau - \cos \theta)^2} (2 \cos \theta \operatorname{ch} \tau + m \sin^2 \theta - 2) r_m$$

then

$$\frac{\partial r_m}{\partial \tau} = \frac{1}{(\operatorname{ch} \tau - \cos \theta) \operatorname{sh} \tau} (\operatorname{ch}^2 \tau + (m-1) \operatorname{ch} \tau \cos \theta - m) r_m$$

and

$$\begin{aligned} \frac{\partial^2 r_m}{\partial \tau^2} = & \frac{1}{4(\operatorname{ch} \tau - \cos \theta)^2 \operatorname{sh}^2 \tau} [\operatorname{ch}^4 \tau - 2 \operatorname{ch}^3 \tau \cos \theta + (m-1)^2 \operatorname{ch}^2 \tau \cos^2 \theta + \\ & + 2(m-1) \operatorname{ch}^2 \tau + (4-2m^2) \operatorname{ch} \tau \cos \theta + 2(m-1) \cos^2 \theta + m(m-2)] r_m. \end{aligned}$$

The equation

$$L_{m,x,y} u = 0$$

can be rewritten as

$$\begin{aligned} r_m \left(\frac{\partial^2 v_m}{\partial \tau^2} + \frac{\partial^2 v_m}{\partial \theta^2} \right) + \frac{\partial v_m}{\partial \tau} \left(2 \frac{\partial r_m}{\partial \tau} + \frac{m}{\operatorname{sh} \tau} \frac{1 - \operatorname{ch} \tau \cos \theta}{\operatorname{ch} \tau - \cos \theta} r_m \right) + \\ + \frac{\partial v_m}{\partial \theta} \left(2 \frac{\partial r_m}{\partial \theta} - \frac{m \sin \theta}{\operatorname{ch} \tau - \cos \theta} r_m \right) \\ + v_m \left(\frac{\partial^2 r_m}{\partial \tau^2} + \frac{\partial^2 r_m}{\partial \theta^2} + \frac{m(1 - \operatorname{ch} \tau \cos \theta)}{\operatorname{sh} \tau (\operatorname{ch} \tau - \cos \theta)} \frac{\partial r_m}{\partial \tau} - \frac{m \sin \theta}{\operatorname{ch} \tau - \cos \theta} \frac{\partial r_m}{\partial \theta} \right) = 0 \end{aligned}$$

with

$$\begin{aligned} 2 \frac{\partial r_m}{\partial \tau} + \frac{m}{\operatorname{sh} \tau} \frac{1 - \operatorname{ch} \tau \cos \theta}{\operatorname{ch} \tau - \cos \theta} r_m &= r_m \coth \tau, \\ 2 \frac{\partial r_m}{\partial \theta} - \frac{m \sin \theta}{\operatorname{ch} \tau - \cos \theta} r_m &= 0 \end{aligned}$$

and

$$\frac{\partial^2 r_m}{\partial \tau^2} + \frac{\partial^2 r_m}{\partial \theta^2} + \frac{m(1 - \operatorname{ch} \tau \cos \theta)}{\operatorname{sh} \tau (\operatorname{ch} \tau - \cos \theta)} \frac{\partial r_m}{\partial \tau} - \frac{m \sin \theta}{\operatorname{ch} \tau - \cos \theta} \frac{\partial r_m}{\partial \theta} = \left(\frac{1}{4} - \frac{(m-1)^2}{4 \operatorname{sh}^2 \tau} \right) r_m.$$

And this completes the proof. \square

We seek v_m by separation of variables : $v_m(\tau, \theta) = A_m(\tau)B_m(\theta)$. From the equation satisfied by v_m (see Theorem 6.1), we obtain

$$\frac{A_m''}{A_m} + \coth \tau \frac{A_m'}{A_m} + \frac{1}{4} - \frac{(m-1)^2}{4 \operatorname{sh}^2 \tau} = -\frac{B_m''}{B_m}.$$

The term on the right depends only of θ and the left one depends only of τ , thus we deduce that both are constant. Let $n \in \mathbb{C}$ such that this constant is equal to n^2 . We then have

$$\begin{cases} A_m'' + \coth \tau A_m' + \left(\frac{1}{4} - \frac{(m-1)^2}{4 \operatorname{sh}^2 \tau} - n^2 \right) A_m = 0, \\ B_m'' + n^2 B_m = 0. \end{cases}$$

B_m is naturally a 2π -periodic function (because θ represents an angle), therefore n should necessarily be an integer.

To examine the equation satisfied by A_m , we carry out the following change of function

$$A_m(\tau) = C_m(\operatorname{ch} \tau).$$

Then, C_m satisfies

$$\operatorname{sh}^2 \tau C_m''(\operatorname{ch} \tau) + 2 \operatorname{ch} \tau C_m'(\operatorname{ch} \tau) + \left(\frac{1}{4} - n^2 - \frac{(m-1)^2}{4 \operatorname{sh}^2 \tau} \right) C_m(\operatorname{ch} \tau) = 0$$

which can be rewritten as

$$(1 - \operatorname{ch}^2 \tau) C_m''(\operatorname{ch} \tau) - 2 \operatorname{ch} \tau C_m'(\operatorname{ch} \tau) + \left(n^2 - \frac{1}{4} - \frac{((m-1)/2)^2}{1 - \operatorname{ch}^2 \tau} \right) C_m(\operatorname{ch} \tau) = 0. \quad (LAH)$$

This equation is named *Hyperbolic Associated Legendre equation*.

Note that if we put $z = \operatorname{ch} \tau$ and $u(z) = C_m(\operatorname{ch} \tau)$, then

$$(1 - z^2)u'' - 2zu' + \left[\nu(\nu + 1) - \frac{\mu^2}{1 - z^2} \right] u = 0 \quad (LA)$$

where

$$\nu = n - \frac{1}{2} \quad \text{and} \quad \mu = \frac{m-1}{2}.$$

This equation is named *Associated Legendre equation*, and it can be reduced to the Legendre equation if $\mu = 0$:

$$(1 - z^2)u'' - 2zu' + \nu(\nu + 1)u = 0. \quad (L)$$

Two independent solutions of (LA) are given in section 8 and denoted $P_\nu^\mu(\operatorname{ch} \tau)$ and $Q_\nu^\mu(\operatorname{ch} \tau)$.

Starting from this investigation of solutions in the form of separate variables, we can state the following theorem

Theorem 6.2. *Let $m \in \mathbb{C}$. Let $0 < \tau_0$. Let u be a smooth solution of $L_m u = 0$ on the disk $\tau \geq \tau_0$ and let v be a smooth solution of $L_m v = 0$ on $\mathbb{H}^+ \setminus \{\tau > \tau_0\}$ which is the complement on \mathbb{H}^+ of the disk $\{\tau > \tau_0\}$ and we assume that $\lim_{\partial \mathbb{H}^+} v = 0$. Then there are two sequences $(a_n)_{n \in \mathbb{Z}}$ and $(b_n)_{n \in \mathbb{Z}}$ of $\ell^2(\mathbb{Z})$ (which are even rapidly decreasing) such that :*

$$u = \sum_{n=-\infty}^{+\infty} a_n Q_{n-\frac{1}{2}}^{\frac{m-1}{2}}(\operatorname{ch} \tau) \operatorname{sh}^{\frac{1-m}{2}} \tau (\operatorname{ch} \tau - \cos \theta)^{\frac{m}{2}} e^{in\theta}$$

and

$$v = \sum_{n=-\infty}^{+\infty} b_n P_{n-\frac{1}{2}}^{\frac{m-1}{2}}(\operatorname{ch} \tau) \operatorname{sh}^{\frac{1-m}{2}} \tau (\operatorname{ch} \tau - \cos \theta)^{\frac{m}{2}} e^{in\theta}.$$

The sequence (a_n) is unique. In addition, the convergence of the first series is uniform on every compact set $[\tau_1, \tau_2]$ of the disk $\tau > \tau_0$ with $\tau_0 \leq \tau_1 < \tau_2$. And the convergence of the second one is uniform on every compact set $[\tau_3, \tau_4]$ of the complement of the disk $\tau > \tau_0$ on \mathbb{H}^+ with $0 < \tau_3 < \tau_4 \leq \tau_0$.

If $\operatorname{Re} m < 1$, then the sequence (b_n) is unique.

Proof. Indeed, decomposing the function

$$\theta \mapsto u(\tau_0, \theta)(\operatorname{ch} \tau_0 - \cos \theta)^{-m/2} \operatorname{sh}^{\frac{m-1}{2}} \tau_0$$

by Fourier series with respect to the variable θ , to yield the Fourier expansion for $u(\tau_0, \cdot)$

$$u(\tau_0, \theta) = \operatorname{sh}^{\frac{1-m}{2}} \tau_0 (\operatorname{ch} \tau_0 - \cos \theta)^{\frac{m}{2}} \sum_{n=-\infty}^{+\infty} a_n e^{in\theta},$$

where $a_n \in \ell^2(\mathbb{Z})$ satisfies

$$a_n = \frac{1}{2\pi} \int_0^{2\pi} (\operatorname{ch} \tau_0 - \cos \theta)^{-m/2} \operatorname{sh}^{\frac{m-1}{2}} \tau_0 u(\tau_0, s) e^{-ins} ds.$$

This function is a smooth function of the variable θ , therefore we deduce that the sequence $(a_n)_n$ is rapidly decreasing when $|n| \rightarrow +\infty$. The function

$$\tilde{u}(\tau, \theta) = \operatorname{sh}^{\frac{1-m}{2}} \tau (\operatorname{ch} \tau - \cos \theta)^{\frac{m}{2}} \sum_{n=-\infty}^{+\infty} a_n \frac{Q_{n-\frac{1}{2}}^{\frac{m-1}{2}}(\operatorname{ch} \tau)}{Q_{n-\frac{1}{2}}^{\frac{m-1}{2}}(\operatorname{ch} \tau_0)} e^{in\theta}$$

coincides with u on the circle $\tau = \tau_0$.

Moreover, thanks to the Proposition 8.1, we have when $|n| \rightarrow +\infty$,

$$\frac{Q_{n-\frac{1}{2}}^{\frac{m-1}{2}}(\operatorname{ch} \tau)}{Q_{n-\frac{1}{2}}^{\frac{m-1}{2}}(\operatorname{ch} \tau_0)} \sim \sqrt{\frac{\operatorname{sh} \tau_0}{\operatorname{sh} \tau}} e^{|n|(\tau_0 - \tau)}$$

and this equivalence is uniform in all compact set $[\tau_1, \tau_2]$ with $0 < \tau_0 \leq \tau_1 < \tau_2$.

It follows that the series of functions which defines \tilde{u} is normally converging on any compact sets $[\tau_1, \tau_2]$ of the disk $\tau \geq \tau_0$. So does same for derivatives with respect to τ and θ (which are expressed also with the Associated Legendre functions as mentioned in the section 8).

Particularly, the function \tilde{u} is well defined on the disk $\tau \geq \tau_0$ and coincides with u on the circle $\tau = \tau_0$.

Due to the fact that the solution of an elliptic equation is uniquely determined by its boundary values (this follows from the maximum principle), we deduce that \tilde{u} the unique axisymmetric potential on the disk $\tau \geq \tau_0$ which coincides with u on the circle $\tau = \tau_0$.

For v , the proof is completely similar.

Indeed, decomposing the function

$$\theta \mapsto v(\tau_0, \theta)(\operatorname{ch} \tau_0 - \cos \theta)^{-m/2} \operatorname{sh}^{\frac{m-1}{2}} \tau_0$$

by Fourier series with respect to the variable θ , to yield the Fourier expansion for $v(\tau_0, \cdot)$

$$v(\tau_0, \theta) = \text{sh}^{\frac{1-m}{2}} \tau_0 (\text{ch} \tau_0 - \cos \theta)^{\frac{m}{2}} \sum_{n=-\infty}^{+\infty} b_n e^{in\theta},$$

where $b_n \in \ell^2(\mathbb{Z})$ satisfies

$$b_n = \frac{1}{2\pi} \int_0^{2\pi} (\text{ch} \tau_0 - \cos \theta)^{-m/2} \text{sh}^{\frac{m-1}{2}} \tau_0 v(\tau_0, s) e^{-ins} ds.$$

This function is a smooth function of the variable θ , therefore we deduce that the sequence $(b_n)_n$ is rapidly decreasing when $|n| \rightarrow +\infty$. The function

$$\tilde{v}(\tau, \theta) = \text{sh}^{\frac{1-m}{2}} \tau (\text{ch} \tau - \cos \theta)^{\frac{m}{2}} \sum_{n=-\infty}^{+\infty} b_n \frac{P_{n-\frac{1}{2}}^{\frac{m-1}{2}}(\text{ch} \tau)}{P_{n-\frac{1}{2}}^{\frac{m-1}{2}}(\text{ch} \tau_0)} e^{in\theta}$$

coincides with v on the circle $\tau = \tau_0$.

Moreover, thanks to the Proposition 8.1, we have when $|n| \rightarrow +\infty$,

$$\frac{P_{n-\frac{1}{2}}^{\frac{m-1}{2}}(\text{ch} \tau)}{P_{n-\frac{1}{2}}^{\frac{m-1}{2}}(\text{ch} \tau_0)} \sim \sqrt{\frac{\text{sh} \tau_0}{\text{sh} \tau}} e^{|n|(\tau-\tau_0)}$$

and this equivalence is uniform in all compact set $[\tau_1, \tau_2]$ with $0 < \tau_1 < \tau_2 \leq \tau_0$.

It follows that the series of functions which defines \tilde{v} is normally converging on any compact sets $[\tau_1, \tau_2]$ of the complementary of the disc $\tau > \tau_0$. So does same for derivatives with respect to τ and θ .

Particularly, the function \tilde{v} is well defined on the complementary of the disk $\tau > \tau_0$ and coincides with v on the circle $\tau = \tau_0$.

We will show that

$$\lim_{\tau \rightarrow 0^+} \tilde{v} = 0.$$

If $\text{Re } m < 1$, we have when $n \in \mathbb{N}$ and thanks to the formula (8.1)

$$P_{n-\frac{1}{2}}^{\frac{m-1}{2}}(\text{ch} \tau) = \frac{2^{\frac{m-1}{2}}}{\sqrt{\pi} \Gamma(1 - \frac{m}{2})} \text{sh}^{\frac{1-m}{2}} \tau \int_0^\pi (\text{ch} \tau + \text{sh} \tau \cos \theta)^{n+\frac{m}{2}-1} \sin^{1-m} \theta d\theta$$

then

$$\lim_{\tau \rightarrow 0^+} P_{n-\frac{1}{2}}^{\frac{m-1}{2}}(\text{ch} \tau) = 0$$

and in addition, for $n > 1 - \frac{\text{Re } m}{2}$, we have

$$\begin{aligned} \left| P_{n-\frac{1}{2}}^{\frac{m-1}{2}}(\text{ch} \tau) \right| &\leq \frac{2^{\frac{\text{Re } m-1}{2}} \text{sh}^{\frac{1-\text{Re } m}{2}} \tau}{\sqrt{\pi} |\Gamma(1 - \frac{m}{2})|} \int_0^\pi (\text{ch} \tau + \text{sh} \tau \cos \theta)^{n+\frac{\text{Re } m}{2}-1} \sin^{1-\text{Re } m} \theta d\theta \\ &\leq \frac{2^{\frac{\text{Re } m-1}{2}} \text{sh}^{\frac{1-\text{Re } m}{2}} \tau}{\sqrt{\pi} |\Gamma(1 - \frac{m}{2})|} \int_0^\pi (\text{ch} \tau + \text{sh} \tau)^{n+\frac{\text{Re } m}{2}-1} \sin^{1-\text{Re } m} \theta d\theta \leq C_m \text{sh}^{\frac{1-\text{Re } m}{2}} \tau e^{(n+\frac{\text{Re } m}{2})\tau} \end{aligned}$$

thus

$$\sum_{n > 1 - \frac{\operatorname{Re} m}{2}} \sup_{\tau \in [0, \frac{\tau_0}{2}]} \left| b_n \frac{P_{n-\frac{1}{2}}^{\frac{m-1}{2}}(\operatorname{ch} \tau)}{P_{n-\frac{1}{2}}^{\frac{m-1}{2}}(\operatorname{ch} \tau_0)} e^{in\theta} \right| < +\infty$$

by the Proposition 8.1, we obtain

$$P_{n-\frac{1}{2}}^{\frac{m-1}{2}}(\operatorname{ch} \tau_0) \sim_{n \rightarrow +\infty} \frac{n^{\frac{m}{2}-1}}{\sqrt{2\pi \operatorname{sh} \tau_0}} e^{n\tau_0}.$$

So, we can deduce that $\lim_{\tau \rightarrow 0+} \tilde{v} = 0$.

It remains to prove the uniqueness of the previous decomposition where $\operatorname{Re} m < 1$. This will result in the next paragraph which will establish the fact that the family

$$\begin{aligned} \mathcal{A} &:= \left(\frac{Q_{n-\frac{1}{2}}^{\frac{m-1}{2}}(\operatorname{ch} \tau)}{Q_{n-\frac{1}{2}}^{\frac{m-1}{2}}(\operatorname{ch} \tau_0)} \frac{(\operatorname{ch} \tau - \cos \theta)^{m/2}}{\operatorname{sh}^{\frac{m-1}{2}} \tau} e^{in\theta} \right)_{n \in \mathbb{Z}} := (a_n)_{n \in \mathbb{Z}} \\ \mathcal{B} &:= \left(\frac{P_{n-\frac{1}{2}}^{\frac{m-1}{2}}(\operatorname{ch} \tau)}{P_{n-\frac{1}{2}}^{\frac{m-1}{2}}(\operatorname{ch} \tau_1)} \frac{(\operatorname{ch} \tau - \cos \theta)^{m/2}}{\operatorname{sh}^{\frac{m-1}{2}} \tau} e^{in\theta} \right)_{n \in \mathbb{Z}} := (b_n)_{n \in \mathbb{Z}} \end{aligned}$$

is a Riesz basis. \square

Corollary 6.3. *The solution of the Dirichlet problem for $L_m u = 0$ on $D((a, 0), R)$ where $u = \varphi$ on $\partial D((a, 0), R)$ is given by*

$$u(\tau, \theta) = \operatorname{sh}^{\frac{1-m}{2}} \tau (\operatorname{ch} \tau - \cos \theta)^{\frac{m}{2}} \sum_{n=-\infty}^{+\infty} c_n \frac{Q_{n-\frac{1}{2}}^{\frac{m-1}{2}}(\operatorname{ch} \tau)}{Q_{n-\frac{1}{2}}^{\frac{m-1}{2}}(\operatorname{ch} \tau_0)} e^{in\theta}$$

where $\{\tau = \tau_0\}$ corresponds to the circle of center $(a, 0)$ and radius R and where

$$c_n = \frac{1}{2\pi} \int_0^{2\pi} (\operatorname{ch} \tau_0 - \cos \theta)^{-m/2} \operatorname{sh}^{\frac{m-1}{2}} \tau_0 \varphi(a + R \cos s, R \sin s) e^{-ins} ds.$$

Similarly,

$$v(\tau, \theta) = \operatorname{sh}^{\frac{1-m}{2}} \tau (\operatorname{ch} \tau - \cos \theta)^{\frac{m}{2}} \sum_{n=-\infty}^{+\infty} c_n \frac{P_{n-\frac{1}{2}}^{\frac{m-1}{2}}(\operatorname{ch} \tau)}{P_{n-\frac{1}{2}}^{\frac{m-1}{2}}(\operatorname{ch} \tau_0)} e^{in\theta}$$

is a solution of $L_m v = 0$ on $\mathbb{H}^+ \setminus D((a, 0), R)$, which is equal to φ on $\partial D((a, 0), R)$ where

$$c_n = \frac{1}{2\pi} \int_0^{2\pi} (\operatorname{ch} \tau_0 - \cos \theta)^{-m/2} \operatorname{sh}^{\frac{m-1}{2}} \tau_0 \varphi(a + R \cos s, R \sin s) e^{-ins} ds.$$

Moreover, if $Rem < 1$, then v satisfies $\lim_{\partial\mathbb{H}^+} v = 0$, and the function v constructed above is the unique solution of the Dirichlet problem $L_m v = 0$ on $\mathbb{H}^+ \setminus D((a, 0), R)$ which vanishes on $\partial\mathbb{H}^+$.

7. RIESZ BASIS

We will prove that the half part of the following family

$$\left(\frac{(\operatorname{ch} \tau - \cos \theta)^{m/2}}{\operatorname{sh} \frac{m-1}{2} \tau} \begin{Bmatrix} \cos(n\theta) \\ \sin(n\theta) \end{Bmatrix} \begin{Bmatrix} P_{n-\frac{1}{2}}^{\frac{m-1}{2}}(\operatorname{ch} \tau) \\ Q_{n-\frac{1}{2}}^{\frac{m-1}{2}}(\operatorname{ch} \tau) \end{Bmatrix} \right)_{n \in \mathbb{Z}}$$

is a basis of solutions on the disk $\tau \geq \tau_1$ and the other half part is a basis of solutions on $\tau \leq \tau_0$, which is the complement on \mathbb{H}^+ of a some disk, with $0 < \tau_0 < \tau_1$. This fact is known for $m = -1$, namely for $\mu = 1$. We extend this result for complex values of m .

Let us recall the definition of a Riesz basis. $(x_n)_{n \in \mathbb{N}}$ is a quasi-orthogonal or Riesz sequence of a Hilbert space X if there are two constants $c, C > 0$ such that for all sequences $(a_n)_{n \in \mathbb{Z}}$ in ℓ^2 , we have

$$c^2 \sum_n |a_n|^2 \leq \left\| \sum_n a_n x_n \right\|^2 \leq C^2 \sum_n |a_n|^2.$$

If the family $(x_n)_{n \in \mathbb{Z}}$ is complete, it is a *Riesz basis*. The matrix of scalar product $\{\langle x_i, x_j \rangle\}_{i,j}$ is named Gram matrix associated to $\{x_i\}_i$.

To prove that $\{x_i\}_i$ is a Riesz basis, a convenient characterization with the Gram matrix is the following property :

Property ([40, p. 170]). *A family $\{x_i\}_i$ is a Riesz basis for a some Hilbert space if $\{x_i\}_i$ is complete on this Hilbert space and if the Gram matrix associated to $\{x_i\}_i$ defines a bounded and invertible operator on $\ell^2(\mathbb{N})$.*

Let \mathcal{A} and \mathcal{B} the two families of solutions of $L_m[u] = 0$, respectively inside the disk $\tau > \tau_0$ and outside the other one $\tau > \tau_1$, with $0 < \tau_0 < \tau_1$

$$\mathcal{A} := \left(\frac{Q_{n-\frac{1}{2}}^{\frac{m-1}{2}}(\operatorname{ch} \tau)}{Q_{n-\frac{1}{2}}^{\frac{m-1}{2}}(\operatorname{ch} \tau_0)} \frac{(\operatorname{ch} \tau - \cos \theta)^{m/2}}{\operatorname{sh} \frac{m-1}{2} \tau} e^{in\theta} \right)_{n \in \mathbb{Z}} := (a_n)_{n \in \mathbb{Z}}$$

$$\mathcal{B} := \left(\frac{P_{n-\frac{1}{2}}^{\frac{m-1}{2}}(\operatorname{ch} \tau)}{P_{n-\frac{1}{2}}^{\frac{m-1}{2}}(\operatorname{ch} \tau_1)} \frac{(\operatorname{ch} \tau - \cos \theta)^{m/2}}{\operatorname{sh} \frac{m-1}{2} \tau} e^{in\theta} \right)_{n \in \mathbb{Z}} := (b_n)_{n \in \mathbb{Z}}$$

Let \mathcal{C} the union of the two previous families :

$$\mathcal{C} := (c_n)_{n \in \mathbb{Z}} := (c_{2n} = a_n \text{ et } c_{2n+1} = b_n)_{n \in \mathbb{Z}}$$

The annulus defined in terms of bipolar coordinates $\{0 < \tau_0 < \tau < \tau_1\}$ will be denoted \mathbb{A} . The space $L^2(\partial\mathbb{A})$ is equipped of the following inner product : for $f, g \in L^2(\partial\mathbb{A})$,

$$\begin{aligned} \langle f, g \rangle &= \frac{1}{2\pi} \int_0^{2\pi} f(\tau_0, \theta) \overline{g(\tau_0, \theta)} \frac{\text{sh}^{\text{Re } m-1} \tau_0}{(\text{ch } \tau_0 - \cos \theta)^{\text{Re } m}} d\theta \\ &\quad + \frac{1}{2\pi} \int_0^{2\pi} f(\tau_1, \theta) \overline{g(\tau_1, \theta)} \frac{\text{sh}^{\text{Re } m-1} \tau_1}{(\text{ch } \tau_1 - \cos \theta)^{\text{Re } m}} d\theta. \end{aligned}$$

We have the following proposition :

Proposition 7.1. \mathcal{C} is a Riesz basis in the Hilbert space $L^2(\partial\mathbb{A})$.

Proof. Indeed, in order to build the Gram matrix of \mathcal{C} , we first calculate all its scalar products. We obtain for all $n \in \mathbb{Z}$,

$$\begin{aligned} \langle c_{2n}, c_{2n} \rangle &= 1 + \left| \frac{Q_{n-\frac{1}{2}}^{\frac{m-1}{2}}(\text{ch } \tau_1)}{Q_{n-\frac{1}{2}}^{\frac{m-1}{2}}(\text{ch } \tau_0)} \right|^2 \\ \langle c_{2n+1}, c_{2n+1} \rangle &= 1 + \left| \frac{P_{n-\frac{1}{2}}^{\frac{m-1}{2}}(\text{ch } \tau_0)}{P_{n-\frac{1}{2}}^{\frac{m-1}{2}}(\text{ch } \tau_1)} \right|^2 \\ \langle c_{2n}, c_{2n+1} \rangle &= \overline{\left(\frac{P_{n-\frac{1}{2}}^{\frac{m-1}{2}}(\text{ch } \tau_0)}{P_{n-\frac{1}{2}}^{\frac{m-1}{2}}(\text{ch } \tau_1)} \right)} + \frac{Q_{n-\frac{1}{2}}^{\frac{m-1}{2}}(\text{ch } \tau_1)}{Q_{n-\frac{1}{2}}^{\frac{m-1}{2}}(\text{ch } \tau_0)} \\ \langle c_{2n+1}, c_{2n} \rangle &= \overline{\langle c_{2n}, c_{2n+1} \rangle} \end{aligned}$$

In all other cases, the inner product is zero, the Gram matrix is diagonal by blocks and each blocks is expressed as the 2×2 matrix :

$$M_n = \begin{pmatrix} 1 + \left| \frac{Q_{n-\frac{1}{2}}^{\frac{m-1}{2}}(\text{ch } \tau_1)}{Q_{n-\frac{1}{2}}^{\frac{m-1}{2}}(\text{ch } \tau_0)} \right|^2 & \overline{\left(\frac{P_{n-\frac{1}{2}}^{\frac{m-1}{2}}(\text{ch } \tau_0)}{P_{n-\frac{1}{2}}^{\frac{m-1}{2}}(\text{ch } \tau_1)} \right)} + \frac{Q_{n-\frac{1}{2}}^{\frac{m-1}{2}}(\text{ch } \tau_1)}{Q_{n-\frac{1}{2}}^{\frac{m-1}{2}}(\text{ch } \tau_0)} \\ \frac{P_{n-\frac{1}{2}}^{\frac{m-1}{2}}(\text{ch } \tau_0)}{P_{n-\frac{1}{2}}^{\frac{m-1}{2}}(\text{ch } \tau_1)} + \overline{\left(\frac{Q_{n-\frac{1}{2}}^{\frac{m-1}{2}}(\text{ch } \tau_1)}{Q_{n-\frac{1}{2}}^{\frac{m-1}{2}}(\text{ch } \tau_0)} \right)} & 1 + \left| \frac{P_{n-\frac{1}{2}}^{\frac{m-1}{2}}(\text{ch } \tau_0)}{P_{n-\frac{1}{2}}^{\frac{m-1}{2}}(\text{ch } \tau_1)} \right|^2 \end{pmatrix}$$

The Gram matrix G can be written as

$$G = \begin{pmatrix} M_0 & 0 & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ 0 & M_{-1} & 0 & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ \vdots & 0 & M_1 & 0 & \cdots & \cdots & \cdots & \cdots & \cdots \\ \vdots & \ddots & 0 & M_{-2} & 0 & \cdots & \cdots & \cdots & \cdots \\ \vdots & \ddots & \ddots & 0 & \ddots & \ddots & \cdots & \cdots & \cdots \\ \vdots & \ddots & \ddots & \ddots & \ddots & M_{-n} & \ddots & \cdots & \cdots \\ \vdots & \ddots & \ddots & \ddots & \ddots & \ddots & M_n & \ddots & \cdots \\ \vdots & \ddots & \ddots & \ddots & \ddots & \ddots & \ddots & M_n & \ddots \\ \vdots & \ddots & \ddots & \ddots & \ddots & \ddots & \ddots & \ddots & \ddots \end{pmatrix}$$

and the determinant of M_n is

$$\det(M_n) = \left| 1 - \frac{Q_{n-\frac{1}{2}}^{\frac{m-1}{2}}(\text{ch } \tau_1) P_{n-\frac{1}{2}}^{\frac{m-1}{2}}(\text{ch } \tau_0)}{Q_{n-\frac{1}{2}}^{\frac{m-1}{2}}(\text{ch } \tau_0) P_{n-\frac{1}{2}}^{\frac{m-1}{2}}(\text{ch } \tau_1)} \right|^2$$

Let's show that M_n is invertible. Suppose the contrary, if M_n is not invertible, then $\det(M_n) = 0$, which is equivalent to

$$Q_{n-\frac{1}{2}}^{\frac{m-1}{2}}(\text{ch } \tau_1) P_{n-\frac{1}{2}}^{\frac{m-1}{2}}(\text{ch } \tau_0) = Q_{n-\frac{1}{2}}^{\frac{m-1}{2}}(\text{ch } \tau_0) P_{n-\frac{1}{2}}^{\frac{m-1}{2}}(\text{ch } \tau_1).$$

The previous equality can be written as follows

$$\begin{vmatrix} Q_{n-\frac{1}{2}}^{\frac{m-1}{2}}(\text{ch } \tau_0) & P_{n-\frac{1}{2}}^{\frac{m-1}{2}}(\text{ch } \tau_0) \\ Q_{n-\frac{1}{2}}^{\frac{m-1}{2}}(\text{ch } \tau_1) & P_{n-\frac{1}{2}}^{\frac{m-1}{2}}(\text{ch } \tau_1) \end{vmatrix} = 0, \text{ with } P_{n-\frac{1}{2}}^{\frac{m-1}{2}}(\text{ch } \tau_0), Q_{n-\frac{1}{2}}^{\frac{m-1}{2}}(\text{ch } \tau_1) \neq 0.$$

Therefore, there is $\lambda \in \mathbb{C} \setminus \{0\}$ (which depends on m, n, τ_0 and τ_1) such that

$$\begin{cases} Q_{n-\frac{1}{2}}^{\frac{m-1}{2}}(\text{ch } \tau_0) = \lambda P_{n-\frac{1}{2}}^{\frac{m-1}{2}}(\text{ch } \tau_0) \\ Q_{n-\frac{1}{2}}^{\frac{m-1}{2}}(\text{ch } \tau_1) = \lambda P_{n-\frac{1}{2}}^{\frac{m-1}{2}}(\text{ch } \tau_1) \end{cases}$$

Then, by the asymptotic of Associated Legendre functions (see Proposition 8.1 in the Annex), on the one hand, we have both

$$\lambda \underset{n \rightarrow +\infty}{\sim} \pi e^{i\pi \frac{m-1}{2}} e^{-2n\tau_0}$$

and on the other hand, we have

$$\lambda \underset{n \rightarrow +\infty}{\sim} \pi e^{i\pi \frac{m-1}{2}} e^{-2n\tau_1},$$

then it implies that $\tau_0 = \tau_1$, it is not possible.

We deduce that the matrix M_n is invertible and this completes the proof. \square

8. ANNEX : ASSOCIATED LEGENDRE FUNCTIONS OF FIRST AND SECOND KIND

In this section, we provide the main formulas of integral representation for the Associated Legendre function of the first and the second kind with $z = \text{ch } \tau > 1$ (see [2, 37, 46]) :

$$P_{\nu}^{\mu}(\text{ch } \tau) = \frac{2^{-\nu} \text{sh }^{-\mu} \tau}{\Gamma(-\mu - \nu) \Gamma(\nu + 1)} \int_0^{\infty} (\text{ch } \tau + \text{ch } \theta)^{\mu - \nu - 1} \text{sh }^{2\nu + 1} \theta \, d\theta$$

with $\text{Re } \nu > -1$ and $\text{Re}(\mu + \nu) < 0$.

$$P_{\nu}^{\mu}(\text{ch } \tau) = \frac{2^{\mu} \text{sh }^{-\mu} \tau}{\sqrt{\pi} \Gamma(\frac{1}{2} - \mu)} \int_0^{\pi} \frac{(\text{ch } \tau + \text{sh } \tau \cos \theta)^{\mu + \nu}}{\sin^{2\mu} \theta} d\theta \quad (8.1)$$

with $\text{Re } \mu < \frac{1}{2}$.

$$P_{\nu}^{\mu}(\text{ch } \tau) = \sqrt{\frac{2}{\pi}} \frac{\text{sh }^{\mu} \tau}{\Gamma(\frac{1}{2} - \mu)} \int_0^{\tau} \frac{\text{ch } [(\nu + \frac{1}{2}) \theta]}{(\text{ch } \tau - \text{ch } \theta)^{\mu + 1/2}} d\theta$$

with $\text{Re } \mu < \frac{1}{2}$.

$$Q_{\nu}^{\mu}(\text{ch } \tau) = \frac{e^{i\pi\mu} \sqrt{\pi}}{2^{\mu}} \frac{\text{sh }^{\mu} \tau \Gamma(\nu + \mu + 1)}{\Gamma(\nu - \mu + 1) \Gamma(\mu + 1/2)} \int_0^{\infty} \frac{\text{sh }^{2\mu} \theta}{(\text{ch } \tau + \text{sh } \tau \text{ch } \theta)^{\nu + \mu + 1}} d\theta$$

with $\text{Re } \mu > -\frac{1}{2}$, $\text{Re}(\mu - \nu - 1) < 0$ and $\mu + \nu + 1 \notin \mathbb{Z}^{-}$.

$$Q_{\nu}^{\mu}(\text{ch } \tau) = \sqrt{\frac{\pi}{2}} e^{i\pi\mu} \frac{\text{sh }^{\mu} \tau}{\Gamma(\frac{1}{2} - \mu)} \int_{\tau}^{\infty} \frac{e^{-(\nu + \frac{1}{2})\theta}}{(\text{ch } \theta - \text{ch } \tau)^{\mu + 1/2}} d\theta$$

with $\text{Re } \mu < \frac{1}{2}$ et $\text{Re}(\mu + \nu + 1) > 0$.

$$Q_{\nu}^{\mu}(\text{ch } \tau) = e^{i\pi\mu} 2^{-\nu - 1} \frac{\Gamma(\nu + \mu + 1)}{\Gamma(\nu + 1)} \text{sh }^{-\mu} \tau \int_0^{\pi} (\text{ch } \tau + \cos \theta)^{\mu - \nu - 1} \sin^{2\nu + 1} \theta \, d\theta$$

with $\text{Re } \nu > -1$ and $\mu + \nu + 1 \notin \mathbb{Z}^{-}$ (see [46] pages 4, 5 and 6).

We have also the following relations satisfied by the Legendre functions (see [46] page 6 and [2], formula 8.2.2)

$$P_{\nu}^{\mu} = P_{-\nu - 1}^{\mu}.$$

$$Q_{-\nu - 1}^{\mu}(z) = \frac{-\pi e^{i\pi\mu} \cos(\pi\nu) P_{\nu}^{\mu} + \sin[\pi(\nu + \mu)] Q_{\nu}^{\mu}}{\sin[\pi(\nu - \mu)]}$$

for $\nu - \mu \notin \mathbb{Z}$. (in particular for $\nu = n - \frac{1}{2}$ with $n \in \mathbb{Z}$, we have

$$Q_{-\nu - 1}^{\mu} = Q_{\nu}^{\mu}$$

for all $\mu \in \mathbb{C}$),

$$e^{i\pi\mu} \Gamma(\nu + \mu + 1) Q_{\nu}^{-\mu} = e^{-i\pi\mu} \Gamma(\nu - \mu + 1) Q_{\nu}^{\mu},$$

$$P_\nu^{-\mu} = \frac{\Gamma(\nu - \mu + 1)}{\Gamma(\nu + \mu + 1)} \left[P_\nu^\mu - \frac{2}{\pi} e^{-i\pi\mu} \sin(\pi\mu) Q_\nu^\mu \right],$$

In addition, we have the Whipple formulas connecting the associated Legendre functions of first and second kind (see [46] page 6)

$$Q_\nu^\mu(\operatorname{ch} \tau) = e^{i\pi\mu} \sqrt{\frac{\pi}{2}} \frac{\Gamma(\mu + \nu + 1)}{\sqrt{\operatorname{sh} \tau}} P_{-\mu - \frac{1}{2}}^{-\nu - \frac{1}{2}}(\operatorname{coth} \tau),$$

$$P_\nu^\mu(\operatorname{ch} \tau) = \frac{ie^{i\pi\nu}}{\Gamma(-\nu - \mu)} \sqrt{\frac{2}{\pi}} \frac{1}{\sqrt{\operatorname{sh} \tau}} Q_{-\mu - \frac{1}{2}}^{-\nu - \frac{1}{2}}(\operatorname{coth} \tau).$$

We also have the recursion formulas (see [46] pages 6 et 7)

$$P_\nu^{\mu+1}(\operatorname{ch} \tau) = \frac{(\nu - \mu) \operatorname{ch} \tau P_\nu^\mu(\operatorname{ch} \tau) - (\nu + \mu) P_{\nu-1}^\mu(\operatorname{ch} \tau)}{\operatorname{sh} \tau}$$

$$(\nu - \mu + 1) P_{\nu+1}^\mu(\operatorname{ch} \tau) = (2\nu + 1) \operatorname{ch} \tau P_\nu^\mu(\operatorname{ch} \tau) - (\nu + \mu) P_{\nu-1}^\mu(\operatorname{ch} \tau).$$

$$(z^2 - 1) \frac{dP_\nu^\mu(z)}{dz} = (\nu + \mu)(\nu - \mu + 1)(z^2 - 1)^{1/2} P_\nu^{\mu-1}(z) - \mu z P_\nu^\mu(z).$$

$$(z^2 - 1) \frac{dP_\nu^\mu(z)}{dz} = \nu z P_\nu^\mu(z) - (\nu + \mu) P_{\nu-1}^\mu(z).$$

All of these formulas are used to explicitly calculate the values of $P_\nu^\mu(\operatorname{ch} \tau)$ and $Q_\nu^\mu(\operatorname{ch} \tau)$ for all $\tau > 0$ and $(\mu, \nu) \in \mathbb{C}^2$.

If μ and τ are fixed, the following proposition collects the behavior of Associated Legendre functions of the first and second kind when $\nu = n - \frac{1}{2}$ with $n \in \mathbb{Z}$ and $|n| \rightarrow +\infty$.

Proposition 8.1. *We fix $\tau > 0$ and $\mu \in \mathbb{C}$. Then if $\nu = n - \frac{1}{2}$ with $n \in \mathbb{Z}$, we have :*

$$\text{when } \nu \rightarrow +\infty, \quad P_\nu^\mu(\operatorname{ch} \tau) \sim \frac{e^{\tau/2}}{\sqrt{2\pi \operatorname{sh} \tau}} \nu^{\mu-1/2} e^{\tau\nu}$$

$$\text{when } \nu \rightarrow -\infty, \quad P_\nu^\mu(\operatorname{ch} \tau) \sim \frac{e^{-\tau/2}}{\sqrt{2\pi \operatorname{sh} \tau}} (-\nu)^{\mu-1/2} e^{-\tau\nu}$$

$$\text{when } \nu \rightarrow +\infty, \quad Q_\nu^\mu(\operatorname{ch} \tau) \sim e^{i\pi\mu} e^{-\tau/2} \sqrt{\frac{\pi}{2 \operatorname{sh} \tau}} \nu^{\mu-1/2} e^{-\tau\nu}$$

$$\text{when } \nu \rightarrow -\infty, \quad Q_\nu^\mu(\operatorname{ch} \tau) \sim e^{i\pi\mu} e^{\tau/2} \sqrt{\frac{\pi}{2 \operatorname{sh} \tau}} (-\nu)^{\mu-1/2} e^{\tau\nu}.$$

These equivalences are locally uniform with respect to the variable τ , that is to say uniform on all interval $[\tau_0, \tau_1]$ with $0 < \tau_0 < \tau_1$.

Proof. If $\nu = n - \frac{1}{2}$ with $n \in \mathbb{N}$ (see [46] page 48), we have

$$P_\nu^\mu(\operatorname{ch} \tau) = \frac{\Gamma(\nu + 1)}{\Gamma(\nu - \mu + 1)} \frac{1}{\sqrt{2\pi(\nu + 1)\operatorname{sh} \tau}} \left[e^{(\nu + \frac{1}{2})\tau} + e^{-\pi i(\mu - \frac{1}{2}) - (\nu + \frac{1}{2})\tau} \right] \left[1 + \mathcal{O}\left(\frac{1}{\nu}\right) \right].$$

A straightforward application of the Stirling formula shows that when $\nu \rightarrow +\infty$

$$\begin{aligned} \frac{\Gamma(\nu+1)}{\Gamma(\nu-\mu+1)} &\sim \frac{\sqrt{2\pi}\nu^{\nu+1/2}e^{-\nu}}{\sqrt{2\pi}(\nu-\mu)^{\nu-\mu+1/2}e^{-\nu+\mu}} = \left(\frac{\nu}{\nu-\mu}\right)^{\nu+1/2} (\nu-\mu)^\mu e^{-\mu} \\ &= (\nu-\mu)^\mu e^{-\mu} \exp\left(-\left(\nu+\frac{1}{2}\right)\ln\left(1-\frac{\mu}{\nu}\right)\right) \sim \nu^\mu \end{aligned}$$

consequently,

$$P_\nu^\mu(\operatorname{ch} \tau) \sim \nu^\mu \frac{1}{\sqrt{2\pi\nu\operatorname{sh} \tau}} e^{\frac{\tau}{2}} e^{\tau\nu} = \frac{e^{\tau/2}}{\sqrt{2\pi\operatorname{sh} \tau}} \nu^{\mu-1/2} e^{\tau\nu},$$

which gives us the first estimate.

The second one is obtained directly thanks to the relation $P_\nu^\mu = P_{-\nu-1}^\mu$. The third estimate follows directly from the formula (8.3) of [46] :

$$Q_\nu^\mu(\operatorname{ch} \tau) \sim \sqrt{\frac{\pi}{2\operatorname{sh} \tau}} \nu^{\mu-1/2} e^{i\pi\mu} e^{-\tau(\nu+1/2)}$$

and the last estimation arises from the fact that for $\nu = n - \frac{1}{2}$ with $n \in \mathbb{Z}$, we have

$$Q_{-\nu-1}^\mu = Q_\nu^\mu.$$

The locally uniform character of these equivalences come from explicit expressions of P_ν^μ and Q_ν^μ in terms of hypergeometric functions ([17], tables pages 124-138) and estimations of these special functions (always locally uniform with respect to their parameters ([46], pages 178-182)). \square

REFERENCES

- [1] M. J. Ablowitz and A. S. Fokas. *Complex Variables : Introduction and Applications*. Cambridge Texts in Applied Mathematics. Cambridge University Press, Cambridge, 1997.
- [2] M. Abramowitz and Irene A. Stegun. *Handbook of mathematical functions*. Dover, 1972.
- [3] F. Alladio and F. Crisanti. Analysis of mhd equilibria by toroidal multipolar expansions,. *Nuclear Fusion*, 26:1143–1164, 1986.
- [4] L. Baratchart, A. Borichev, and S. Chaabi. Pseudo-holomorphic functions at the critical exponent.
- [5] L. Baratchart, Y. Fischer, and J. Leblond. Dirichlet/Neumann problems and Hardy classes for the planar conductivity equation. *to appear in Complex variables and elliptic equations*.
- [6] L. Baratchart, J. Leblond, S. Rigat, and E. Russ. Hardy spaces of the conjugated Beltrami equation. *Journal of Functional Analysis*, 259:384–427, 2010.
- [7] M. Brelot. Équation de Weinstein et potentiels de Marcel Riesz. *Lecture Notes in Math, Springer*, 681:1838, 1978.
- [8] B. Brelot-Collin and M. Brelot. Représentation intégrale des solutions positives de l'équation

$$L_k(u) = \sum_1^n \frac{\partial^2 u}{\partial x_i^2} + \frac{k}{x_n} \frac{\partial u}{\partial x_n} = 0$$

(k constante réelle) dans le demi-espace $E(x_n > 0)$, de \mathbb{R}^n . *Acad. Roy. Belg. Bull. Cl. Sci. (5)*, 58:317–326., 1972.

- [9] B. BreLOT-COLLIN and M. BreLOT. Allure à la frontière des solutions positives de l'équation de weinstein

$$L_k(u) = \Delta u + kx_n^{-1} \frac{\partial u}{\partial x_n} = 0$$

dans le demi-espace $E(x_n > 0)$ de \mathbb{R}^n ($n \geq 2$). *Acad. Roy. Belg. Bull. Cl. Sci. (5)*, 59:1100–1117, 1973.

- [10] B. BreLOT-COLLIN and M. BreLOT. tude la frontire des solutions locales positives de l'equation

$$(1) \quad L_k(u) = \Delta u + kx_n^{-1} \frac{\partial u}{\partial x_n} = 0$$

dans le demi-espace $E(x_n > 0)$ de \mathbb{R}^n ($n \geq 2$). *Acad. Roy. Belg. Bull. Cl. Sci. (5)*, 62:322340, 1976.

- [11] Slah Chaabi. *Analyse complexe et problèmesmes de Dirichlet dans le plan : équation de Weinstein et autres conductivités non-bornées*. PhD thesis, Aix-Marseille université, 2013.
- [12] E. T. Copson. On sound waves of finite amplitude. *Proc. Roy. Soc. London. Ser. A.*, 216:539–547, 1953.
- [13] E. T. Copson. On hadamard's elementary solution. (1970/71), 1927. *Proc. Roy. Soc. Edinburgh Sect. A*, 69:19–27, 1970/71.
- [14] E. T. Copson. *Partial differential equations*. Cambridge University Press, Cambridge-New York-Melbourne, 1975.
- [15] R. Dautray and J-L Lions. *Analyse mathématique et calcul numérique pour les sciences et techniques, L'opérateur de Laplace, Tome 2*. Masson, 1987.
- [16] J. B. Diaz and A. Weinstein. On the fundamental solutions of a singular beltrami operator. *Studies in mathematics and mechanics presented to Richard von Mises, Academic Press Inc*, pages 97–102, 1954.
- [17] A. Erdélyi. *Higher Transcendental Functions, 1*. McGraw-Hill, Newyork, Toronto, London, 1953.
- [18] A. Erdélyi. Singularities of generalized axially symmetric potentials. *Comm. Pure Appl. Math.*, 9:403–414, 1956.
- [19] A. Erdélyi. An application of fractional integrals. *J. Analyse Math.* 14, 14:113–126, 1965.
- [20] A. Erdélyi. Symmetric potentials and fractional integration. *J. of the Soc. for Ind. and Applied Math*, Vol. 13, no 1:216–228, 1965.
- [21] Y. Fischer and J. Leblond. Solutions to conjugate Beltrami equations and approximation in generalized Hardy spaces. *Adv. Pure Appl. Math.*, 2(1):47–63, 2011.
- [22] Yannick Fischer, Juliette Leblond, Jonathan R. Partington, and Eva Sincich. Bounded extremal problems in Hardy spaces for the conjugate Beltrami equation in simply-connected domains. *Appl. Comput. Harmon. Anal.*, 31(2):264–285, 2011.
- [23] Yannick Fischer, Benjamin Marteau, and Yannick Privat. Some inverse problems around the Tokamak *tore supra*. *Commun. Pure Appl. Anal.*, 11(6):2327–2349, 2012.
- [24] A. S. Fokas and J. Lenells. Boundary-value problems for the stationary axisymmetric einstein equations: a rotating disc. *Nonlinearity*, 24:177–206, 2011.
- [25] P. R. Garabedian. *Partial Differential Equations*. AMS Chelsea Publishing, Providence, RI, Reprint of the 1964 original, 1998.
- [26] R. P. Gilbert. Some properties of generalized axially symmetric potentials. *Amer. J. Math.*, 84:475–484, 1962.
- [27] R. P. Gilbert. On generalized axially symmetric potentials. *J. Reine Angew. Math.*, 212:158–168, 1963.
- [28] R. P. Gilbert. Poisson's equation and generalized axially potential theory. *Ann. Mat. Pura Appl.*, 61:337–348, 1963.

- [29] R. P. Gilbert. Bergman's integral operator method in generalized axially symmetric potential theory. *J. Math. Physics*, 5:983–997, 1964.
- [30] D. W. Weinacht R. J. Hall, N. S. Quinn. Poisson integral formulas in generalized bi-axially symmetric potential theory. *SIAM J. Math. Anal.*, 5:111–118, 1974.
- [31] P. Henrici. On the domain of regularity of generalized axially symmetric potentials. *Proc. Amer. Math. Soc.*, 8:29–31, 1957.
- [32] Lars Hörmander. *The Analysis of Linear Partial Differential Operators I: Distribution Theory And Fourier Analysis*. Springer, 1990.
- [33] A.R.P. Rau1 H.M. Srivastava2 H.S. Cohl1, J.E. Tohline1. Developments in determining the gravitational potential using toroidal functions. *Astron. Nachr.*, 321:363–372, 2000.
- [34] A. Huber. A theorem of phragmn-lindelf type. *Proc. Amer. Math. Soc.*, 4:852–857, 1953.
- [35] A. Huber. On the uniqueness of generalized axially symmetric potentials. *Annals of Mathematics*, Vol. 60, no 2:351–358, 1954.
- [36] A. Huber. Some results on generalized axially symmetric potentials. *Proceedings of the conference on differential equations (dedicated to A. Weinstein)*, pages 147–155, 1956.
- [37] N.N. Lebedev. *Special functions and their applications*. Prentice-Hall, 1965.
- [38] Hong Liu. The cauchy problem for an axially symmetric equation and the schwarz potential conjecture for the torus. *Journal of Mathematical Analysis and Applications*, 250:387–405, 2000.
- [39] J. D. Love. The dielectric ring in a uniform, axial, electrostatic field. *J. Mathematical Phys.*, 13:1297–1304, 1972.
- [40] N. K. Nikolski. *Operators, Functions, and Systems: An Easing Reading, Volume 2*. AMS, 2002.
- [41] T. V. Savina. On splitting up singularities of fundamental solutions to elliptic equations in \mathbb{C}^2 . *Central European Journal of Mathematics*, 5:733740, 2007.
- [42] J. Segura and A. Gil. Evaluation of toroidal harmonics. *Computer Physics Communications*, 124:104–122, 2000.
- [43] G. Ch. Shushkevich. Electrostatic problem for a torus and a disk. *Zh. Tekh. Fiz.*, 67:123–125, 1997.
- [44] B. Ph. van Milligen and A. Lopez Fraguas. Expansion of vacuum magnetic fields in toroidal harmonics. *Computer Physics Communications*, 81:74–90, 1994.
- [45] I.N. Vekua. *New methods for solving elliptic equations*. North-Holland Publishing Co., Amsterdam; Interscience Publishers John Wiley & Sons, Inc., New York, 1967.
- [46] Nina Virchenko and Iryna Fedotova. *Generalized associated Legendre functions and their applications*. World Scientific Publishing Co. Inc., River Edge, NJ, 2001. With a foreword by Semyon Yakubovich.
- [47] R. J. Weinacht. Fundamental solutions for a class of singular equations. *Contributions to Differential equations*, 3:43–55, 1964.
- [48] R. J. Weinacht. A mean value theorem in generalized axially symmetric potential theory. *Atti Accad. Naz. Lincei Rend. Cl. Sci. Fis. Mat. Natur. (8)*, 38:610–613, 1965.
- [49] R. J. Weinacht. Fundamental solutions for a class of equations with several singular coefficients. *J. Austral. Math. Soc.*, 8:574–583, 1968.
- [50] A. Weinstein. Discontinuous integrals and generalized potential theory. *Trans. Amer. Math. Soc.*, 63:342–354, 1948.
- [51] A. Weinstein. On generalized potential theory and on the torsion of shafts. *Studies and essays Presented to R. Courant on his 60th Birthday*, pages 451–460, 1948.
- [52] A. Weinstein. On the torsion of shafts of revolution. *Proc. Seventh Internat. Congress Appl. Mech.*, 1:108–119, 1948.

- [53] A. Weinstein. Transonic flow and generalized axially symmetric potential theory. *Symposium on theoretical compressible flow*, 28:73–82, 1949.
- [54] A. Weinstein. On tricomi's equation and generalized axially symmetric potential theory. *Acad. Roy. Belg. Bull. Cl. Sci. (5)*, 37:348–358, 1951.
- [55] A. Weinstein. Generalized axially symmetric potential theory. *Bull. Amer. Math. Soc*, 59:20–38, 1953.
- [56] A. Weinstein. Elliptic and hyperbolic axially symmetric problems. *Proceedings of the International Congress of Mathematicians*, III:264–269, 1954.
- [57] A. Weinstein. The method of axial symmetry in partial differential equations. *Convegno Internazionale sulle Equazioni Lineari alle Derivate Parziali, Trieste*, pages 86–96, 1954.
- [58] A. Weinstein. The singular solutions and the cauchy problem for generalized tricomi equations. *Comm. Pure Appl. Math.*, 7:105–116, 1954.
- [59] A. Weinstein. The generalized radiation problem and the euler-poisson-darboux equation. *Summa. Brasil. Math*, 3:125–147, 1955.
- [60] A. Weinstein. On a class of partial differential equations of even order. *Ann. Mat. Pura Appl.*, 39:245–254, 1955.
- [61] A. Weinstein. Sur une classe d'équations aux dérivées partielles singulières. *La théorie des équations aux dérivées partielles, colloques Internationaux du CNRS*, LXXI:179–186, 1956.
- [62] A. Weinstein. On a singular differential operator. *Ann. Mat. Pura Appl.*, 49:359365., 1960.
- [63] A. Weinstein. Singular partial differential equations and their applications. *Fluid Dynamics and Applied Mathematics (Proc. Sympos., Univ. of Maryland, 1961)*, pages 29–49, 1962.
- [64] A. Weinstein. Some applications of theory of generalized axially symmetric potential theory to continuum mechanics. *Appl. Theory of Functions in Continuum Mechanics (Proc. Internat. Sympos., Tbilisi, 1963), Fluid and Gas Mechanics, Math. Methods (Russian)*, Vol. II:440–453, 1965.

S. CHAABI, AIX-MARSEILLE UNIVERSITÉ, I2M, UMR 7373, CMI, 39 RUE FRÉDÉRIC JOLIOT-CURIE 13453 MARSEILLE, FRANCE
E-mail address: slah.chaabi@univ-amu.fr

S. RIGAT, AIX-MARSEILLE UNIVERSITÉ, I2M, UMR 7373, CMI, 39 RUE FRÉDÉRIC JOLIOT-CURIE 13453 MARSEILLE, FRANCE
E-mail address: stephane.rigat@univ-amu.fr