

On the Distribution of Values and Zeros of Polynomial Systems over Arbitrary Sets

BRYCE KERR

Department of Computing, Macquarie University
Sydney, NSW 2109, Australia
bryce.kerr@mq.edu.au

IGOR E. SHPARLINSKI*

Department of Computing, Macquarie University
Sydney, NSW 2109, Australia
igor.shparlinski@mq.edu.au

Abstract

Let $G_1, \dots, G_n \in \mathbb{F}_p[X_1, \dots, X_m]$ be n polynomials in m variables over the finite field \mathbb{F}_p of p elements. A result of É. Fouvry and N. M. Katz shows that under some natural condition, for any fixed ε and sufficiently large prime p the vectors of fractional parts

$$\left(\left\{ \frac{G_1(\mathbf{x})}{p} \right\}, \dots, \left\{ \frac{G_n(\mathbf{x})}{p} \right\} \right), \quad \mathbf{x} \in \Gamma,$$

are uniformly distributed in the unit cube $[0, 1]^n$ for any cube $\Gamma \in [0, p-1]^m$ with the side length $h \geq p^{1/2}(\log p)^{1+\varepsilon}$. Here we use this result to show the above vectors remain uniformly distributed, when \mathbf{x} runs through a rather general set. We also obtain new results about the distribution of solutions to system of polynomial congruences.

*This work was supported in part by ARC Grant DP1092835

Mathematics Subject Classification (2010): 11D79, 11K38, 11L07

1 Introduction

Let p be a prime and let \mathbb{F}_p be the finite field of p elements, which we assume to be represented by the set $\{0, 1, \dots, p-1\}$.

Given n polynomials $G_j(X_1, \dots, X_m) \in \mathbb{F}_p[X_1, \dots, X_m]$, $j = 1, \dots, n$, in m variables with integer coefficients, we consider the following points formed by fractional parts:

$$\left(\left\{ \frac{G_1(\mathbf{x})}{p} \right\}, \dots, \left\{ \frac{G_n(\mathbf{x})}{p} \right\} \right), \quad \mathbf{x} = (x_1, \dots, x_m) \in \mathbb{F}_p^m. \quad (1)$$

We say that the polynomials G_1, \dots, G_n are *degree 2 independent over \mathbb{F}_p* if any non-trivial linear combinations $a_1 G_1 + \dots + a_n G_n$ is a polynomial of degree at least 2 over \mathbb{F}_p .

Let $\mathfrak{G}_{m,n,p}$ denote the family of polynomial systems $\{G_1, \dots, G_n\}$ of n polynomials in m variables that are degree 2 independent over \mathbb{F}_p .

Fouvry and Katz [3] have shown that for any $\{G_1, \dots, G_n\} \in \mathfrak{G}_{m,n,p}$, the points (1) are uniformly distributed in the unit cube $[0, 1]^n$, where \mathbf{x} runs through the integral points in any cube $\Gamma \in [0, p-1]^m$ with side length $h \geq p^{1/2}(\log p)^{1+\varepsilon}$. Here we use several of the results from [3] combined with some ideas of Schmidt [7] to obtain a similar uniformity of distribution result when \mathbf{x} runs through a set from a very general family. For example, this holds for \mathbf{x} that belong to the dilate $p\Omega$ of a convex set $\Omega \in [0, 1]^m$ of Lebesgue measure at least $p^{-1/2+\varepsilon}$ for any fixed $\varepsilon > 0$ and sufficiently large p . We note that standard way of moving from boxes to arbitrary convex sets, via the isotropic discrepancy, see [7, Theorem 2], leads to a much weaker result which is nontrivial only for sets $\Omega \in [0, 1]^m$ of Lebesgue measure at least $p^{-1/2m+\varepsilon}$.

As in [8], it is crucial for our approach that the error term in the aforementioned asymptotic formula of [3] depends on the size of the cube $\Gamma \in [0, p-1]^m$ and decreases rapidly together with the size of Γ . We note that a similar idea has also recently been used in [4] in combination with a new upper bound on the number of zeros of multivariate polynomial congruences in small cubes.

Furthermore, given n polynomials $F_j(X_1, \dots, X_m) \in \mathbb{Z}[X_1, \dots, X_m]$, $j = 1, \dots, n$, we consider the distribution of points in the set \mathcal{X}_p , of solutions $\mathbf{x} = (x_1, \dots, x_m) \in \mathbb{F}_p^m$ to the system of congruences

$$F_j(\mathbf{x}) \equiv 0 \pmod{p}, \quad j = 1, \dots, n. \quad (2)$$

Let $\mathfrak{F}_{m,n}$ denote the family of polynomial systems $\{F_1, \dots, F_n\}$ of n polynomials in $m \geq n+1$ variables with integer coefficients, such that the solution set of the system of equations (over \mathbb{C})

$$F_j(\mathbf{x}) = 0, \quad j = 1, \dots, n,$$

has at least one absolutely irreducible component of dimension $m - n$ and no absolutely irreducible component is contained in a hyperplane. For sufficiently large p all absolutely irreducible components remain of the same dimension and are absolutely irreducible modulo p , so by the Lang-Weil theorem [6] we have

$$\#\mathcal{X}_p = \nu p^{m-n} + O(p^{m-n-1/2}), \quad (3)$$

where ν is the number of absolutely irreducible components of \mathcal{X}_p of dimension $m - n$. It is shown in [8], that for a rather general class of sets Ω , including all convex sets, we have

$$T_p(\Omega) = \#\mathcal{X}_p(\mu(\Omega) + O(p^{-1/2(n+1)} \log p)) \quad (4)$$

with

$$T_p(\Omega) = \#(\mathcal{X}_p \cap \Omega).$$

The asymptotic formula (4) is based on a combination of a result of Fouvry [2] and Schmidt [7].

Here we show that for a more restricted class of sets, which includes such natural sets as m -dimensional balls, one can improve (4) and obtain an asymptotic formula which is nontrivial provided that

$$\mu(\Omega) \geq p^{-1/2+\varepsilon}$$

for any fixed $\varepsilon > 0$ and a sufficiently large p , while (4) is nontrivial only under the condition $\mu(\Omega) \geq p^{-1/2(n+1)+\varepsilon}$ (but applies to a wider class of sets).

2 Well and Very Well Shaped Sets

Let $\mathbb{T}_s = (\mathbb{R}/\mathbb{Z})^s$ be the s -dimensional unit torus.

We define the distance between a vector $\mathbf{u} \in \mathbb{T}_m$ and a set $\Omega \subseteq \mathbb{T}_m$ by

$$\text{dist}(\mathbf{u}, \Omega) = \inf_{\mathbf{w} \in \Omega} \|\mathbf{u} - \mathbf{w}\|,$$

where $\|\mathbf{v}\|$ denotes the Euclidean norm of \mathbf{v} . Given $\varepsilon > 0$ and a set $\Omega \subseteq \mathbb{T}_m$, we define the sets

$$\Omega_\varepsilon^+ = \{\mathbf{u} \in \mathbb{T}_m \setminus \Omega : \text{dist}(\mathbf{u}, \Omega) < \varepsilon\}$$

and

$$\Omega_\varepsilon^- = \{\mathbf{u} \in \Omega : \text{dist}(\mathbf{u}, \mathbb{T}_m \setminus \Omega) < \varepsilon\}.$$

We say that a set Ω is *well shaped* if

$$\mu(\Omega_\varepsilon^\pm) \leq C\varepsilon, \tag{5}$$

for some constant C , where μ is the Lebesgue measure on \mathbb{T}_m .

It is known that any convex set is well shaped, see [7, Lemma 1].

Finally, a very general result of Weyl [9, Equation (2)] (taken with $n = m$ and $\nu = n - 1$), that actually expresses $\mu(\Omega_\varepsilon^\pm)$ via a finite sum of powers ε^i , $i = 1, \dots, m$ in the case when the boundary of Ω is manifold of dimension $n - 1$. Examining the constants in this expansion we see that any such set with a bounded surface size is well shaped.

Furthermore, we say that a set $\Omega \subseteq \mathbb{T}_m$ is *very well shaped* if for every $\varepsilon > 0$ the measures of the sets Ω_ε^\pm exist and satisfy

$$\mu(\Omega_\varepsilon^\pm) \leq C(\mu(\Omega)^{1-1/m}\varepsilon + \varepsilon^m) \tag{6}$$

for some $C > 0$, the most natural example of a very well shaped set being a Euclidian ball.

We recall that the notation $A(t) \ll B(t)$ is equivalent to $A(t) = O(B(t))$, which means that there exists some absolute constant, α , such that $|A(t)| \leq \alpha B(t)$ for all values of t within a certain range. Throughout the paper, the implied constants in symbols ‘ O ’ and ‘ \ll ’ may depend on the constant C in (5) and (6) and it may also depend on the polynomial system $\{F_1, \dots, F_n\} \in \mathfrak{F}_{m,n}$ (but does not depend on the polynomial system $\{G_1, \dots, G_n\} \in \mathfrak{G}_{m,n,p}$).

3 Discrepancy

Given a sequence Ξ of N points

$$\Xi = \{(\xi_{k,1}, \dots, \xi_{k,n})\}_{k=1}^N, \quad (7)$$

in \mathbb{T}_n , we define its *discrepancy* as

$$\Delta(\Xi) = \sup_{\Pi \subseteq \mathbb{T}_n} \left| \frac{\#A(\Xi, \Pi)}{N} - \lambda(\Pi) \right|,$$

where $A(\Xi, \Pi)$ is the number of $k \leq N$ such that $(\xi_{k,1}, \dots, \xi_{k,n}) \in \Pi$, λ is the Lebesgue measure on \mathbb{T}_n and the supremum is taken over all boxes

$$\Pi = [\alpha_1, \beta_1) \times \dots \times [\alpha_n, \beta_n) \subseteq \mathbb{T}_n, \quad (8)$$

see [1, 5].

We also define the discrepancy of an empty sequence as 1.

4 Main Results

For a set $\Omega \subseteq \mathbb{T}_m$ let $D(\Omega)$ be the discrepancy of the points

$$\left(\left\{ \frac{G_1(\mathbf{x})}{p} \right\}, \dots, \left\{ \frac{G_n(\mathbf{x})}{p} \right\} \right), \quad \mathbf{x} \in p\Omega.$$

Theorem 1. *For any polynomial system $\{G_1, \dots, G_n\} \in \mathfrak{G}_{m,n,p}$ and any well shaped set $\Omega \in \mathbb{T}_m$, we have*

$$D(\Omega) \ll \mu(\Omega)^{-1} p^{-1/2} (\log p)^{n+2}.$$

We can get a sharper error term for the case of very well shaped sets.

Theorem 2. *For any polynomial system $\{G_1, \dots, G_n\} \in \mathfrak{G}_{m,n,p}$ and any very well shaped set $\Omega \in \mathbb{T}_m$, we have*

$$D(\Omega) \ll \mu(\Omega)^{-1/m} p^{-1/2} (\log p)^{n+2}.$$

We prove the following

Theorem 3. *For any polynomial system $\{F_1, \dots, F_n\} \in \mathfrak{F}_{m,n}$ and any very well shaped set $\Omega \in \mathbb{T}_m$, we have*

$$T_p(\Omega) = \#\mathcal{X}_p(\mu(\Omega) + O(\mu(\Omega)^{1-1/m} p^{-1/2(n+1)} \log p + p^{-1/2} (\log p)^{n+2})).$$

5 Exponential Sum and Congruences

Typically the bounds on the discrepancy of a sequence are derived from bounds of exponential sums with elements of this sequence. The relation is made explicit in the celebrated *Koksma–Szűsz inequality*, see [1, Theorem 1.21], which we present in the following form.

Lemma 4. *Suppose that for the sequence of points (7) for some integer $L \geq 1$ and the real number S we have*

$$\left| \sum_{k=1}^N \exp \left(2\pi i \sum_{j=1}^n a_j \xi_{k,j} \right) \right| \leq S,$$

for all integers $-L \leq a_j \leq L$, $j = 1, \dots, n$, not all equal to zero. Then,

$$D(\Gamma) \ll \frac{1}{L} + \frac{(\log L)^n}{N} S,$$

where the implied constant depends only on n .

To use Lemma 4 we need the following bound of Fouvry and Katz [3, Equation (10.6)]

Lemma 5. *For any polynomial system $\{G_1, \dots, G_n\} \in \mathfrak{G}_{m,n,p}$ and arbitrary integers u and w with $1 \leq w < p$, uniformly over all non-zero modulo p integer vectors (a_1, \dots, a_n) we have*

$$\sum_{x_1, \dots, x_m = u}^{u+w} \exp \left(\frac{2\pi i}{p} \sum_{j=1}^n a_j G_j(x_1, \dots, x_m) \right) \ll p^{1/2} w^{m-1} \log p.$$

Proof. The bound in [3, Equation (10.6)], that gives the desired result for $u = 0$ is uniform in polynomials G_1, \dots, G_m . It now remains to notice that the property of being degree 2 independent is preserved under the change of variables $X_j \rightarrow X_j + u$, $j = 1, \dots, m$. \square

The proof of Theorem 3 is based on the following bound for $T_p(\mathcal{C})$ for a cube \mathcal{C} which is essentially a result of Fouvry [2]

Lemma 6. For any polynomial system $\{F_1, \dots, F_n\} \in \mathfrak{F}_{m,n}$ and any cubic box

$$\mathcal{C} = \left[\gamma_1 + \frac{u_1}{k}, \gamma_1 + \frac{u_1 + 1}{k} \right] \times \dots \times \left[\gamma_m + \frac{u_m}{k}, \gamma_m + \frac{u_m + 1}{k} \right] \subseteq \mathbb{R}^m,$$

where $u_1, \dots, u_m \in \mathbb{Z}$, of side length $1/k$, we have

$$\begin{aligned} T_p(\mathcal{C}) &= \#\mathcal{X}_p \left(\frac{1}{k} \right)^m \\ &\quad + O \left(p^{(m-n)/2} (\log p)^m + \left(\frac{1}{k} \right)^{m-n-1} p^{m-n-1/2} (\log p)^{n+1} \right). \end{aligned}$$

6 Proof of Theorem 1

For a set $\Omega \subseteq \mathbb{T}_m$ and a box $\Pi \subseteq \mathbb{T}_n$ of the form (8) let $N(\Omega; \Pi)$ be the number of integer vectors $\mathbf{x} \in p\Omega$ for which the points (1) belong to Π .

In particular, let $N(\Omega) = N(\Omega; \mathbb{T}_m)$ be the number of integer vectors $\mathbf{x} \in p\Omega$. A simple geometric argument shows that if $\Omega = \Gamma \subseteq \mathbb{T}_m$ is a cube then

$$N(\Gamma) = \mu(\Gamma)p^m + O \left(p^{m-1} \mu(\Gamma)^{(m-1)/m} \right). \quad (9)$$

We start with deriving a lower bound on $N(\Omega; \Pi)$.

We now recall some constructions and arguments from the proof of [7, Theorem 2]. Pick a point $\boldsymbol{\gamma} = (\gamma_1, \dots, \gamma_m) \in \mathbb{T}_m$ such that all its coordinates are irrational. For positive k , let $\mathfrak{C}(k)$ be the set of cubes of the form

$$\left[\gamma_1 + \frac{u_1}{k}, \gamma_1 + \frac{u_1 + 1}{k} \right] \times \dots \times \left[\gamma_m + \frac{u_m}{k}, \gamma_m + \frac{u_m + 1}{k} \right] \subseteq \mathbb{R}^m,$$

where $u_1, \dots, u_m \in \mathbb{Z}$. Note that the above irrationality condition guarantees that the points $p^{-1}\mathbf{x}$ with $\mathbf{x} \in \mathbb{Z}^m$ all belong to the interior of the cubes from $\mathfrak{C}(k)$.

Furthermore, let $\mathcal{C}(k)$ be the set of cubes from $\mathfrak{C}(k)$ that are contained inside of Ω . By [7, Equation (9)], for any well shaped set $\Omega \in \mathbb{T}_m$, we have

$$\#\mathcal{C}(k) = k^m \mu(\Omega) + O(k^{m-1}). \quad (10)$$

Let $\mathcal{B}_1 = \mathcal{C}(2)$ and for $i = 2, 3, \dots$, let \mathcal{B}_i be the set of cubes $\Gamma \in \mathcal{C}(2^i)$ that are not contained in any cube from $\mathcal{C}(2^{i-1})$. Clearly

$$2^{-im} \#\mathcal{B}_i + 2^{-(i-1)m} \#\mathcal{C}(2^{i-1}) \leq \mu(\Omega), \quad i = 2, 3, \dots$$

We now see from (10) that

$$\#\mathcal{B}_i \ll 2^{i(m-1)} \quad (11)$$

and also for any integer $M \geq 1$,

$$\Omega \setminus \Omega_\varepsilon^- \subseteq \bigcup_{i=1}^M \bigcup_{\Gamma \in \mathcal{B}_i} \Gamma \subseteq \Omega$$

with $\varepsilon = m^{1/2}2^{-M}$. Since Ω is well shaped, we obtain

$$\mu \left(\bigcup_{i=1}^M \bigcup_{\Gamma \in \mathcal{B}_i} \Gamma \right) = \mu(\Omega) + O(2^{-M}). \quad (12)$$

Using Lemma 4 (taken with $L = (p-1)/2$) and recalling the bound of Lemma 5 we see that the discrepancy $D(\Gamma)$ of the points (1) with $\mathbf{x} \in p\Gamma$, for a cube Γ satisfies

$$D(\Gamma) \ll \frac{p^{1/2} (p\mu(\Gamma)^{1/m})^{m-1}}{\mu(\Gamma)p^m} (\log p)^{n+1} = p^{-1/2} \mu(\Gamma)^{-1/m} (\log p)^{n+1}.$$

Therefore, using (9), we derive

$$\begin{aligned} N(\Gamma; \Pi) &= \lambda(\Pi)N(\Gamma) + O(N(\Gamma)D(\Gamma)) \\ &= \lambda(\Pi)\mu(\Gamma)p^m + O(p^{m-1/2}\mu(\Gamma)^{(m-1)/m}(\log p)^{n+1}). \end{aligned} \quad (13)$$

Hence

$$N(\Omega; \Pi) \geq \sum_{i=1}^M \sum_{\Gamma \in \mathcal{B}_i} N(\Gamma; \Pi) = \lambda(\Pi)p^m \sum_{i=1}^M \sum_{\Gamma \in \mathcal{B}_i} \mu(\Gamma) + O(R), \quad (14)$$

where

$$R = p^{m-1/2}(\log p)^{n+1} \sum_{i=1}^M \#\mathcal{B}_i 2^{-i(m-1)}.$$

We see from (12) that

$$\sum_{i=1}^M \sum_{\Gamma \in \mathcal{B}_i} \mu(\Gamma) = \mu \left(\bigcup_{i=1}^M \bigcup_{\Gamma \in \mathcal{B}_i} \Gamma \right) = \mu(\Omega) + O(2^{-M}). \quad (15)$$

Furthermore, using (11), we derive

$$R \ll Mp^{m-1/2}(\log p)^{n+1}. \quad (16)$$

We now choose M to satisfy

$$2^M \leq p^{1/2} < 2^{(M+1)}.$$

Now, substituting (15) and (16) in (14) with the above choice of M , we obtain

$$N(\Omega; \Pi) \geq \lambda(\Pi)\mu(\Omega)p^m + O(p^{m-1/2}(\log p)^{n+2}). \quad (17)$$

Since the complementary set $\bar{\Omega} = \mathbb{T}_m \setminus \Omega$ is also well shaped, we also have

$$N(\bar{\Omega}; \Pi) \leq \lambda(\Pi)\mu(\bar{\Omega})p^m + O(p^{m-1/2}(\log p)^{n+2}). \quad (18)$$

Note that by (13) we have

$$N(\mathbb{T}_m; \Pi) = \lambda(\Pi)p^m + O(p^{m-1/2}(\log p)^{n+1}).$$

Now, since

$$N(\bar{\Omega}; \Pi) = N(\mathbb{T}_m; \Pi) - N(\Omega; \Pi) \quad \text{and} \quad \mu(\bar{\Omega}) = 1 - \mu(\Omega),$$

we now see that (18) implies that upper bound

$$N(\Omega; \Pi) \leq \lambda(\Pi)\mu(\Omega)p^m + O(p^{m-1/2}(\log p)^{n+2})$$

together with (17) leads to the desired asymptotic formula

$$N(\Omega; \Pi) = \lambda(\Pi)\mu(\Omega)p^m + O(p^{m-1/2}(\log p)^{n+2}).$$

Since $D(\Omega) \leq 1$, we can assume that

$$\mu(\Omega) \geq c_0 p^{-1/2}(\log p)^{n+2}.$$

for a sufficiently large constant $c_0 > 0$ as otherwise the result is trivial. Thus

$$\frac{N(\Omega; \Pi)}{N(\Omega)} = \lambda(\Pi) + O(\mu(\Omega)^{-1}p^{-1/2}(\log p)^{n+2})$$

which concludes the proof.

7 Proof of Theorem 2

If Ω is very well shaped we may use the same method as the proof of Theorem 1 to replace the bounds (11) and (12) with

$$\#\mathcal{B}_i \ll 1 + \mu(\Omega)^{(m-1)/m} 2^{i(m-1)} \quad (19)$$

and

$$\mu\left(\bigcup_{i=1}^M \bigcup_{\Gamma \in \mathcal{B}_i} \Gamma\right) = \mu(\Omega) + O(\mu(\Omega)^{(m-1)/m} 2^{-M} + 2^{-Mm}). \quad (20)$$

Recalling the lower bound (14)

$$N(\Omega; \Pi) \geq \lambda(\Pi) p^m \sum_{i=1}^M \sum_{\Gamma \in \mathcal{B}_i} \mu(\Gamma) + O(R),$$

where

$$R = p^{m-1/2} (\log p)^{n+1} \sum_{i=1}^M \#\mathcal{B}_i 2^{-i(m-1)}.$$

We use (19) to bound the term R . First we note if $\#\mathcal{B}_i > 0$ we must have $2^{-im} \leq \mu(\Omega)$. Hence

$$\begin{aligned} \sum_{i=1}^M \#\mathcal{B}_i 2^{-i(m-1)} &= \sum_{i \geq -\log \mu(\Omega)/(m \log 2)}^M \#\mathcal{B}_i 2^{-i(m-1)} \\ &\ll \sum_{i \geq -\log \mu(\Omega)/(m \log 2)}^M (\mu(\Omega)^{(m-1)/m} + 2^{-i(m-1)}) \\ &\ll M \mu(\Omega)^{(m-1)/m} + \mu(\Omega)^{(m-1)/m} \\ &\ll M \mu(\Omega)^{(m-1)/m} \end{aligned}$$

so that

$$R \ll M \mu(\Omega)^{(m-1)/m} p^{m-1/2} (\log p)^{n+1}.$$

By (20) we have,

$$\sum_{i=1}^M \sum_{\Gamma \in \mathcal{B}_i} \mu(\Gamma) = \mu\left(\bigcup_{i=1}^M \bigcup_{\Gamma \in \mathcal{B}_i} \Gamma\right) = \mu(\Omega) + O(\mu(\Omega)^{(m-1)/m} 2^{-M} + 2^{-Mm}).$$

Hence

$$N(\Omega; \Pi) \geq \lambda(\Pi)\mu(\Omega)p^m + O(\mu(\Omega)^{(m-1)/m}p^m2^{-M} + p^m2^{-Mm}) \\ + M\mu(\Omega)^{(m-1)/m}p^{m-1/2}(\log p)^{n+1}.$$

Since $D(\Omega) \leq 1$, we can assume that

$$\mu(\Omega) \geq c_0p^{-m/2}(\log p)^{m(n+2)}$$

for a sufficiently large constant $c_0 > 0$. Thus, choosing M so that

$$2^M \leq p \leq 2^{(M+1)},$$

gives

$$N(\Omega; \Pi) \geq \lambda(\Pi)\mu(\Omega)p^m + O(\mu(\Omega)^{(m-1)/m}p^{m-1/2}(\log p)^{n+2}).$$

The upper bounds for $N(\Omega; \Pi)$ and $D(\Omega)$ follow the same method as in the proof of Theorem 1.

8 Proof of Theorem 3

Given Ω very well shaped, we consider the same constructions in the proof of Theorem 1. As in Theorem 2 we have the bound

$$\#\mathcal{B}_i \ll 1 + \mu(\Omega)^{1-1/m}2^{i(m-1)}. \quad (21)$$

The set inclusions

$$\Omega \setminus \Omega_\varepsilon^- \subseteq \bigcup_{i=1}^M \bigcup_{\Gamma \in \mathcal{B}_i} \Gamma \subseteq \Omega \quad (22)$$

give the approximation

$$\mu\left(\bigcup_{i=1}^M \bigcup_{\Gamma \in \mathcal{B}_i} \Gamma\right) = \mu(\Omega) + O\left(\frac{\mu(\Omega)^{1-1/m}}{2^M} + \frac{1}{2^{mM}}\right). \quad (23)$$

Using Lemma 6 and (22),

$$T_p(\Omega) \geq \sum_{i=1}^M \sum_{\Gamma \in \mathcal{B}_i} T_p(\Gamma) = \#\mathcal{X}_p \sum_{n=1}^M \sum_{\Gamma \in \mathcal{B}_i} \mu(\Gamma) + O(R),$$

where

$$R = \sum_{i=1}^M \#B_i \left(p^{(m-n)/2} (\log p)^m + 2^{-i(m-n-1)} p^{m-n-1/2} (\log p)^{n+1} \right).$$

By (23)

$$\#\mathcal{X}_p \sum_{n=1}^M \sum_{\Gamma \in \mathcal{B}_i} \mu(\Gamma) = \#\mathcal{X}_p \left(\mu(\Omega) + O \left(\frac{\mu(\Omega)^{1-1/m}}{2^M} + \frac{1}{2^{mM}} \right) \right)$$

and using (3) we have

$$\#\mathcal{X}_p \sum_{n=1}^M \sum_{\Gamma \in \mathcal{B}_i} \mu(\Gamma) = \#\mathcal{X}_p \mu(\Omega) + O \left(\frac{p^{m-n} \mu(\Omega)^{1-1/m}}{2^M} + \frac{p^{m-n}}{2^{mM}} \right).$$

For the term R , by (21)

$$\begin{aligned} R &\ll \sum_{i=1}^M \left(p^{(m-n)/2} (\log p)^m + (p2^{-i})^{m-n-1} p^{1/2} (\log p)^{n+1} \right) \\ &\quad + \sum_{i=1}^M \left(\mu(\Omega)^{1-1/m} 2^{i(m-1)} p^{(m-n)/2} (\log p)^m + (p2^{-i})^{m-n-1} p^{1/2} (\log p)^{n+1} \right) \\ &\ll Mp^{(m-n)/2} (\log p)^m + p^{m-n-1/2} (\log p)^{n+1} \sum_{i=1}^M 2^{-i(m-n-1)} \\ &\quad + \mu(\Omega)^{1-1/m} p^{(m-n)/2} \sum_{i=1}^M 2^{i(m-1)} (\log p)^m \\ &\quad + \mu(\Omega)^{1-1/m} p^{m-n-1/2} \sum_{i=1}^M 2^{in} (\log p)^{n+1} \\ &\ll M \left(p^{(m-n)/2} (\log p)^m + p^{m-n-1/2} (\log p)^{n+1} \right) \\ &\quad + \mu(\Omega)^{1-1/m} \left(2^{M(m-1)} p^{(m-n)/2} (\log p)^m + 2^{Mn} p^{m-n-1/2} (\log p)^{n+1} \right). \end{aligned}$$

Hence we have

$$T_p(\Omega) \geq \#\mathcal{X}_p \mu(\Omega) + O(R_1 + R_2 + R_3) \tag{24}$$

with

$$\begin{aligned}
R_1 &= \frac{p^{m-n}\mu(\Omega)^{1-1/m}}{2^M} + \frac{p^{m-n}}{2^{mM}}, \\
R_2 &= \mu(\Omega)^{1-1/m} (2^{M(m-1)}p^{(m-n)/2}(\log p)^m + 2^{Mn}p^{m-n-1/2}(\log p)^{n+1}), \\
R_3 &= Mp^{m-n-1/2}(\log p)^{n+1}.
\end{aligned}$$

It is clear that for the bound to be nontrivial we have to choose $M = O(\log p)$, under which condition we have

$$R_3 = p^{m-n-1/2}(\log p)^{n+2}.$$

Now considering all four ways of balancing the terms of R_1 and R_2 , after straight forward calculations we conclude that the optimal choice of M is defined by the condition

$$2^{-M} \leq p^{-1/2(n+1)} \log p < 2^{-M+1}.$$

that balances the first term of R_1 and the second term of R_2 . This gives

$$\begin{aligned}
R &\ll p^{m-n-1/2(n+1)}\mu(\Omega)^{1-1/m} \log p + p^{m-n-m/2(n+1)}(\log p)^m \\
&\quad + p^{(m-n)-1/2(n+1)-n(m-1-n)/2(n+1)}\mu(\Omega)^{1-1/m} \log p \\
&\quad \quad \quad + p^{m-n-1/2}(\log p)^{n+2} \\
&\ll p^{m-n-1/2(n+1)}\mu(\Omega)^{1-1/m} \log p + p^{m-n-1/2}(\log p)^{n+2}.
\end{aligned} \tag{25}$$

Hence by (3),

$$\begin{aligned}
T_p(\Omega) &\geq \#\mathcal{X}_p(\mu(\Omega) + O(\mu(\Omega)^{1-1/m}p^{-1/2(n+1)} \log p + p^{-1/2}(\log p)^{n+2})).
\end{aligned} \tag{26}$$

Although since

$$(\mathbb{T}_m \setminus \Omega)_\varepsilon^- = \Omega_\varepsilon^+ \leq C(\mu(\Omega)^{1-1/m}\varepsilon + \varepsilon^m)$$

we may repeat the above argument to get,

$$\begin{aligned}
T_p(\mathbb{T}_m \setminus \Omega) &\geq \#\mathcal{X}_p\mu(\mathbb{T}_m \setminus \Omega) \\
&\quad + O(\#\mathcal{X}_p(\mu(\Omega)^{1-1/m}p^{-1/2(n+1)} \log p + p^{-1/2}(\log p)^{n+2})).
\end{aligned} \tag{27}$$

Finally, combining (3), (26) and (27), gives the desired result.

References

- [1] M. Drmota and R. Tichy, *Sequences, discrepancies and applications*, Springer-Verlag, Berlin, 1997.
- [2] É. Fouvry, ‘Consequences of a result of N. Katz and G. Laumon concerning trigonometric sums’, *Israel J. Math.*, **120** (2000), 81–96.
- [3] É. Fouvry and N. Katz, ‘A general stratification theorem for exponential sums, and applications’, *J. Reine Angew. Math.*, **540** (2001), 115–166.
- [4] B. Kerr, ‘Solutions to polynomial congruences in well shaped sets’, *Preprint*, (available from <http://arxiv.org/abs/1210.1623>).
- [5] L. Kuipers and H. Niederreiter, *Uniform distribution of sequences*, Wiley-Intersci., New York, 1974.
- [6] S. Lang and A. Weil, ‘Number of points of varieties in finite fields’, *Amer. J. Math.*, **76** (1954), 819–827.
- [7] W. Schmidt, ‘Irregularities of distribution. IX’, *Acta Arith.*, **27** (1975), 385–396.
- [8] I. E. Shparlinski, ‘On the distribution of solutions to polynomial congruences’, *Archiv Math.*, (to appear).
- [9] H. Weyl, ‘On the volume of tubes’, *Amer. J. Math.*, **61** (1939), 461–472.