

# SOME EXTENSIONS OF THE OPEN DOOR LEMMA

MING LI AND TOSHIYUKI SUGAWA

ABSTRACT. Miller and Mocanu proved in their 1997 paper a greatly useful result which is now known as the Open Door Lemma. It provides a sufficient condition for an analytic function on the unit disk to have positive real part. Kuroki and Owa modified the lemma when the initial point is non-real. In the present note, by extending their methods, we give a sufficient condition for an analytic function on the unit disk to take its values in a given sector.

## 1. INTRODUCTION

We denote by  $\mathcal{H}$  the class of holomorphic functions on the unit disk  $\mathbb{D} = \{z : |z| < 1\}$  of the complex plane  $\mathbb{C}$ . For  $a \in \mathbb{C}$  and  $n \in \mathbb{N}$ , let  $\mathcal{H}[a, n]$  denote the subclass of  $\mathcal{H}$  consisting of functions  $h$  of the form  $h(z) = a + c_n z^n + c_{n+1} z^{n+1} + \dots$ . Here,  $\mathbb{N} = \{1, 2, 3, \dots\}$ . Let also  $\mathcal{A}_n$  be the set of functions  $f$  of the form  $f(z) = zh(z)$  for  $h \in \mathcal{H}[1, n]$ .

A function  $f \in \mathcal{A}_1$  is called *starlike* (resp. *convex*) if  $f$  is univalent on  $\mathbb{D}$  and if the image  $f(\mathbb{D})$  is starlike with respect to the origin (resp. convex). It is well known (cf. [1]) that  $f \in \mathcal{A}_1$  is starlike precisely if  $q_f(z) = zf'(z)/f(z)$  has positive real part on  $|z| < 1$ , and that  $f \in \mathcal{A}_1$  is convex precisely if  $\varphi_f(z) = 1 + zf''(z)/f'(z)$  has positive real part on  $|z| < 1$ . Note that the following relation holds for those quantities:

$$\varphi_f(z) = q_f(z) + \frac{zq'_f(z)}{q_f(z)}.$$

It is geometrically obvious that a convex function is starlike. This, in turn, means the implication

$$\operatorname{Re} \left[ q(z) + \frac{zq'(z)}{q(z)} \right] > 0 \text{ on } |z| < 1 \quad \Rightarrow \quad \operatorname{Re} q(z) > 0 \text{ on } |z| < 1$$

for a function  $q \in \mathcal{H}[1, 1]$ . Interestingly, it looks highly nontrivial. Miller and Mocanu developed a theory (now called *differential subordination*) which enables us to deduce such a result systematically. See a monograph [4] written by them for details.

The set of functions  $q \in \mathcal{H}[1, 1]$  with  $\operatorname{Re} q > 0$  is called the Carathéodory class and will be denoted by  $\mathcal{P}$ . It is well recognized that the function

---

1991 *Mathematics Subject Classification*. Primary 30C45; Secondary 30C80.

*Key words and phrases*. open door function, subordination.

The authors were supported in part by JSPS Grant-in-Aid for Scientific Research (B) 22340025.

$q_0(z) = (1+z)/(1-z)$  (or its rotation) maps the unit disk univalently onto the right half-plane and is extremal in many problems. One can observe that the function

$$\varphi_0(z) = q_0(z) + \frac{zq_0'(z)}{q_0(z)} = \frac{1+z}{1-z} + \frac{2z}{1-z^2} = \frac{1+4z+z^2}{1-z^2}$$

maps the unit disk onto the slit domain  $V(-\sqrt{3}, \sqrt{3})$ , where

$$V(A, B) = \mathbb{C} \setminus \{iy : y \leq A \text{ or } y \geq B\}$$

for  $A, B \in \mathbb{R}$  with  $A < B$ . Note that  $V(A, B)$  contains the right half-plane and has the “window”  $(Ai, Bi)$  in the imaginary axis to the left half-plane. The Open Door Lemma of Miller and Mocanu asserts for a function  $q \in \mathcal{H}[1, 1]$  that, if  $q(z) + zq'(z)/q(z) \in V(-\sqrt{3}, \sqrt{3})$  for  $z \in \mathbb{D}$ , then  $q \in \mathcal{P}$ . Indeed, Miller and Mocanu [3] (see also [4]) proved it in a more general form. For a complex number  $c$  with  $\operatorname{Re} c > 0$  and  $n \in \mathbb{N}$ , we consider the positive number

$$C_n(c) = \frac{n}{\operatorname{Re} c} \left[ |c| \sqrt{\frac{2\operatorname{Re} c}{n} + 1} + \operatorname{Im} c \right].$$

In particular,  $C_n(c) = \sqrt{n(n+2c)}$  when  $c$  is real. The following is a version of the Open Door Lemma modified by Kuroki and Owa [2].

**Theorem A** (Open Door Lemma). *Let  $c$  be a complex number with positive real part and  $n$  be an integer with  $n \geq 1$ . Suppose that a function  $q \in \mathcal{H}[c, n]$  satisfies the condition*

$$q(z) + \frac{zq'(z)}{q(z)} \in V(-C_n(c), C_n(\bar{c})), \quad z \in \mathbb{D}.$$

*Then  $\operatorname{Re} q > 0$  on  $\mathbb{D}$ .*

**Remark 1.1.** In the original statement of the Open Door Lemma in [3], the slit domain was erroneously described as  $V(-C_n(c), C_n(c))$ . Since  $C_n(\bar{c}) < C_n(c)$  when  $\operatorname{Im} c > 0$ , we see that  $V(-C_n(\bar{c}), C_n(\bar{c})) \subset V(-C_n(c), C_n(\bar{c})) \subset V(-C_n(c), C_n(c))$  for  $\operatorname{Im} c \geq 0$  and the inclusions are strict if  $\operatorname{Im} c > 0$ . As the proof will suggest us, seemingly the domain  $V(-C_n(c), C_n(\bar{c}))$  is maximal for the assertion, which means that the original statement in [3] and the form of the associated open door function are incorrect for a non-real  $c$ . This, however, does not decrease so much the value of the original article [3] by Miller and Mocanu because the Open Door Lemma is mostly applied when  $c$  is real. We also note that the Open Door Lemma deals with the function  $p = 1/q \in \mathcal{H}[1/c, n]$  instead of  $q$ . The present form is adopted for convenience of our aim.

The Open Door Lemma gives a sufficient condition for  $q \in \mathcal{H}[c, n]$  to have positive real part. We extend it so that  $|\arg q| < \pi\alpha/2$  for a

given  $0 < \alpha \leq 1$ . First we note that the Möbius transformation

$$g_c(z) = \frac{c + \bar{c}z}{1 - z}$$

maps  $\mathbb{D}$  onto the right half-plane in such a way that  $g_c(0) = c$ , where  $c$  is a complex number with  $\operatorname{Re} c > 0$ . In particular, one can take an analytic branch of  $\log g_c$  so that  $|\operatorname{Im} \log g_c| < \pi/2$ . Therefore, the function  $q_0 = g_c^\alpha = \exp(\alpha \log g_c)$  maps  $\mathbb{D}$  univalently onto the sector  $|\arg w| < \pi\alpha/2$  in such a way that  $q_0(0) = c^\alpha$ . The present note is based mainly on the following result, which will be deduced from a more general result of Miller and Mocanu (see Section 2).

**Theorem 1.2.** *Let  $c$  be a complex number with  $\operatorname{Re} c > 0$  and  $\alpha$  be a real number with  $0 < \alpha \leq 1$ . Then the function*

$$R_{\alpha,c,n}(z) = g_c(z)^\alpha + \frac{n\alpha z g'_c(z)}{g_c(z)} = \left( \frac{c + \bar{c}z}{1 - z} \right)^\alpha + \frac{2n\alpha(\operatorname{Re} c)z}{(1 - z)(c + \bar{c}z)}$$

is univalent on  $|z| < 1$ . If a function  $q \in \mathcal{H}[c^\alpha, n]$  satisfies the condition

$$q(z) + \frac{zq'(z)}{q(z)} \in R_{\alpha,c,n}(\mathbb{D}), \quad z \in \mathbb{D},$$

then  $|\arg q| < \pi\alpha/2$  on  $\mathbb{D}$ .

We remark that the special case when  $\alpha = 1$  reduces to Theorem 1 (see the paragraph right after Lemma 3.3 below). Also, the case when  $c = 1$  is already proved by Mocanu [5] even under the weaker assumption that  $0 < \alpha \leq 2$  (see Remark 3.6). Since the shape of  $R_{\alpha,c,n}(\mathbb{D})$  is not very clear, we will deduce more concrete results as corollaries of Theorem 1.2 in Section 3. This is our principal aim in the present note.

## 2. PRELIMINARIES

We first recall the notion of subordination. A function  $f \in \mathcal{H}$  is said to be *subordinate* to  $F \in \mathcal{H}$  if there exists a function  $\omega \in \mathcal{H}[0, 1]$  such that  $|\omega| < 1$  on  $\mathbb{D}$  and that  $f = F \circ \omega$ . We write  $f \prec F$  or  $f(z) \prec F(z)$  for subordination. When  $F$  is univalent,  $f \prec F$  precisely if  $f(0) = F(0)$  and if  $f(\mathbb{D}) \subset F(\mathbb{D})$ .

Miller and Mocanu [3, Theorem 5] (see also [4, Theorem 3.2h]) proved the following general result, from which we will deduce Theorem 1.2 in the next section.

**Lemma 2.1** (Miller and Mocanu). *Let  $\mu, \nu \in \mathbb{C}$  with  $\mu \neq 0$  and  $n$  be a positive integer. Let  $q_0 \in \mathcal{H}[c, 1]$  be univalent and assume that  $\mu q_0(z) + \nu \neq 0$  for  $z \in \mathbb{D}$  and  $\operatorname{Re}(\mu c + \nu) > 0$ . Set  $Q(z) = zq'_0(z)/(\mu q_0(z) + \nu)$ , and*

$$(2.1) \quad h(z) = q_0(z) + nQ(z) = q_0(z) + \frac{nzq'_0(z)}{\mu q_0(z) + \nu}.$$

Suppose further that

- (a)  $\operatorname{Re} [zh'(z)/Q(z)] = \operatorname{Re} [h'(z)(\mu q_0(z) + \nu)/q_0'(z)] > 0$ , and
- (b) either  $h$  is convex or  $Q$  is starlike.

If  $p \in \mathcal{H}[c, n]$  satisfies the subordination relation

$$(2.2) \quad q(z) + \frac{zq'(z)}{\mu q(z) + \nu} \prec h(z),$$

then  $q \prec q_0$ , and  $q_0$  is the best dominant. An extremal function is given by  $q(z) = q_0(z^n)$ .

In the investigation of the generalized open door function  $R_{\alpha, c, n}$ , we will need to study the positive solution to the equation

$$(2.3) \quad x^2 + Ax^{1+\alpha} - 1 = 0,$$

where  $A > 0$  and  $0 < \alpha \leq 1$  are constants. Let  $F(x) = x^2 + Ax^{1+\alpha} - 1$ . Then  $F(x)$  is increasing in  $x > 0$  and  $F(0) = -1 < 0$ ,  $F(+\infty) = +\infty$ . Therefore, there is a unique positive solution  $x = \xi(A, \alpha)$  to the equation. We have the following estimates for the solution.

**Lemma 2.2.** *Let  $0 < \alpha \leq 1$  and  $A > 0$ . The positive solution  $x = \xi(A, \alpha)$  to equation (2.3) satisfies the inequalities*

$$(1 + A)^{-1/(1+\alpha)} \leq \xi(A, \alpha) \leq (1 + A)^{-1/2} (< 1).$$

Here, both inequalities are strict when  $0 < \alpha < 1$ .

*Proof.* Set  $\xi = \xi(A, \alpha)$ . Since the above  $F(x)$  is increasing in  $x > 0$ , the inequalities  $F(x_1) \leq 0 = F(\xi) \leq F(x_2)$  imply  $x_1 \leq \xi \leq x_2$  for positive numbers  $x_1, x_2$  and the inequalities are strict when  $x_1 < \xi < x_2$ . Keeping this in mind, we now show the assertion. First we put  $x_2 = (1 + A)^{-1/2}$  and observe

$$F(x_2) = \frac{1}{1 + A} + \frac{A}{(1 + A)^{(1+\alpha)/2}} - 1 \geq \frac{1}{1 + A} + \frac{A}{1 + A} - 1 = 0,$$

which implies the right-hand inequality in the assertion.

Next put  $x_1 = (1 + A)^{-1/(1+\alpha)}$ . Then

$$F(x_1) = \frac{1}{(1 + A)^{2/(1+\alpha)}} + \frac{A}{1 + A} - 1 \leq \frac{1}{1 + A} + \frac{A}{1 + A} - 1 = 0,$$

which implies the left-hand inequality. We note also that  $F(x_1) < 0 < F(x_2)$  when  $\alpha < 1$ . The proof is now complete.  $\square$

### 3. PROOF AND COROLLARIES

Theorem 1.2 can be rephrased in the following.

**Theorem 3.1.** *Let  $c$  be a complex number with  $\operatorname{Re} c > 0$  and  $\alpha$  be a real number with  $0 < \alpha \leq 1$ . Then the function*

$$R_{\alpha, c, n}(z) = g_c(z)^\alpha + \frac{n\alpha z g_c'(z)}{g_c(z)}$$

is univalent on  $|z| < 1$ . If a function  $q \in \mathcal{H}[c^\alpha, n]$  satisfies the subordination condition

$$q(z) + \frac{zq'(z)}{q(z)} \prec R_{\alpha, c, n}(z)$$

on  $\mathbb{D}$ , then  $q(z) \prec g_c(z)^\alpha$  on  $\mathbb{D}$ . The function  $g_c^\alpha$  is the best dominant.

*Proof.* We first show that the function  $Q(z) = \alpha z g_c'(z)/g_c(z)$  is starlike. Indeed, we compute

$$\frac{zQ'(z)}{Q(z)} = 1 - \frac{\bar{c}z}{c + \bar{c}z} + \frac{z}{1-z} = \frac{1}{2} \left[ \frac{c - \bar{c}z}{c + \bar{c}z} + \frac{1+z}{1-z} \right].$$

Thus we can see that  $\operatorname{Re} [zQ'(z)/Q(z)] > 0$  on  $|z| < 1$ . Next we check condition (a) in Lemma 2.1 for the functions  $q_0 = g_c^\alpha, h = R_{\alpha, c, n}$  with the choice  $\mu = 1, \nu = 0$ . We have the expression

$$\frac{zh'(z)}{Q(z)} = q_c(z)^\alpha + n \frac{zQ'(z)}{Q(z)}.$$

Since both terms in the right-hand side have positive real part, we obtain (a). We now apply Lemma 2.1 to obtain the required assertion up to univalence of  $h = R_{\alpha, c, n}$ . In order to show the univalence, we have only to note that the condition (a) implies that  $h$  is close-to-convex, since  $Q$  is starlike. As is well known, a close-to-convex function is univalent (see [1]), the proof has been finished.  $\square$

We now investigate the shape of the image domain  $R_{\alpha, c, n}(\mathbb{D})$  of the generalized open door function  $R_{\alpha, c, n}$  given in Theorem 1.2. Let  $z = e^{i\theta}$  and  $c = re^{it}$  for  $\theta \in \mathbb{R}, r > 0$  and  $-\pi/2 < t < \pi/2$ . Then we have

$$\begin{aligned} R_{\alpha, c, n}(e^{i\theta}) &= \left( \frac{re^{it} + re^{-it}e^{i\theta}}{1 - e^{i\theta}} \right)^\alpha + \frac{2n\alpha e^{i\theta} \cos t}{(1 - e^{i\theta})(e^{it} + e^{-it}e^{i\theta})} \\ &= \left( \frac{r \cos(t - \theta/2)}{\sin(\theta/2)} i \right)^\alpha + \frac{i}{2} \cdot \frac{n\alpha \cos t}{\sin(\theta/2) \cos(t - \theta/2)} \\ &= r^\alpha e^{\pi\alpha i/2} (\cos t \cot(\theta/2) + \sin t)^\alpha + \frac{i}{2} \cdot \frac{n\alpha(1 + \cot^2(\theta/2)) \cos t}{\cos t \cot(\theta/2) + \sin t}. \end{aligned}$$

Let  $x = \cot(\theta/2) \cos t + \sin t$ . When  $x > 0$ , we write  $R_{\alpha, c, n}(e^{i\theta}) = u_+(x) + iv_+(x)$  and get the expressions

$$\begin{cases} u_+(x) = a(rx)^\alpha, \\ v_+(x) = b(rx)^\alpha + \frac{n\alpha}{2 \cos t} \left( x - 2 \sin t + \frac{1}{x} \right), \end{cases}$$

where

$$a = \cos \frac{\beta\pi}{2} \quad \text{and} \quad b = \sin \frac{\beta\pi}{2}.$$

Taking the derivative, we get

$$v'_+(x) = \frac{n\alpha}{2x^2 \cos t} \left[ x^2 + \frac{2br^\alpha \cos t}{n} x^{\alpha+1} - 1 \right].$$

Hence, the minimum of  $v_+(x)$  is attained at  $x = \xi(A, \alpha)$ , where  $A = 2br^\alpha n^{-1} \cos t$ . By using the relation (2.3), we obtain

$$\begin{aligned} \min_{0 < x} v_+(x) &= v_+(\xi) = \frac{n}{2 \cos t} \left( A\xi^\alpha + \alpha\xi + \frac{\alpha}{\xi} \right) - n\alpha \tan t \\ &= \frac{n}{2 \cos t} \left( (\alpha - 1)\xi + \frac{\alpha + 1}{\xi} \right) - n\alpha \tan t = U(\xi), \end{aligned}$$

where

$$U(x) = \frac{n}{2 \cos t} \left( (\alpha - 1)x + \frac{\alpha + 1}{x} \right) - n\alpha \tan t.$$

Since the function  $U(x)$  is decreasing in  $0 < x < 1$ , Lemma 2.2 yields the inequality

$$\begin{aligned} v_+(\xi) &= U(\xi) \geq U((1 + A)^{-1/2}) \\ &= \frac{n}{2 \cos t} \left( \frac{\alpha - 1}{\sqrt{1 + A}} + (\alpha + 1)\sqrt{1 + A} \right) - n\alpha \tan t. \end{aligned}$$

We remark here that

$$U((1 + A)^{-1/2}) > U(1) = \frac{n\alpha(1 - \sin t)}{\cos t} > 0;$$

namely,  $v_+(x) > 0$  for  $x > 0$ . When  $x < 0$ , letting  $y = -x = -\cot(\theta/2) \cos t - \sin t$ , we write  $h(e^{i\theta}) = u_-(y) + iv_-(y)$ . Then, with the same  $a$  and  $b$  as above,

$$\begin{cases} u_-(y) = a(ry)^\alpha, \\ v_-(y) = -b(ry)^\alpha - \frac{n\alpha}{2 \cos t} \left( y + 2 \sin t + \frac{1}{y} \right), \end{cases}$$

We observe here that  $u_+ = u_- > 0$  and, in particular, we obtain the following.

**Lemma 3.2.** *The left half-plane  $\Omega_1 = \{w : \operatorname{Re} w < 0\}$  is contained in  $R_{\alpha, c, n}(\mathbb{D})$ .*

We now look at  $v_-(y)$ . Since

$$v'_-(y) = -\frac{n\alpha}{2y^2 \cos t} \left[ y^2 + \frac{2br^\alpha \cos t}{n} y^{\alpha+1} - 1 \right],$$

in the same way as above, we obtain

$$\begin{aligned} \max_{0 < y} v_-(y) &= v_-(\xi) = -\frac{n}{2 \cos t} \left( (\alpha - 1)\xi + \frac{\alpha + 1}{\xi} \right) - n\alpha \tan t \\ &\leq -\frac{n}{2 \cos t} \left( \frac{\alpha - 1}{\sqrt{1 + A}} + (\alpha + 1)\sqrt{1 + A} \right) - n\alpha \tan t, \end{aligned}$$

where  $\xi = \xi(A, \alpha)$  and  $A = 2br^\alpha n^{-1} \cos t$ . Note also that  $v_-(y) < 0$  for  $y > 0$ .

Since the horizontal parallel strip  $v_-(\xi) < \operatorname{Im} w < v_+(\xi)$  is contained in the image domain  $R_{\alpha,c,n}(\mathbb{D})$  of the generalized open door function, we obtain the following.

**Lemma 3.3.** *The parallel strip  $\Omega_2$  described by*

$$|\operatorname{Im} w + n\alpha \tan t| < \frac{n}{2 \cos t} \left( \frac{\alpha - 1}{\sqrt{1 + A}} + (\alpha + 1)\sqrt{1 + A} \right)$$

is contained in  $R_{\alpha,c,n}(\mathbb{D})$ . Here,  $t = \arg c \in (-\frac{\pi}{2}, \frac{\pi}{2})$  and  $A = \frac{2}{n}|c|^\alpha \sin \frac{\pi\alpha}{2} \cos t$ .

When  $\alpha = 1$ , we have  $u_\pm = 0$ , that is, the boundary is contained in the imaginary axis. Since  $\xi(A, 1) = (1 + A)^{-1/2}$  by Lemma 2.2, the above computations tell us  $\min v_+ = (n/\cos t)(\sqrt{1 + A} - \sin t) = C_n(\bar{c})$ . Similarly, we have  $\max v_- = -(n/\cos t)(\sqrt{1 + A} + \sin t) = -C_n(c)$ . Therefore, we have  $R_{1,c,n}(\mathbb{D}) = V(-C_n(c), C_n(\bar{c}))$ . Note that the open door function then takes the following form

$$\begin{aligned} R_{1,c,n}(z) &= \frac{c + \bar{c}z}{1 - z} + \frac{2n(\operatorname{Re} c)z}{(1 - z)(c + \bar{c}z)} \\ &= \frac{2\operatorname{Re} c + n}{1 + cz/\bar{c}} - \frac{n}{1 - z} - \bar{c}, \end{aligned}$$

which is the same as given by Kuroki and Owa [2, (2.2)]. In this way, we see that Theorem 1.2 contains Theorem 1 as a special case.

**Remark 3.4.** In [2], they proposed another open door function of the form

$$R(z) = \frac{2n|c|}{\operatorname{Re} c} \sqrt{\frac{2\operatorname{Re} c}{n} + 1} \frac{(\zeta - z)(1 - \bar{\zeta}z)}{(1 - \bar{\zeta}z)^2 - (\zeta - z)^2} - \frac{\operatorname{Im} c}{\operatorname{Re} c} i,$$

where

$$\zeta = 1 - \frac{2}{\omega}, \quad \omega = \frac{c}{|c|} \sqrt{\frac{2\operatorname{Re} c}{n} + 1} + 1.$$

It can be checked that  $R(z) = R_{1,c,n}(-\omega z/\bar{\omega})$ . Hence,  $R$  is just a rotation of  $R_{1,c,n}$ .

We next study the argument of the boundary curve of  $R_{\alpha,c,n}(\mathbb{D})$ . We will assume that  $0 < \alpha < 1$  since we have nothing to do when  $\alpha = 1$ .

As we noted above, the boundary is contained in the right half-plane  $\operatorname{Re} w > 0$ . When  $x > 0$ , we have

$$\frac{v_+(x)}{u_+(x)} = \frac{b}{a} + \frac{n\alpha}{2a r^\alpha x^\alpha \cos t} \left[ x + \frac{1}{x} - 2 \sin t \right].$$

We observe now that  $v_+(x)/u_+(x) \rightarrow +\infty$  as  $x \rightarrow 0+$  or  $x \rightarrow +\infty$ . We also have

$$\left( \frac{v_+}{u_+} \right)' (x) = \frac{n\alpha}{2a r^\alpha x^{\alpha+2} \cos t} \left[ (1 - \alpha)x^2 + 2\alpha x \sin t - (1 + \alpha) \right].$$

Therefore,  $v_+(x)/u_+(x)$  takes its minimum at  $x = \xi$ , where

$$\xi = \frac{-\alpha \sin t + \sqrt{1 - \alpha^2 \cos^2 t}}{1 - \alpha}$$

is the positive root of the equation  $(1 - \alpha)x^2 + 2\alpha x \sin t - (1 + \alpha) = 0$ . It is easy to see that  $1 < \xi$  and that

$$\begin{aligned} T_+ &:= \min_{0 < x} \frac{v_+(x)}{u_+(x)} = \frac{v_+(\xi)}{u_+(\xi)} = \frac{b}{a} + \frac{n\alpha}{2ar^\alpha \xi^\alpha \cos t} \left[ \xi + \frac{1}{\xi} - 2 \sin t \right] \\ &= \tan \frac{\pi\alpha}{2} + \frac{n(\xi - \xi^{-1})}{2ar^\alpha \xi^\alpha \cos t}. \end{aligned}$$

When  $x = -y < 0$ , we have

$$\frac{v_-(y)}{u_-(y)} = -\frac{b}{a} - \frac{n\alpha}{2ar^\alpha y^\alpha \cos t} \left[ y + \frac{1}{y} + 2 \sin t \right]$$

and

$$\left( \frac{v_-}{u_-} \right)' (y) = \frac{-n\alpha}{2ar^\alpha y^{\alpha+2} \cos t} [(1 - \alpha)y^2 - 2\alpha y \sin t - (1 + \alpha)].$$

Hence,  $v_-(y)/u_-(y)$  takes its maximum at  $y = \eta$ , where

$$\eta = \frac{\alpha \sin t + \sqrt{1 - \alpha^2 \cos^2 t}}{1 - \alpha}.$$

Note that

$$T_- := \max_{0 < y} \frac{v_-(y)}{u_-(y)} = \frac{v_-(\eta)}{u_-(\eta)} = -\tan \frac{\pi\alpha}{2} - \frac{n(\eta - \eta^{-1})}{2ar^\alpha \eta^\alpha \cos t}.$$

Therefore, the sector  $\{w : T_- < \arg w < T_+\}$  is contained in the image  $h(\mathbb{D})$ . It is easy to check that  $T_- < -\tan(\pi\alpha/2) < \tan(\pi\alpha/2) < T_+$ . In particular  $T_- < \arg c^\alpha = \alpha t < T_+$ . We summarize the above observations, together with Theorem 1.2, in the following form.

**Corollary 3.5.** *Let  $0 < \alpha < 1$  and  $c = re^{it}$  with  $r > 0$ ,  $-\pi/2 < t < \pi/2$ , and  $n$  be a positive integer. If a function  $q \in \mathcal{H}[c^\alpha, n]$  satisfies the condition*

$$-\Theta_- < \arg \left( q(z) + \frac{zq'(z)}{q(z)} \right) < \Theta_+$$

on  $|z| < 1$ , then  $|\arg q| < \pi\alpha/2$  on  $\mathbb{D}$ . Here,

$$\Theta_\pm = \arctan \left[ \tan \frac{\pi\alpha}{2} + \frac{n(\xi_\pm - \xi_\pm^{-1})}{2r^\alpha \xi_\pm^\alpha \cos(\pi\alpha/2) \cos t} \right],$$

and

$$\xi_\pm = \frac{\mp\alpha \sin t + \sqrt{1 - \alpha^2 \cos^2 t}}{1 - \alpha}.$$



It is a simple task to check that  $x^{1-\alpha} - x^{-1-\alpha}$  is increasing in  $0 < x$ . When  $\text{Im } c > 0$ , we see that  $\xi_- > \xi_+$  and thus  $\Theta_- > \Theta_+$ . It might be useful to note the estimates  $\xi_- < \sqrt{(1+\alpha)/(1-\alpha)} < \xi_+$  and  $\xi_- < 1/\sin t$  for  $\text{Im } c > 0$ .

**Remark 3.6.** When  $c = 1$  and  $n = 1$ , we have  $\xi := \xi_{\pm} = \sqrt{(1+\alpha)/(1-\alpha)}$ ,  $\xi - \xi^{-1} = 2\alpha/\sqrt{1-\alpha^2}$ , and thus

$$\begin{aligned} \Theta_{\pm} &= \arctan \left[ \tan \frac{\pi\alpha}{2} + \frac{\xi - \xi^{-1}}{2\xi^{\alpha} \cos \frac{\pi\alpha}{2}} \right] \\ &= \arctan \left[ \tan \frac{\pi\alpha}{2} + \frac{\alpha}{\cos \frac{\pi\alpha}{2} (1-\alpha)^{\frac{1-\alpha}{2}} (1+\alpha)^{\frac{1+\alpha}{2}}} \right] \\ &= \frac{\pi\alpha}{2} + \arctan \left[ \frac{\alpha \cos \frac{\pi\alpha}{2}}{(1-\alpha)^{\frac{1-\alpha}{2}} (1+\alpha)^{\frac{1+\alpha}{2}} + \alpha \sin \frac{\pi\alpha}{2}} \right]. \end{aligned}$$

Therefore, the corollary gives a theorem proved by Mocanu [6].

Since the values  $\Theta_+$  and  $\Theta_-$  are not given in an explicitly way, it might be convenient to have a simpler sufficient condition for  $|\arg q| < \pi\alpha/2$ .

**Corollary 3.7.** *Let  $0 < \alpha \leq 1$  and  $c$  with  $\text{Re } c > 0$  and  $n$  be a positive integer. If a function  $q \in \mathcal{H}[c^{\alpha}, n]$  satisfies the condition*

$$q(z) + \frac{zq'(z)}{q(z)} \in \Omega,$$

*then  $|\arg q| < \pi\alpha/2$  on  $\mathbb{D}$ . Here,  $\Omega = \Omega_1 \cup \Omega_2 \cup \Omega_3$ , and  $\Omega_1$  and  $\Omega_2$  are given in Lemmas 3.2 and 3.3, respectively, and  $\Omega_3 = \{w \in \mathbb{C} : |\arg w| < \pi\alpha/2\}$ .*

*Proof.* Lemmas 3.2 and 3.3 yield that  $\Omega_1 \cup \Omega_2 \subset R_{\alpha, c, n}(\mathbb{D})$ . Since  $\Theta_{\pm} > \pi\alpha/2$ , we also have  $\Omega_3 \subset R_{\alpha, c, n}(\mathbb{D})$ . Thus  $\Omega \subset R_{\alpha, c, n}(\mathbb{D})$ . Now the result follows from Theorem 1.2.  $\square$

See Figure 1 for the shape of the domain  $\Omega$  together with  $R_{\alpha, c, n}(\mathbb{D})$ . We remark that  $\Omega = R_{\alpha, c, n}(\mathbb{D})$  when  $\alpha = 1$ .

## REFERENCES

- [1] Duren, P. L., *Univalent Functions*, Springer-Verlag, 1983.
- [2] Kuroki, K. and Owa, S., *Notes on the open door lemma*, Rend. Semin. Mat. Univ. Padova, to appear.
- [3] Miller, S. S. and Mocanu, P. T., *Briot-Bouquet differential equations and differential subordinations*, Complex Variables Theory Appl. **33** (1997), 217–237.
- [4] ———, *Differential subordinations. Theory and applications*, Marcel Dekker, Inc., New York, 2000.
- [5] Mocanu, P. T., *On strongly-starlike and strongly-convex functions*, Studia Univ. Babeş-Bolyai Math. **31** (1986), 16–21.




FIGURE 1. The image  $R_{\alpha, c, n}(\mathbb{D})$  and  $\Omega$  for  $\alpha = 1/2, c = 4 + 3i, n = 2$ .

- [6] ———, *Alpha-convex integral operator and strongly starlike functions*, Studia Univ. Babeş-Bolyai, Math. **34** (1989), 18–24.

GRADUATE SCHOOL OF INFORMATION SCIENCES,, TOHOKU UNIVERSITY,, AOBA-KU, SENDAI 980-8579, JAPAN  
*E-mail address:* li@ims.is.tohoku.ac.jp

GRADUATE SCHOOL OF INFORMATION SCIENCES,, TOHOKU UNIVERSITY,, AOBA-KU, SENDAI 980-8579, JAPAN  
*E-mail address:* sugawa@math.is.tohoku.ac.jp