

Correction: The Zeros of the Partial Sums of the Exponential Function

D. J. NEWMAN

Department of Mathematics, Yeshiva University, New York, New York 10033

AND

T. J. RIVLIN

IBM Research Center, Yorktown Heights, New York 10598

Communicated by Oved Shisha

Received June 17, 1974

It has been pointed out to us by E. B. Saff that our proof [1], that there is a parabolic domain which is free of zeros of

$$S_n(z) = \sum_{k=0}^n z^k/k! \tag{1}$$

for n sufficiently large, is incorrect. We present here a proof that there is a parabolic domain (smaller than the one claimed in [1]) free of zeros of $S_n(z)$ for all n .

Write $z = x + iy$ and suppose $y^2 \leq cx$, where c is any positive number satisfying $ce^c < (\pi/2)$. For example $c = 0.7$ may be substituted in what follows.

Case (i). $0 \leq x \leq n$.

$$n! S_n(z) e^{-z} = \int_z^\infty t^n e^{-t} dt = \int_x^\infty t^n e^{-t} dt - \int_x^{x+iy} t^n e^{-t} dt, \tag{2}$$

so that

$$\begin{aligned} n! |S_n(z)| &\geq n! S_n(x) - \int_0^{|y|} |x \pm is|^n ds \\ &\geq n! S_n(x) - |y| (x^2 + y^2)^{n/2} \\ &\geq n! S_n(x) - |y| (x^2 + cx)^{n/2}. \end{aligned} \tag{3}$$

We claim next that $0 \leq x \leq n$ implies that

$$S_n(x) \geq \frac{1}{2}e^x. \quad (4)$$

To see this, note that in view of (2) it suffices to show that

$$\int_x^\infty s^n e^{-s} ds \geq \int_0^x s^n e^{-s} ds, \quad (5)$$

moreover, (5) holds for $0 \leq x \leq n$ if it holds for $x = n$. This is, in turn, a consequence of the inequality

$$(n + nu)^n e^{-(n+nu)} \geq (n - nu)^n e^{-(n-nu)}, \quad 0 < u < 1,$$

or

$$(1 + u) e^{-(1+u)} \geq (1 - u) e^{-(1-u)}, \quad 0 < u < 1,$$

or

$$(1 + u)/(1 - u) \geq e^{2u}, \quad 0 < u < 1,$$

which is well known.

Using (4) in (3) we obtain

$$\begin{aligned} n! |S_n(z)| &\geq (n!e^x/2) - (nc)^{1/2}(x + (c/2))^n \\ &\geq (e^x/2)(n! - 2(nc)^{1/2}(x + (c/2))^n e^{-x}). \end{aligned}$$

But

$$(x + (c/2))^n e^{-x} \leq e^{(c/2)}(n/e)^n,$$

while $n! > (2\pi n)^{1/2}(n/e)^n$, and so

$$n! |S_n(z)| \geq (e^x/2) n^{1/2}((2\pi)^{1/2} - 2c^{1/2}e^{c/2})(n/e)^n > 0.$$

Case (ii). $n < x$. It is an easy consequence of the Eneström–Kakeya theorem on polynomials with monotone coefficients (see [2]) that all zeros of $S_n(z)$ lie in $|z| \leq n$, and so the region $x > n$ is free of zeros. This simple observation due to a student of Richard Varga, W. Ni, replaces an elaborate discussion of this case that we had devised.

Thus, we have shown that if $y^2 \leq cx$, $S_n(x + iy) \neq 0$ for any n .

REFERENCES

1. D. J. NEWMAN AND T. J. RIVLIN, The zeros of partial sums of the exponential function, *J. Approximation Theory*, **5** (1972), 405–412.
2. G. PÓLYA AND G. SZEGÖ, "Aufgaben und Lehrsätze," Vol. 1, Abschn. III, No. 23, Springer-Verlag, Berlin, 1954.