

AN APPLICATION OF MILLER AND MOCANU LEMMA

HITOSHI SHIRAISHI AND SHIGEYOSHI OWA

ABSTRACT. Let $\mathcal{H}[a_0, n]$ be the class of functions $f(z) = a_0 + a_n z^n + \dots$ which are analytic in the open unit disk \mathbb{U} . For $f(z) \in \mathcal{H}[a_0, n]$, S. S. Miller and P. T. Mocanu (J. Math. Anal. Appl. **65**(1978), 289-305) have shown Miller and Mocanu lemma which is the generalization of Jack lemma by I. S. Jack (J. London Math. Soc. **3**(1971), 469-474). Applying Miller and Mocanu lemma, an interesting property for $f(z) \in \mathcal{H}[a_0, n]$ and an example are discuss.

1. INTRODUCTION

Let $\mathcal{H}[a_0, n]$ denote the class of functions $f(z)$ of the form

$$f(z) = a_0 + \sum_{k=n}^{\infty} a_k z^k \quad (n = 1, 2, 3, \dots)$$

which are analytic in the open unit disk $\mathbb{U} = \{z \in \mathbb{C} : |z| < 1\}$, where $a_0 \in \mathbb{C}$. Jack [1] has shown the result for analytic functions $w(z)$ in \mathbb{U} with $w(0) = 0$, which is called Jack's lemma. In 1978, Miller and Mocanu [2] have given the generalization theorem for Jack's lemma, which was called Miller and Mocanu lemma.

Lemma 1 (Miller and Mocanu lemma). *Let $f(z) \in \mathcal{H}[a_0, n]$ with $f(z) \neq a_0$. If there exists a point $z_0 \in \mathbb{U}$ such that*

$$\max_{|z| \leq |z_0|} |f(z)| = |f(z_0)|,$$

then

$$\frac{z_0 f'(z_0)}{f(z_0)} = m$$

and

$$\operatorname{Re} \left(\frac{z_0 f''(z_0)}{f'(z_0)} \right) + 1 \geq m,$$

where m is real and

$$m \geq n \frac{|f(z_0) - a_0|^2}{|f(z_0)|^2 - |a_0|^2} \geq n \frac{|f(z_0)| - |a_0|}{|f(z_0)| + |a_0|}.$$

If $a_0 = 0$, then the above lemma becomes Jack's lemma due to Jack [1].

2010 *Mathematics Subject Classification.* 30C45.

Key words and phrases. Analytic, univalent, Jack's lemma, Miller and Mocanu lemma.

2. MAIN THEOREM

Applying Miller and Mocanu lemma, we derive

Theorem 1. *Let $f(z) \in \mathcal{H}[a_0, n]$ with $f(z) \neq 0$ for $z \in \mathbb{U}$. If there exists a point $z_0 \in \mathbb{U}$ such that*

$$\min_{|z| \leq |z_0|} |f(z)| = |f(z_0)|,$$

then

$$\frac{z_0 f'(z_0)}{f(z_0)} = -m \quad (1)$$

and

$$\operatorname{Re} \left(\frac{z_0 f''(z_0)}{f'(z_0)} \right) + 1 \geq -m, \quad (2)$$

where

$$m \geq n \frac{|a_0 - f(z_0)|^2}{|a_0|^2 - |f(z_0)|^2} \geq n \frac{|a_0| - |f(z_0)|}{|a_0| + |f(z_0)|}.$$

Proof. We defined the function $g(z)$ by

$$\begin{aligned} g(z) &= \frac{1}{f(z)} \\ &= c_0 + c_n z^n + c_{n+1} z^{n+1} + \dots \quad \left(c_0 = \frac{1}{a_0} \right). \end{aligned}$$

Then, $g(z)$ is analytic in \mathbb{U} and $g(0) = c_0 \neq 0$. Furthermore, by the assumption of the theorem, $|g(z)|$ takes its maximum value at $z = z_0$ in the closed disk $|z| \leq |z_0|$. It follows from this that

$$|g(z_0)| = \frac{1}{|f(z_0)|} = \frac{1}{\min_{|z| \leq |z_0|} |f(z)|} = \max_{|z| \leq |z_0|} |g(z)|.$$

Therefore, applying Lemma 1 to $g(z)$, we observe that

$$\frac{z_0 g'(z_0)}{g(z_0)} = -\frac{z_0 f'(z_0)}{f(z_0)} = m$$

which shows (1) and

$$\begin{aligned} \operatorname{Re} \left(\frac{z_0 g''(z_0)}{g'(z_0)} \right) + 1 &= \operatorname{Re} \left(\frac{z_0 f''(z_0)}{f'(z_0)} - 2 \frac{z_0 f'(z_0)}{f(z_0)} \right) + 1 \\ &= \operatorname{Re} \left(\frac{z_0 f''(z_0)}{f'(z_0)} \right) + 2m + 1 \\ &\geq m \end{aligned}$$

which implies (2), where

$$m \geq n \frac{|g(z_0) - c_0|^2}{|g(z_0)|^2 - |c_0|^2} = n \frac{|a_0 - f(z_0)|^2}{|a_0|^2 - |f(z_0)|^2} \geq n \frac{|a_0| - |f(z_0)|}{|a_0| + |f(z_0)|}.$$

This completes the assertion of Theorem 1. \square

Example 1. Let us consider the function $f(z)$ given by

$$\begin{aligned} f(z) &= \frac{a_0 + (e^{i \arg(a_0)} - a_0) z^n}{1 - z^n} \\ &= a + e^{i \arg(a_0)} z^n + e^{i \arg(a_0)} z^{2n} + \dots \quad (z \in \mathbb{U}) \end{aligned}$$

for some complex number a_0 with $|a_0| > \frac{1}{2}$. Then, $f(z)$ maps the disk $\mathbb{U}_r = \{z : |z| < r \leq 1\}$ onto the domain

$$\left| f(z) - \left(a_0 + \frac{e^{i \arg(a_0)} r^{2n}}{1 - r^{2n}} \right) \right| \leq \frac{r^n}{1 - r^{2n}}.$$

Thus, we know that there exists a point $z_0 = r e^{i \frac{\pi}{n}} \in \mathbb{U}$ such that

$$\min_{|z| \leq |z_0|} |f(z)| = |f(z_0)| = |a_0| - \frac{r^n}{1 + r^n}.$$

For such a point z_0 , we obtain that

$$\frac{z_0 f'(z_0)}{f(z_0)} = -\frac{n r^n}{(1 + r^n)(|a_0| - (1 - |a_0|) r^n)} = -m$$

where

$$m = \frac{n r^n}{(1 + r^n)(|a_0| - (1 - |a_0|) r^n)} > 0.$$

Therefore, we get that

$$\operatorname{Re} \left(\frac{z_0 f''(z_0)}{f'(z_0)} \right) + 1 = n \frac{1 - r^n}{1 + r^n} > 0 > -m.$$

Furthermore, we obtain that

$$n \frac{|a_0 - f(z_0)|^2}{|a_0|^2 - |f(z_0)|^2} = \frac{n r^n}{2|a_0| + (2|a_0| - 1)r^n} = \frac{n r^n}{2 \left(|a_0| - (1 - |a_0|) r^n + \frac{1}{2} r^n \right)} < m.$$

Putting a_0 with a real number in Example 1, we get Example 2.

Example 2. Let us consider the function

$$\begin{aligned} f(z) &= \frac{a_0 + (1 - a) z^n}{1 - z^n} \\ &= a_0 + z^n + z^{2n} + \dots \quad (z \in \mathbb{U}) \end{aligned}$$

for $a_0 > \frac{1}{2}$. Then, it follows that the function $f(z)$ maps the disk \mathbb{U}_r onto the domain

$$\left| f(z) - \left(a_0 + \frac{r^{2n}}{1 - r^{2n}} \right) \right| \leq \frac{r^n}{1 - r^{2n}}.$$

Thus, there exists a point $z_0 = r e^{i \frac{\pi}{n}} \in \mathbb{U}$ such that

$$\min_{|z| \leq |z_0|} |f(z)| = |f(z_0)| = a_0 - \frac{r^n}{1 + r^n}.$$

For such a point z_0 , we obtain

$$\frac{z_0 f'(z_0)}{f(z_0)} = -\frac{n r^n}{(1 + r^n)(a_0 - (1 - a_0) r^n)} = -m$$

where

$$m = \frac{nr^n}{(1+r^n)(a_0 - (1-a_0)r^n)} > 0.$$

Therefore, we see that

$$\operatorname{Re} \left(\frac{z_0 f''(z_0)}{f'(z_0)} \right) + 1 = n \frac{1-r^n}{1+r^n} > 0 > -m.$$

Moreover, we have that

$$n \frac{|a_0 - f(z_0)|^2}{|a_0|^2 - |f(z_0)|^2} = \frac{nr^n}{2a_0 + (2a_0 - 1)r^n} = \frac{nr^n}{2 \left(a_0 - (1-a_0)r^n + \frac{1}{2}r^n \right)} < m.$$

REFERENCES

- [1] I. S. Jack, *Functions starlike and convex of order α* , J. London Math. Soc. **3**(1971), 469–474.
- [2] S. S. Miller and P. T. Mocanu, *Second-order differential inequalities in the complex plane*, J. Math. Anal. Appl. **65**(1978), 289–305.

HITOSHI SHIRAIISHI

DEPARTMENT OF MATHEMATICS

KINKI UNIVERSITY

HIGASHI-OSAKA, OSAKA 577-8502, JAPAN

E-mail address: shiraishi@math.kindai.ac.jp

SHIGEYOSHI OWA

DEPARTMENT OF MATHEMATICS

KINKI UNIVERSITY

HIGASHI-OSAKA, OSAKA 577-8502, JAPAN

E-mail address: owa@math.kindai.ac.jp