

## PROPERTIES OF CERTAIN ANALYTIC FUNCTIONS ASSOCIATED WITH TWO BOUNDARY POINTS

HITOSHI SHIRAISHI AND SHIGEYOSHI OWA

ABSTRACT. For analytic functions  $f(z)$  in the closed unit disk  $\bar{\mathbb{U}}$ , two boundary points  $z_1$  and  $z_2$  such that  $\alpha = (f'(z_1) + f'(z_2))/2 \in f'(\mathbb{U})$  are considered. The object of the present paper is to discuss some interesting conditions for  $f(z)$  to be  $|f'(z) - 1| < \rho|1 - \alpha|$  in  $\mathbb{U}$  with some examples.

### 1. INTRODUCTION

Let  $\mathcal{A}_n$  denote the class of functions

$$f(z) = z + a_{n+1}z^{n+1} + a_{n+2}z^{n+2} + \dots \quad (n = 1, 2, 3, \dots)$$

that are analytic in the closed unit disk  $\bar{\mathbb{U}} = \{z \in \mathbb{C} : |z| \leq 1\}$  and  $\mathcal{A} = \mathcal{A}_1$ . Also, the open unit disk is denoted by  $\mathbb{U} = \{z \in \mathbb{C} : |z| < 1\}$ .

For two boundary points  $z_1$  and  $z_2$ , let us consider

$$\alpha = \frac{f'(z_1) + f'(z_2)}{2} \in f'(\mathbb{U}).$$

For such  $\alpha$  ( $\alpha \neq 1$ ), if  $f(z) \in \mathcal{A}_n$  satisfies

$$\left| \frac{f'(z) - \alpha}{1 - \alpha} - 1 \right| < \rho \quad (z \in \mathbb{U})$$

for some real  $\rho > 0$ , then

$$|f'(z) - 1| < \rho|1 - \alpha| \quad (z \in \mathbb{U}).$$

Therefore, if  $0 < \rho|1 - \alpha| < 1$ , then  $f(z)$  is close-to-convex (univalent) in  $\mathbb{U}$ . In the present paper, we use such a technique for  $f(z)$ .

The basic tool in proving our results is the following lemma due to Jack [1] (also, due to Miller and Mocanu [2]).

**Lemma 1.** *Let the function  $w(z)$  defined by*

$$w(z) = a_n z^n + a_{n+1} z^{n+1} + a_{n+2} z^{n+2} + \dots \quad (n = 1, 2, 3, \dots)$$

---

2010 *Mathematics Subject Classification.* 30C45.  
*Key words and phrases.* Analytic, univalent, Jack's lemma.

be analytic in  $\mathbb{U}$  with  $w(0) = 0$ . If  $|w(z)|$  attains its maximum value on the circle  $|z| = r$  at a point  $z_0 \in \mathbb{U}$ , then there exists a real number  $k \geq n$  such that

$$\frac{z_0 w'(z_0)}{w(z_0)} = k$$

and

$$\operatorname{Re} \left( \frac{z_0 w''(z_0)}{w'(z_0)} \right) + 1 \geq k.$$

## 2. MAIN RESULTS

Applying Lemma 1, we derive the following results.

**Theorem 1.** *If  $f(z) \in \mathcal{A}_n$  satisfies*

$$\left| \frac{z f''(z)}{f'(z)} \right| < \frac{|1 - \alpha| n \rho}{1 + |1 - \alpha| \rho} \quad (z \in \mathbb{U})$$

for some complex  $\alpha = \frac{f'(z_1) + f'(z_2)}{2} \in f'(\mathbb{U})$  and  $\alpha \neq 1$  such that  $z_1, z_2 \in \partial\mathbb{U}$ , and for some real  $\rho > 1$ , then

$$|f'(z) - 1| < \rho |1 - \alpha| \quad (z \in \mathbb{U}).$$

*Proof.* Let us define  $w(z)$  by

$$\begin{aligned} w(z) &= \frac{f'(z) - \alpha}{1 - \alpha} - 1 \quad (z \in \mathbb{U}) \\ &= \frac{(n+1)a_{n+1}}{1 - \alpha} z^n + \frac{(n+2)a_{n+2}}{1 - \alpha} z^{n+1} + \dots \end{aligned} \quad (1)$$

Then, clearly,  $w(z)$  is analytic in  $\mathbb{U}$  and  $w(0) = 0$ . Differentiating both sides in (1), we obtain

$$\frac{z f''(z)}{f'(z)} = \frac{(1 - \alpha) z w'(z)}{(1 - \alpha) w(z) + 1},$$

and therefore,

$$\left| \frac{z f''(z)}{f'(z)} \right| = \left| \frac{(1 - \alpha) z w'(z)}{(1 - \alpha) w(z) + 1} \right| < \frac{|1 - \alpha| n \rho}{1 + |1 - \alpha| \rho} \quad (z \in \mathbb{U}).$$

If there exists a point  $z_0 \in \mathbb{U}$  such that

$$\max_{|z| \leq |z_0|} |w(z)| = |w(z_0)| = \rho,$$

then Lemma 1 gives us that  $w(z_0) = \rho e^{i\theta}$  and  $z_0 w'(z_0) = k w(z_0)$  ( $k \geq n$ ).

For such a point  $z_0$ , we have

$$\begin{aligned}
 & \left| \frac{z_0 f''(z_0)}{f'(z_0)} \right| \\
 &= \left| \frac{(1-\alpha)z_0 w'(z_0)}{(1-\alpha)w(z_0)+1} \right| \\
 &= \left| \frac{(1-\alpha)kw(z_0)}{(1-\alpha)w(z_0)+1} \right| \\
 &= \frac{|(1-\alpha)k\rho}{|(1-\alpha)\rho e^{i\theta}+1|} \\
 &= \frac{|(1-\alpha)k\rho}{\sqrt{(1+|1-\alpha|\rho \cos \phi)^2 + (|1-\alpha|\rho \sin \phi)^2}} \\
 &= \frac{|(1-\alpha)k\rho}{\sqrt{1+|1-\alpha|^2\rho^2+2|1-\alpha|\rho \cos \phi}} \\
 &\geq \frac{|1-\alpha|k\rho}{1+|1-\alpha|\rho} \\
 &\geq \frac{|1-\alpha|n\rho}{1+|1-\alpha|\rho}.
 \end{aligned}$$

where  $\phi = \theta + \arg(1-\alpha)$ .

This contradicts our condition in the theorem. Therefore, there is no  $z_0 \in \mathbb{U}$  such that  $|w(z_0)| = \rho$ . This means that  $|w(z)| < \rho$  for all  $z \in \mathbb{U}$ . It follows that

$$\left| \frac{f'(z) - \alpha}{1 - \alpha} - 1 \right| < \rho \quad (z \in \mathbb{U})$$

so that  $|f'(z) - 1| < \rho|1 - \alpha|$  in  $\mathbb{U}$ . □

**Example 1.** Let us consider a function

$$f(z) = z + a_{n+1}z^{n+1} \quad (z \in \mathbb{U})$$

with  $|a_{n+1}| < \frac{1}{2(n+1)}$ . Differentiating the function  $f(z)$ , we obtain

$$\frac{zf''(z)}{f'(z)} = \frac{n(n+1)a_{n+1}z^n}{1+(n+1)a_{n+1}z^n},$$

and therefore,

$$\begin{aligned}
 \left| \frac{zf''(z)}{f'(z)} \right| &= \left| \frac{n(n+1)a_{n+1}z^n}{1+(n+1)a_{n+1}z^n} \right| \\
 &< \frac{n(n+1)|a_{n+1}|}{1-(n+1)|a_{n+1}|} \quad (z \in \mathbb{U}).
 \end{aligned}$$

We choose two boundary points  $z_1$  and  $z_2$  such that  $f'(z_1) = 1 + (n+1)|a_{n+1}|$  and  $f'(z_2) = 1 + (n+1)|a_{n+1}|i$ . Then we know that  $z_1 = e^{-i\frac{\arg(a_{n+1})}{n}}$  and  $z_2 = e^{i\frac{\pi-2\arg(a_{n+1})}{2n}}$ . For such  $z_1$  and  $z_2$ , we obtain

$$\alpha = \frac{f'(z_1) + f'(z_2)}{2} = 1 + \frac{(n+1)|a_{n+1}|(1+i)}{2}$$

so that

$$1 - \alpha = -\frac{(n+1)|a_{n+1}|(1+i)}{2}.$$

Now, we consider some  $\rho > 1$  such that

$$\frac{n(n+1)|a_{n+1}|}{1 - (n+1)|a_{n+1}|} \leq \frac{|1 - \alpha|n\rho}{1 + |1 - \alpha|\rho} = \frac{n(n+1)|a_{n+1}|\rho}{\sqrt{2} + (n+1)|a_{n+1}|\rho}.$$

This gives us that

$$\rho \geq \frac{\sqrt{2}}{1 - 2(n+1)|a_{n+1}|}.$$

For such  $\alpha$  and  $\rho$ , we know that  $f(z)$  satisfies

$$|f'(z) - 1| < (n+1)|a_{n+1}| \leq \frac{(n+1)|a_{n+1}|}{1 - 2(n+1)|a_{n+1}|} \leq \rho|1 - \alpha|.$$

If we defined

$$\max_{z \in \mathbb{U}} |f'(z) - \alpha| = M_\alpha, \quad (2)$$

then we can have

$$\begin{aligned} |f'(z) - 1| &< \rho|1 - \alpha| \\ &= \rho|f'(0) - \alpha| \\ &\leq \rho M_\alpha \quad (z \in \mathbb{U}). \end{aligned}$$

So, we get

**Corollary 1.** *If  $f(z) \in \mathcal{A}_n$  satisfies*

$$\left| \frac{zf''(z)}{f'(z)} \right| < \frac{|1 - \alpha|n\rho}{1 + |1 - \alpha|\rho} \quad (z \in \mathbb{U})$$

for some complex  $\alpha = \frac{f'(z_1) + f'(z_2)}{2} \in f'(\mathbb{U})$  and  $\alpha \neq 1$  such that  $z_1, z_2 \in \partial\mathbb{U}$ , and for some real  $\rho > 1$ , then

$$|f'(z) - 1| < \rho M_\alpha \quad (z \in \mathbb{U}).$$

Putting

$$\alpha = \frac{f'(z_1) + f'(z_2) + \dots + f'(z_m)}{m} \quad (3)$$

for  $z_1, z_2, \dots, z_m \in \partial\mathbb{U}$ , we obtain

**Corollary 2.** *If  $f(z) \in \mathcal{A}_n$  satisfies*

$$\left| \frac{zf''(z)}{f'(z)} \right| < \frac{|1 - \alpha|n\rho}{1 + |1 - \alpha|\rho} \quad (z \in \mathbb{U})$$

for some complex  $\alpha = \frac{f'(z_1) + f'(z_2) + \dots + f'(z_m)}{m} \in f'(\mathbb{U})$  such that  $z_1, z_2, \dots, z_m \in \partial\mathbb{U}$  and  $\alpha \neq 1$ , and for some real  $\rho > 1$ , then

$$|f'(z) - 1| < \rho|1 - \alpha| \quad (z \in \mathbb{U}).$$

We also derive

**Theorem 2.** *If  $f(z) \in \mathcal{A}_n$  satisfies*

$$\left| z f''(z) - \frac{z f''(z)}{f'(z)} \right| < \frac{|1 - \alpha|^2 n \rho^2}{1 + |1 - \alpha| \rho} \quad (z \in \mathbb{U})$$

for some complex  $\alpha = \frac{f'(z_1) + f'(z_2)}{2} \in f'(\mathbb{U})$  and  $\alpha \neq 1$  such that  $z_1, z_2 \in \partial\mathbb{U}$ , and for some real  $\rho > 1$ , then

$$|f'(z) - 1| < \rho |1 - \alpha| \quad (z \in \mathbb{U}).$$

*Proof.* Define  $w(z)$  by

$$\begin{aligned} w(z) &= \frac{f'(z) - \alpha}{1 - \alpha} - 1 \quad (z \in \mathbb{U}) \\ &= \frac{(n+1)a_{n+1}}{1 - \alpha} z^n + \frac{(n+2)a_{n+2}}{1 - \alpha} z^{n+1} + \dots \end{aligned} \quad (4)$$

Evidently,  $w(z)$  is analytic in  $\mathbb{U}$  and  $w(0) = 0$ . Differentiating (4) logarithmically and simplifying, we have

$$\begin{aligned} z f''(z) - \frac{z f''(z)}{f'(z)} &= z f''(z) \left( 1 - \frac{1}{f'(z)} \right) \\ &= \frac{(1 - \alpha)^2 z w'(z) w(z)}{(1 - \alpha) w(z) + 1}, \end{aligned}$$

and hence,

$$\left| z f''(z) - \frac{z f''(z)}{f'(z)} \right| = \left| \frac{(1 - \alpha)^2 z w'(z) w(z)}{(1 - \alpha) w(z) + 1} \right| < \frac{|1 - \alpha|^2 n \rho^2}{1 + |1 - \alpha| \rho} \quad (z \in \mathbb{U}).$$

If there exists a point  $z_0 \in \mathbb{U}$  such that

$$\max_{|z| \leq |z_0|} |w(z)| = |w(z_0)| = \rho,$$

then Lemma 1 gives us that  $w(z_0) = \rho e^{i\theta}$  and  $z_0 w'(z_0) = k w(z_0)$  ( $k \geq n$ ).

For such a point  $z_0$ , we have

$$\begin{aligned}
& \left| z_0 f''(z_0) - \frac{z_0 f''(z_0)}{f'(z_0)} \right| \\
&= \left| \frac{(1-\alpha)^2 z_0 w'(z_0) w(z_0)}{(1-\alpha)w(z_0) + 1} \right| \\
&= \frac{|1-\alpha|^2 \rho^2 k}{|(1-\alpha)\rho e^{i\theta} + 1|} \\
&= \frac{|1-\alpha|^2 \rho^2 k}{\sqrt{(1+|1-\alpha|\rho \cos \phi)^2 + (|1-\alpha|\rho \sin \phi)^2}} \\
&= \frac{|1-\alpha|^2 \rho^2 k}{\sqrt{1+|1-\alpha|^2 \rho^2 + 2|1-\alpha|\rho \cos \phi}} \\
&\geq \frac{|1-\alpha|^2 k \rho^2}{1+|1-\alpha|\rho} \\
&\geq \frac{|1-\alpha|^2 n \rho^2}{1+|1-\alpha|\rho}.
\end{aligned}$$

where  $\phi = \theta + \arg(1-\alpha)$ .

This contradicts our condition in the theorem. Therefore, there is no  $z_0 \in \mathbb{U}$  such that  $|w(z_0)| = \rho$ . This means that  $|w(z)| < \rho$  for all  $z \in \mathbb{U}$ . This implies that

$$\left| \frac{f'(z) - \alpha}{1-\alpha} - 1 \right| < \rho \quad (z \in \mathbb{U}).$$

□

**Example 2.** We consider a function  $f(z)$  given by

$$f(z) = z + a_{n+1} z^{n+1} \quad (z \in \mathbb{U})$$

with  $|a_{n+1}| < \frac{1}{n+1}$ . Differentiating  $f(z)$ , we obtain that

$$z f''(z) - \frac{z f''(z)}{f'(z)} = \frac{n(n+1)^2 a_{n+1}^2 z^{2n}}{1+(n+1)a_{n+1}z^n},$$

that is, that

$$\begin{aligned}
\left| z f''(z) - \frac{z f''(z)}{f'(z)} \right| &= \left| \frac{n(n+1)^2 a_{n+1}^2 z^{2n}}{1+(n+1)a_{n+1}z^n} \right| \\
&< \frac{n(n+1)^2 |a_{n+1}|^2}{1-(n+1)|a_{n+1}|} \quad (z \in \mathbb{U}).
\end{aligned}$$

Choosing same points  $z_1$  and  $z_2$  in Example 1, we see that

$$1-\alpha = -\frac{(n+1)|a_{n+1}|(1+i)}{2}.$$

If we consider some  $\rho > 1$  such that

$$\frac{n(n+1)^2 |a_{n+1}|^2}{1-(n+1)|a_{n+1}|} \leq \frac{|1-\alpha|^2 n \rho^2}{1+|1-\alpha|\rho},$$

we have that

$$\rho \geq \frac{\sqrt{2}}{1 - (n+1)|a_{n+1}|}.$$

For such  $\alpha$  and  $\rho$ , we know that

$$|f'(z) - 1| < (n+1)|a_{n+1}| \leq \frac{(n+1)|a_{n+1}|}{1 - (n+1)|a_{n+1}|} \leq \rho|1 - \alpha|.$$

Further, we obtain

**Corollary 3.** *If  $f(z) \in \mathcal{A}_n$  satisfies*

$$\left| z f''(z) - \frac{z f''(z)}{f'(z)} \right| < \frac{|1 - \alpha|^2 n \rho^2}{1 + |1 - \alpha| \rho} \quad (z \in \mathbb{U})$$

for some complex  $\alpha = \frac{f'(z_1) + f'(z_2)}{2} \in f'(\mathbb{U})$  and  $\alpha \neq 1$  such that  $z_1, z_2 \in \partial\mathbb{U}$ , and for some real  $\rho > 1$ , then

$$|f'(z) - 1| < \rho M_\alpha \quad (z \in \mathbb{U}).$$

**Corollary 4.** *If  $f(z) \in \mathcal{A}_n$  satisfies*

$$\left| z f''(z) - \frac{z f''(z)}{f'(z)} \right| < \frac{|1 - \alpha|^2 n \rho^2}{1 + |1 - \alpha| \rho} \quad (z \in \mathbb{U})$$

for some complex  $\alpha = \frac{f'(z_1) + f'(z_2) + \dots + f'(z_m)}{m} \in f'(\mathbb{U})$  such that  $z_1, z_2, \dots, z_m \in \partial\mathbb{U}$  and  $\alpha \neq 1$ , and for some real  $\rho > 1$ , then

$$|f'(z) - 1| < \rho|1 - \alpha| \quad (z \in \mathbb{U}).$$

Further, we discuss a new application for Lemma 1.

**Theorem 3.** *If  $f(z) \in \mathcal{A}_n$  satisfies*

$$\operatorname{Re} \left( \frac{z(z f''(z))'}{f'(z) - 1} \right) < n^2 \quad (z \in \mathbb{U})$$

for some complex  $\alpha = \frac{f'(z_1) + f'(z_2)}{2} \in f'(\mathbb{U})$  and  $\alpha \neq 1$  such that  $z_1, z_2 \in \partial\mathbb{U}$ , and for some real  $\rho > 1$ , then

$$|f'(z) - 1| < \rho|1 - \alpha| \quad (z \in \mathbb{U}).$$

*Proof.* Defining the function  $w(z)$  by

$$\begin{aligned} w(z) &= \frac{f'(z) - \alpha}{1 - \alpha} - 1 \quad (z \in \mathbb{U}) \\ &= \frac{(n+1)a_{n+1}}{1 - \alpha} z^n + \frac{(n+2)a_{n+2}}{1 - \alpha} z^{n+1} + \dots, \end{aligned}$$

we have that  $w(z)$  is analytic in  $\mathbb{U}$  with  $w(0) = 0$ . Since,

$$\frac{z(zf''(z))'}{f'(z) - 1} = \frac{zw'(z)}{w(z)} + \frac{z^2w''(z)}{w(z)},$$

we obtain that

$$\begin{aligned} \operatorname{Re} \left( \frac{z(zf''(z))'}{f'(z) - 1} \right) &= \operatorname{Re} \left( \frac{zw'(z)}{w(z)} + \frac{z^2w''(z)}{w(z)} \right) \\ &= \operatorname{Re} \left( \frac{zw'(z)}{w(z)} \left( 1 + \frac{zw''(z)}{w'(z)} \right) \right) < n^2 \quad (z \in \mathbb{U}). \end{aligned}$$

If there exists a point  $z_0 \in \mathbb{U}$  such that

$$\max_{|z| \leq |z_0|} |w(z)| = |w(z_0)| = \rho,$$

then Lemma 1 gives us that  $w(z_0) = \rho e^{i\theta}$  and  $z_0w'(z_0) = kw(z_0)$  ( $k \geq n$ ).

Thus we have

$$\begin{aligned} \operatorname{Re} \left( \frac{z_0(z_0f''(z_0))'}{f'(z_0) - 1} \right) &= \operatorname{Re} \left( \frac{z_0w'(z_0)}{w(z_0)} \left( 1 + \frac{z_0w''(z_0)}{w'(z_0)} \right) \right) \\ &= k \left( 1 + \operatorname{Re} \left( \frac{z_0w''(z_0)}{w'(z_0)} \right) \right) \\ &= k^2 \\ &\geq n^2. \end{aligned}$$

This contradicts our condition in the theorem. Therefore, there is no  $z_0 \in \mathbb{U}$  such that  $|w(z_0)| = \rho$ . This means that  $|w(z)| < \rho$  for all  $z \in \mathbb{U}$ .  $\square$

We also have the following corollaries.

**Corollary 5.** *If  $f(z) \in \mathcal{A}_n$  satisfies*

$$\operatorname{Re} \left( \frac{z(zf''(z))'}{f'(z) - 1} \right) < n^2 \quad (z \in \mathbb{U})$$

*for some complex  $\alpha = \frac{f'(z_1) + f'(z_2)}{2} \in f'(\mathbb{U})$  and  $\alpha \neq 1$  such that  $z_1, z_2 \in \partial\mathbb{U}$ , and for some real  $\rho > 1$ , then*

$$|f'(z) - 1| < \rho M_\alpha \quad (z \in \mathbb{U}).$$

**Corollary 6.** *If  $f(z) \in \mathcal{A}_n$  satisfies*

$$\operatorname{Re} \left( \frac{z(zf''(z))'}{f'(z) - 1} \right) < n^2 \quad (z \in \mathbb{U})$$

*for some complex  $\alpha = \frac{f'(z_1) + f'(z_2) + \dots + f'(z_m)}{m} \in f'(\mathbb{U})$  such that  $z_1, z_2, \dots, z_m \in \partial\mathbb{U}$  and  $\alpha \neq 1$ , and for some real  $\rho > 1$ , then*

$$|f'(z) - 1| < \rho |1 - \alpha| \quad (z \in \mathbb{U}).$$



Next our result is contained in

**Theorem 4.** *If  $f(z) \in \mathcal{A}_n$  satisfies*

$$\frac{zf''(z)}{f'(z) - 1} \neq k \quad (z \in \mathbb{U})$$

for some complex  $\alpha = \frac{f'(z_1) + f'(z_2)}{2} \in f'(\mathbb{U})$  and  $\alpha \neq 1$  such that  $z_1, z_2 \in \partial\mathbb{U}$ , some real  $\rho > 1$ , and for all real  $k \geq n$ , then

$$|f'(z) - 1| < \rho|1 - \alpha| \quad (z \in \mathbb{U}).$$

*Proof.* Let us define the function  $w(z)$  by

$$\begin{aligned} w(z) &= \frac{f'(z) - \alpha}{1 - \alpha} - 1 \quad (z \in \mathbb{U}) \\ &= \frac{(n+1)a_{n+1}}{1 - \alpha} z^n + \frac{(n+2)a_{n+2}}{1 - \alpha} z^{n+1} + \dots \end{aligned} \quad (5)$$

Clearly,  $w(z)$  is analytic in  $\mathbb{U}$  with  $w(0) = 0$ . We want to prove that  $|w(z)| < \rho$  in  $\mathbb{U}$ . Note that

$$\frac{zf''(z)}{f'(z) - 1} = \frac{zw'(z)}{w(z)} \quad (z \in \mathbb{U}).$$

If there exists a point  $z_0 \in \mathbb{U}$  such that

$$\max_{|z| \leq |z_0|} |w(z)| = |w(z_0)| = \rho,$$

then Lemma 1 gives us that  $w(z_0) = \rho e^{i\theta}$  and  $z_0 w'(z_0) = kw(z_0)$  ( $k \geq n$ ).

Thus we have

$$\begin{aligned} \frac{z_0 f''(z_0)}{f'(z_0) - 1} &= \frac{z_0 w'(z_0)}{w(z_0)} \\ &= k \end{aligned}$$

This contradicts the condition in the theorem. Therefore, there is no  $z_0 \in \mathbb{U}$  such that  $|w(z_0)| = \rho$ . This means that  $|w(z)| < \rho$  for all  $z \in \mathbb{U}$ . We conclude that  $|f'(z) - 1| < \rho|1 - \alpha|$  in  $\mathbb{U}$ .  $\square$

**Corollary 7.** *If  $f(z) \in \mathcal{A}_n$  satisfies*

$$\frac{zf''(z)}{f'(z) - 1} \neq k \quad (z \in \mathbb{U})$$

for some complex  $\alpha = \frac{f'(z_1) + f'(z_2)}{2} \in f'(\mathbb{U})$  and  $\alpha \neq 1$  such that  $z_1, z_2 \in \partial\mathbb{U}$ , some real  $\rho > 1$ , and for all real  $k \geq n$ , then

$$|f'(z) - 1| < \rho M_\alpha \quad (z \in \mathbb{U}).$$

**Corollary 8.** *If  $f(z) \in \mathcal{A}_n$  satisfies*

$$\frac{zf''(z)}{f'(z)-1} \neq k \quad (z \in \mathbb{U})$$

*for some complex  $\alpha = \frac{f'(z_1) + f'(z_2) + \dots + f'(z_m)}{m} \in f'(\mathbb{U})$  such that  $z_1, z_2, \dots, z_m \in \partial\mathbb{U}$ , some real  $\rho > 1$ , and for all real  $k \geq n$ , then*

$$|f'(z) - 1| < \rho|1 - \alpha| \quad (z \in \mathbb{U}).$$

Finally, we derive

**Theorem 5.** *If  $f(z) \in \mathcal{A}_n$  satisfies*

$$\left| \frac{zf'(z)}{f(z)} - 1 \right| < \frac{|1 - \beta|n\rho}{1 + |1 - \beta|\rho} \quad (z \in \mathbb{U})$$

*for some complex  $\beta = \frac{F(z_1) + F(z_2)}{2} \in F(\mathbb{U})$  such that  $z_1, z_2 \in \partial\mathbb{U}$  and  $\beta \neq 1$ , and for some real  $\rho > 1$ , where  $F(z) = \frac{f(z)}{z}$ , then*

$$\left| \frac{f(z)}{z} - 1 \right| < \rho|1 - \beta| \quad (z \in \mathbb{U}).$$

*Proof.* Let us define  $w(z)$  by

$$\begin{aligned} w(z) &= \frac{\frac{f(z)}{z} - \beta}{1 - \beta} - 1 \quad (z \in \mathbb{U}) \\ &= \frac{a_{n+1}}{1 - \beta} z^n + \frac{a_{n+2}}{1 - \beta} z^{n+1} + \dots \end{aligned} \tag{6}$$

Then, we have that  $w(z)$  is analytic in  $\mathbb{U}$  and  $w(0) = 0$ . Differentiating (6) in both side logarithmically and simplifying, we obtain

$$\frac{zf'(z)}{f(z)} = \frac{(1 - \beta)zw'(z)}{(1 - \beta)w(z) + 1},$$

and hence,

$$\left| \frac{zf'(z)}{f(z)} \right| = \left| \frac{(1 - \beta)zw'(z)}{(1 - \beta)w(z) + 1} \right| < \frac{|1 - \beta|n\rho}{1 + |1 - \beta|\rho} \quad (z \in \mathbb{U}).$$

Using the same process of the proof in Theorem 1, we complete the proof of the theorem.  $\square$

**Example 3.** We consider a function

$$f(z) = z + a_{n+1}z^{n+1} \quad (z \in \mathbb{U})$$

with  $|a_{n+1}| < \frac{1}{2}$ . Differentiating the function, we obtain

$$\frac{zf'(z)}{f(z)} - 1 = \frac{na_{n+1}z^n}{1 + a_{n+1}z^n},$$

and therefore,

$$\begin{aligned} \left| \frac{zf'(z)}{f(z)} - 1 \right| &= \left| \frac{na_{n+1}z^n}{1+a_{n+1}z^n} \right| \\ &< \frac{n|a_{n+1}|}{1-|a_{n+1}|} \quad (z \in \mathbb{U}). \end{aligned}$$

Consider two boundary points  $z_1 = e^{-i\frac{\arg(a_{n+1})}{n}}$  and  $z_2 = e^{i\frac{\pi-2\arg(a_{n+1})}{2n}}$ . Then, since  $F(z_1) = 1 + |a_{n+1}|$  and  $F(z_2) = 1 + |a_{n+1}|i$ , we see that

$$1 - \beta = -\frac{|a_{n+1}|(1+i)}{2}.$$

For such  $\beta$ , we consider some  $\rho$  such that

$$\frac{n|a_{n+1}|}{1-|a_{n+1}|} \leq \frac{|1-\beta|n\rho}{1+|1-\beta|\rho}.$$

This gives us that

$$\rho \geq \frac{\sqrt{2}}{1-2|a_{n+1}|}.$$

Therefore, we have that

$$\left| \frac{f(z)}{z} - 1 \right| < |a_{n+1}| \leq \frac{|a_{n+1}|}{1-2|a_{n+1}|} \leq \rho|1-\beta|.$$

Defining  $M_\beta$  by

$$\max_{z \in \mathbb{U}} \left| \frac{f(z)}{z} - \beta \right| = M_\beta,$$

we have the following corollary.

**Corollary 9.** *If  $f(z) \in \mathcal{A}_n$  satisfies*

$$\left| \frac{zf'(z)}{f(z)} - 1 \right| < \frac{|1-\beta|n\rho}{1+|1-\beta|\rho} \quad (z \in \mathbb{U})$$

for some complex  $\beta = \frac{F(z_1)+F(z_2)}{2} \in F(\mathbb{U})$  such that  $z_1, z_2 \in \partial\mathbb{U}$  and  $\beta \neq 1$ , and

for some real  $\rho > 1$ , where  $F(z) = \frac{f(z)}{z}$ , then

$$\left| \frac{f(z)}{z} - 1 \right| < \rho M_\beta \quad (z \in \mathbb{U}).$$

Also considering

$$\beta = \frac{\frac{f(z_1)}{z_1} + \frac{f(z_2)}{z_2} + \dots + \frac{f(z_m)}{z_m}}{m}$$

for  $z_1, z_2, \dots, z_m \in \partial\mathbb{U}$ , we obtain

**Corollary 10.** *If  $f(z) \in \mathcal{A}_n$  satisfies*

$$\left| \frac{zf'(z)}{f(z)} - 1 \right| < \frac{|1-\beta|n\rho}{1+|1-\beta|\rho} \quad (z \in \mathbb{U})$$

for some complex  $\beta = \frac{F(z_1) + F(z_2) + \dots + F(z_m)}{m} \in F(\mathbb{U})$  such that  $z_1, z_2, \dots, z_m \in \partial\mathbb{U}$  and  $\beta \neq 1$ , and for some real  $\rho > 1$  where  $F(z) = \frac{f(z)}{z}$ , then

$$\left| \frac{f(z)}{z} - 1 \right| < \rho |1 - \beta| \quad (z \in \mathbb{U}).$$

## REFERENCES

- [1] I. S. Jack, *Functions starlike and convex of order  $\alpha$* , J. London Math. Soc. **3**(1971), 469–474.
- [2] S. S. Miller and P. T. Mocanu, *Second-order differential inequalities in the complex plane*, J. Math. Anal. Appl. **65**(1978), 289–305.
- [3] H. Shiraishi and S. Owa, *Some sufficient problems for certain univalent functions*, Far East J. Math. Sci. **30**(2008), 147–155.

HITOSHI SHIRAISHI  
 DEPARTMENT OF MATHEMATICS  
 KINKI UNIVERSITY  
 HIGASHI-OSAKA, OSAKA 577-8502, JAPAN  
*E-mail address:* shiraishi@math.kindai.ac.jp

SHIGEYOSHI OWA  
 DEPARTMENT OF MATHEMATICS  
 KINKI UNIVERSITY  
 HIGASHI-OSAKA, OSAKA 577-8502, JAPAN  
*E-mail address:* owa@math.kindai.ac.jp