

**STARLIKENESS AND CONVEXITY
FOR ANALYTIC FUNCTIONS
CONCERNED WITH JACK'S LEMMA**

HITOSHI SHIRAISHI AND SHIGEYOSHI OWA

ABSTRACT. There are many results for sufficient conditions of functions $f(z)$ which are analytic in the open unit disc \mathbb{U} to be starlike and convex in \mathbb{U} . The object of the present paper is to derive some interesting sufficient conditions for $f(z)$ to be starlike of order α and convex of order α in \mathbb{U} concerned with Jack's lemma. Some examples for our results are also considered with the help of Mathematica 5.2.

1. INTRODUCTION

Let \mathcal{A} denote the class of functions $f(z)$ that are analytic in the open unit disk $\mathbb{U} = \{z \in \mathbb{C} : |z| < 1\}$, so that $f(0) = f'(0) - 1 = 0$.

We denote by \mathcal{S} the subclass of \mathcal{A} consisting of univalent functions $f(z)$ in \mathbb{U} . Let $\mathcal{S}^*(\alpha)$ be the subclass of \mathcal{A} consisting of all functions $f(z)$ which satisfy

$$\operatorname{Re} \left(\frac{zf'(z)}{f(z)} \right) > \alpha \quad (z \in \mathbb{U})$$

for some $0 \leq \alpha < 1$. A function $f(z) \in \mathcal{S}^*(\alpha)$ is said to be starlike of order α in \mathbb{U} . We denote by $\mathcal{S}^* = \mathcal{S}^*(0)$.

Also, let $\mathcal{K}(\alpha)$ denote the subclass of \mathcal{A} consisting of functions $f(z)$ which satisfy

$$\operatorname{Re} \left(1 + \frac{zf''(z)}{f'(z)} \right) > \alpha \quad (z \in \mathbb{U})$$

for some $0 \leq \alpha < 1$. A function $f(z)$ in $\mathcal{K}(\alpha)$ is said to be convex of order α in \mathbb{U} . We say that $\mathcal{K} = \mathcal{K}(0)$. From the definitions for $\mathcal{S}^*(\alpha)$ and $\mathcal{K}(\alpha)$, we know that $f(z) \in \mathcal{K}(\alpha)$ if and only if $zf'(z) \in \mathcal{S}^*(\alpha)$.

Let $f(z)$ and $g(z)$ be analytic in \mathbb{U} . Then $f(z)$ is said to be subordinate to $g(z)$ if there exists an analytic function $w(z)$ in \mathbb{U} satisfying $w(0) = 0$, $|w(z)| < 1$ ($z \in \mathbb{U}$) and $f(z) = g(w(z))$.

We denote this subordination by

$$f(z) \prec g(z) \quad (z \in \mathbb{U}).$$

The basic tool in proving our results is the following lemma due to Jack [1] (also, due to Miller and Mocanu [2]).

2000 *Mathematics Subject Classification.* 30C45.

Key words and phrases. Analytic, univalent, starlike of order α , convex of order α .

Lemma 1. *Let $w(z)$ be analytic in \mathbb{U} with $w(0) = 0$. Then if $|w(z)|$ attains its maximum value on the circle $|z| = r$ at a point $z_0 \in \mathbb{U}$, then we have $z_0 w'(z_0) = kw(z_0)$, where $k \geq 1$ is a real number.*

2. MAIN RESULTS

Applying Lemma 1, we drive the following result for the class \mathcal{C} .

Theorem 1. *If $f(z) \in \mathcal{A}$ satisfies*

$$\operatorname{Re} \left(1 + \frac{zf''(z)}{f'(z)} \right) < \frac{\beta + 1}{2(\beta - 1)} \quad (z \in \mathbb{U})$$

for some real $2 \leq \beta < 3$, or

$$\operatorname{Re} \left(1 + \frac{zf''(z)}{f'(z)} \right) < \frac{5\beta - 1}{2(\beta + 1)} \quad (z \in \mathbb{U})$$

for some real $1 < \beta \leq 2$, then

$$\frac{zf'(z)}{f(z)} \prec \frac{\beta(1-z)}{\beta-z} \quad (z \in \mathbb{U})$$

and

$$\left| \frac{zf'(z)}{f(z)} - \frac{\beta}{\beta+1} \right| < \frac{\beta}{\beta+1} \quad (z \in \mathbb{U}).$$

This implies that $f(z) \in \mathcal{S}^*$ and $\int_0^z \frac{f(t)}{t} dt \in \mathcal{K}$.

Proof. Let us define the function $w(z)$ by

$$\frac{zf'(z)}{f(z)} = \frac{\beta(1-w(z))}{\beta-w(z)} \quad (w(z) \neq \beta).$$

Clearly, $w(z)$ is analytic in \mathbb{U} and $w(0) = 0$. We want to prove that $|w(z)| < 1$ in \mathbb{U} . Since

$$1 + \frac{zf''(z)}{f'(z)} = \frac{\beta(1-w(z))}{\beta-w(z)} - \frac{zw'(z)}{1-w(z)} + \frac{zw'(z)}{\beta-w(z)},$$

we see that

$$\begin{aligned} \operatorname{Re} \left(1 + \frac{zf''(z)}{f'(z)} \right) &= \operatorname{Re} \left(\frac{\beta(1-w(z))}{\beta-w(z)} - \frac{zw'(z)}{1-w(z)} + \frac{zw'(z)}{\beta-w(z)} \right) \\ &< \frac{\beta+1}{2(\beta-1)} \quad (z \in \mathbb{U}) \end{aligned}$$

for $2 \leq \beta < 3$ and

$$\begin{aligned} \operatorname{Re} \left(1 + \frac{zf''(z)}{f'(z)} \right) &= \operatorname{Re} \left(\frac{\beta(1-w(z))}{\beta-w(z)} - \frac{zw'(z)}{1-w(z)} + \frac{zw'(z)}{\beta-w(z)} \right) \\ &< \frac{5\beta-1}{2(\beta+1)} \quad (z \in \mathbb{U}) \end{aligned}$$

for $1 < \beta \leq 2$. If there exists a point $z_0 \in \mathbb{U}$ such that

$$\max_{|z| \leq |z_0|} |w(z)| = |w(z_0)| = 1,$$

then Lemma 1 gives us that $w(z_0) = e^{i\theta}$ and $z_0 w'(z_0) = kw(z_0)$ ($k \geq 1$). Thus we have

$$\begin{aligned} 1 + \frac{z_0 f''(z_0)}{f'(z_0)} &= \frac{\beta(1-w(z_0))}{\beta-w(z_0)} - \frac{z_0 w'(z_0)}{1-w(z_0)} + \frac{z_0 w'(z_0)}{\beta-w(z_0)} \\ &= \beta + \beta(1-\beta+k) \frac{1}{\beta-e^{i\theta}} - \frac{k}{1-e^{i\theta}}. \end{aligned}$$

It follows that

$$\begin{aligned} \operatorname{Re} \left(\frac{1}{\beta-w(z_0)} \right) &= \operatorname{Re} \left(\frac{1}{\beta-e^{i\theta}} \right) \\ &= \frac{1}{2\beta} + \frac{\beta^2-1}{2\beta(1+\beta^2-2\cos\theta)} \end{aligned}$$

and

$$\begin{aligned} \operatorname{Re} \left(\frac{1}{1-w(z_0)} \right) &= \operatorname{Re} \left(\frac{1}{1-e^{i\theta}} \right) \\ &= \frac{1}{2}. \end{aligned}$$

Therefore, we have

$$\operatorname{Re} \left(1 + \frac{z_0 f''(z_0)}{f'(z_0)} \right) = \frac{1+\beta}{2} + \frac{(\beta^2-1)(1-\beta+k)}{2(1+\beta^2-2\beta\cos\theta)}.$$

This implies that, for $2 \leq \beta < 3$,

$$\begin{aligned} \operatorname{Re} \left(1 + \frac{z_0 f''(z_0)}{f'(z_0)} \right) &\geq \frac{1+\beta}{2} + \frac{(\beta+1)(1-\beta+k)}{2(\beta-1)} \\ &\geq \frac{1+\beta}{2} + \frac{(\beta+1)(2-\beta)}{2(\beta-1)} \\ &= \frac{\beta+1}{2(\beta-1)} \end{aligned}$$

and, for $1 < \beta \leq 2$,

$$\begin{aligned} \operatorname{Re} \left(1 + \frac{z_0 f''(z_0)}{f'(z_0)} \right) &\geq \frac{1+\beta}{2} + \frac{(\beta-1)(1-\beta+k)}{2(\beta+1)} \\ &\geq \frac{1+\beta}{2} + \frac{(\beta-1)(2-\beta)}{2(\beta+1)} \\ &= \frac{5\beta-1}{2(\beta+1)}. \end{aligned}$$

This contradicts the condition in the theorem. Therefore, there is no $z_0 \in \mathbb{U}$ such that $|w(z_0)| = 1$. This means that $|w(z)| < 1$ for all $z \in \mathbb{U}$, so that

$$\frac{z f'(z)}{f(z)} \prec \frac{\beta(1-z)}{\beta-z} \quad (z \in \mathbb{U}).$$

Furthermore, since

$$w(z) = \frac{\beta \left(\frac{zf'(z)}{f(z)} - 1 \right)}{\frac{zf'(z)}{f(z)} - \beta} \quad (z \in \mathbb{U})$$

and $|w(z)| < 1$ ($z \in \mathbb{U}$), we conclude that

$$\left| \frac{zf'(z)}{f(z)} - \frac{\beta}{\beta+1} \right| < \frac{\beta}{\beta+1} \quad (z \in \mathbb{U}),$$

which implies that $f(z) \in \mathcal{S}^*$. Furthermore, we see that $f(z) \in \mathcal{S}^*$ if and only if $\int_0^z \frac{f(t)}{t} dt \in \mathcal{K}$. \square

Taking $\beta = 2$ in the theorem, we have following corollary due to Singh and Singh [3].

Corollary 1. *If $f(z) \in \mathcal{A}$ satisfies*

$$\operatorname{Re} \left(1 + \frac{zf''(z)}{f'(z)} \right) < \frac{3}{2} \quad (z \in \mathbb{U}),$$

then

$$\frac{zf'(z)}{f(z)} \prec \frac{2(1-z)}{2-z} \quad (z \in \mathbb{U})$$

and

$$\left| \frac{zf'(z)}{f(z)} - \frac{2}{3} \right| < \frac{3}{2} \quad (z \in \mathbb{U}).$$

With Theorem 1, we give the following example.

Example 1. For $2 \leq \beta < 3$, we consider the function $f(z)$ given by

$$f(z) = \frac{\beta-1}{2} \left(1 - (1-z)^{\frac{2}{\beta-1}} \right) \quad (z \in \mathbb{U}).$$

It follows that

$$\frac{zf'(z)}{f(z)} = \frac{2z(1-z)^{\frac{3-\beta}{\beta-1}}}{(\beta-1) \left(1 - (1-z)^{\frac{2}{\beta-1}} \right)} \quad (z \in \mathbb{U})$$

and

$$\begin{aligned} \operatorname{Re} \left(1 + \frac{zf''(z)}{f'(z)} \right) &= \operatorname{Re} \left(\frac{\beta-1-2z}{(\beta-1)(1-z)} \right) \\ &= \operatorname{Re} \left(\frac{2}{\beta-1} - \frac{3-\beta}{(\beta-1)(1-z)} \right) \\ &< \frac{\beta+1}{2(\beta-1)} \quad (z \in \mathbb{U}). \end{aligned}$$

Therefore, the function $f(z)$ satisfies the condition in Theorem 1. If we define the function $w(z)$ by

$$\frac{zf'(z)}{f(z)} = \frac{\beta(1-w(z))}{\beta-w(z)} \quad (w(z) \neq \beta),$$

then we see that $w(z)$ is analytic in \mathbb{U} , $w(0) = 0$ and $|w(z)| < 1$ ($z \in \mathbb{U}$) with Mathematica 5.2. This implies that

$$\frac{zf'(z)}{f(z)} \prec \frac{\beta(1-z)}{\beta-z} \quad (z \in \mathbb{U}).$$

For $1 < \beta \leq 2$, we consider

$$f(z) = \frac{\beta+1}{2(2\beta-1)} \left(1 - (1-z)^{\frac{2(2\beta-1)}{\beta+1}} \right) \quad (z \in \mathbb{U}).$$

Then we have that

$$\frac{zf'(z)}{f(z)} = \frac{2(2\beta-1)z(1-z)^{\frac{3(\beta-1)}{\beta+1}}}{(\beta+1) \left(1 - (1-z)^{\frac{2(2\beta-1)}{\beta+1}} \right)}$$

and

$$\operatorname{Re} \left(1 + \frac{zf''(z)}{f'(z)} \right) = \operatorname{Re} \left(\frac{\beta+1-2(2\beta-1)z}{(\beta+1)(1-z)} \right) < \frac{5\beta-1}{2(\beta+1)} \quad (z \in \mathbb{U}).$$

Thus, the function $f(z)$ satisfies the condition in Theorem 1. Define the function $w(z)$ by

$$\frac{zf'(z)}{f(z)} = \frac{\beta(1-w(z))}{\beta-w(z)} \quad (w(z) \neq \beta).$$

Then $w(z)$ is analytic in \mathbb{U} , $w(0) = 0$ and $|w(z)| < 1$ ($z \in \mathbb{U}$) with Mathematica 5.2. Therefore, we have that

$$\frac{zf'(z)}{f(z)} \prec \frac{\beta(1-z)}{\beta-z} \quad (z \in \mathbb{U}).$$

In particular, if we take $\beta = 2$ in this example, then $f(z)$ becomes

$$f(z) = z - \frac{1}{2}z^2 \in \mathcal{S}^*.$$

Theorem 2. *If $f(z) \in \mathcal{A}$ satisfies*

$$\operatorname{Re} \left(1 + \frac{zf''(z)}{f'(z)} \right) > -\frac{\beta+1}{2\beta(\beta-1)} \quad (z \in \mathbb{U})$$

for some real $\beta \leq -1$, or

$$\operatorname{Re} \left(1 + \frac{zf''(z)}{f'(z)} \right) > \frac{3\beta+1}{2\beta(\beta+1)} \quad (z \in \mathbb{U})$$

for some real $\beta > 1$, then

$$\frac{f(z)}{zf'(z)} \prec \frac{\beta(1-z)}{\beta-z} \quad (z \in \mathbb{U})$$

and

$$f(z) \in \mathcal{S}^* \left(\frac{\beta+1}{2\beta} \right).$$

This implies that $\int_0^z \frac{f(t)}{t} dt \in \mathcal{K} \left(\frac{\beta+1}{2\beta} \right)$.

Proof. Let us define the function $w(z)$ by

$$\frac{f(z)}{zf'(z)} = \frac{\beta(1-w(z))}{\beta-w(z)} \quad (w(z) \neq \beta). \quad (1)$$

Then, we have that $w(z)$ is analytic in \mathbb{U} and $w(0) = 0$. We want to prove that $|w(z)| < 1$ in \mathbb{U} . Differentiating (1) in both side logarithmically and simplifying, we obtain

$$1 + \frac{zf''(z)}{f'(z)} = \frac{\beta-w(z)}{\beta(1-w(z))} + \frac{zw'(z)}{1-w(z)} - \frac{zw'(z)}{\beta-w(z)},$$

and, hence

$$\begin{aligned} \operatorname{Re} \left(1 + \frac{zf''(z)}{f'(z)} \right) &= \operatorname{Re} \left(\frac{\beta-w(z)}{\beta(1-w(z))} + \frac{zw'(z)}{1-w(z)} - \frac{zw'(z)}{\beta-w(z)} \right) \\ &> -\frac{\beta+1}{2\beta(\beta-1)} \quad (z \in \mathbb{U}) \end{aligned}$$

for $\beta \leq -1$, or

$$\begin{aligned} \operatorname{Re} \left(1 + \frac{zf''(z)}{f'(z)} \right) &= \operatorname{Re} \left(\frac{\beta-w(z)}{\beta(1-w(z))} + \frac{zw'(z)}{1-w(z)} - \frac{zw'(z)}{\beta-w(z)} \right) \\ &> \frac{3\beta+1}{2\beta(\beta+1)} \quad (z \in \mathbb{U}) \end{aligned}$$

for $\beta > 1$. If there exists a point $z_0 \in \mathbb{U}$ such that

$$\max_{|z| \leq |z_0|} |w(z)| = |w(z_0)| = 1,$$

then Lemma 1 gives us that $w(z_0) = e^{i\theta}$ and $z_0 w'(z_0) = kw(z_0)$ ($k \geq 1$). Thus we have

$$\begin{aligned} 1 + \frac{z_0 f''(z_0)}{f'(z_0)} &= \frac{\beta-w(z_0)}{\beta(1-w(z_0))} + \frac{z_0 w'(z_0)}{1-w(z_0)} - \frac{z_0 w'(z_0)}{\beta-w(z_0)} \\ &= \frac{1}{\beta} + \frac{\beta-1}{\beta(1-e^{i\theta})} + \frac{k}{1-e^{i\theta}} - \frac{k\beta}{\beta-e^{i\theta}}. \end{aligned}$$

Therefore, we have

$$\operatorname{Re} \left(1 + \frac{z_0 f''(z_0)}{f'(z_0)} \right) = \frac{1}{2} + \frac{1}{2\beta} - \frac{k(\beta^2-1)}{2(1+\beta^2-2\beta \cos \theta)}.$$

This implies that, for $\beta \leq -1$,

$$\begin{aligned} \operatorname{Re} \left(1 + \frac{z_0 f''(z_0)}{f'(z_0)} \right) &\leq \frac{1}{2} + \frac{1}{2\beta} - \frac{k(\beta+1)}{2(\beta-1)} \\ &\leq \frac{1}{2} + \frac{1}{2\beta} - \frac{\beta+1}{2(\beta-1)} \\ &= -\frac{\beta+1}{2\beta(\beta-1)}. \end{aligned}$$

and, for $\beta > 1$,

$$\begin{aligned} \operatorname{Re} \left(1 + \frac{z_0 f''(z_0)}{f'(z_0)} \right) &\leq \frac{1}{2} + \frac{1}{2\beta} - \frac{k(\beta-1)}{2(\beta+1)} \\ &\leq \frac{1}{2} + \frac{1}{2\beta} - \frac{\beta-1}{2(\beta+1)} \\ &= \frac{3\beta+1}{2\beta(\beta+1)}. \end{aligned}$$

This contradicts the condition in the theorem. Therefore, there is no $z_0 \in \mathbb{U}$ such that $|w(z_0)| = 1$. This means that $|w(z)| < 1$ for all $z \in \mathbb{U}$, this is, that

$$\frac{f(z)}{zf'(z)} \prec \frac{\beta(1-z)}{\beta-z} \quad (z \in \mathbb{U}).$$

Furthermore, since

$$w(z) = \frac{\beta \left(1 - \frac{zf'(z)}{f(z)} \right)}{1 - \beta \frac{zf'(z)}{f(z)}} \quad (z \in \mathbb{U})$$

and $|w(z)| < 1$ ($z \in \mathbb{U}$), we conclude that

$$f(z) \in \mathcal{S}^* \left(\frac{\beta+1}{2\beta} \right).$$

Noting that $f(z) \in \mathcal{S}^*(\alpha)$ if and only if $\int_0^z \frac{f(t)}{t} dt \in \mathcal{K}(\alpha)$, we complete the proof of the theorem. \square

For Theorem 2, we give the following example.

Example 2. For $\beta > 1$, we take

$$f(z) = \frac{\beta(\beta+1)}{-\beta^2+2\beta+1} \left(1 - (1-z)^{\frac{-\beta^2+2\beta+1}{\beta(\beta+1)}} \right) \quad (z \in \mathbb{U}).$$

Then, $f(z)$ satisfies

$$\frac{zf'(z)}{f(z)} = \frac{(-\beta^2+2\beta+1)z}{\beta(\beta+1)(1-z)^{\frac{2\beta^2-\beta-1}{\beta(\beta+1)}} \left(1 - (1-z)^{\frac{-\beta^2+2\beta+1}{\beta(\beta+1)}} \right)}$$

and

$$\begin{aligned} \operatorname{Re} \left(1 + \frac{zf''(z)}{f'(z)} \right) &= \operatorname{Re} \left(\frac{\beta(\beta+1) + (\beta^2 - 2\beta - 1)z}{\beta(\beta+1)(1-z)} \right) \\ &> \frac{3\beta+1}{2\beta(\beta+1)} \quad (z \in \mathbb{U}). \end{aligned}$$

Therefore, $f(z)$ satisfies the condition of Theorem 2. Let us define the function $w(z)$ by

$$\frac{f(z)}{zf'(z)} = \frac{\beta(1-w(z))}{\beta-w(z)} \quad (w(z) \neq \beta).$$

Then $w(z)$ is analytic in \mathbb{U} , $w(0) = 0$ and $|w(z)| < 1$ $z \in \mathbb{U}$ with Mathematica 5.2. It follows that

$$\frac{f(z)}{zf'(z)} \prec \frac{\beta(1-z)}{\beta-z} \quad (z \in \mathbb{U}).$$

Furthermore, for $\beta \leq -1$, we consider the following function

$$f(z) = -\frac{\beta(\beta-1)}{\beta^2+1} \left(1 - (1-z)^{-\frac{\beta^2+1}{\beta(\beta-1)}} \right).$$

Note that

$$\frac{zf'(z)}{f(z)} = \frac{-(\beta^2+1)z}{\beta(\beta-1)(1-z)^{\frac{2\beta^2-\beta+1}{\beta(\beta-1)}} \left(1 - (1-z)^{-\frac{\beta^2+1}{\beta(\beta-1)}} \right)}$$

and

$$\begin{aligned} \operatorname{Re} \left(1 + \frac{zf''(z)}{f'(z)} \right) &= \operatorname{Re} \left(\frac{\beta(\beta-1) + (\beta^2+1)z}{\beta(\beta-1)(1-z)} \right) \\ &> \frac{\beta+1}{2\beta(\beta-1)} \quad (z \in \mathbb{U}). \end{aligned}$$

This implies that $f(z)$ satisfies the condition of Theorem 2. Defining the function $w(z)$ by

$$\frac{f(z)}{zf'(z)} = \frac{\beta(1-w(z))}{\beta-w(z)} \quad (w(z) \neq \beta),$$

we see that $w(z)$ is analytic in \mathbb{U} , $w(0) = 0$ and $|w(z)| < 1$ ($z \in \mathbb{U}$) with Mathematica 5.2.

Thus we have that

$$\frac{f(z)}{zf'(z)} \prec \frac{\beta(1-z)}{\beta-z} \quad (z \in \mathbb{U}).$$

Making $\beta = -1$ for $f(z)$, we have

$$f(z) = \frac{z}{1-z} \in \mathcal{K}.$$

REFERENCES

- [1] I. S. Jack, *Functions starlike and convex of order α* , J. London Math. Soc. **3**(1971), 469–474.
- [2] S. S. Miller and P. T. Mocanu, *Second-order differential inequalities in the complex plane*, J. Math. Anal. Appl. **65**(1978), 289–305.
- [3] R. Singh and S. Singh, *Some sufficient conditions for univalence and starlikeness*, Coll. Math. **47**(1982), 309–314.

HITOSHI SHIRAISHI

DEPARTMENT OF MATHEMATICS

KINKI UNIVERSITY

HIGASHI-OSAKA, OSAKA 577-8502, JAPAN

E-mail address: 0733310104x@kindai.ac.jp

SHIGEYOSHI OWA

DEPARTMENT OF MATHEMATICS

KINKI UNIVERSITY

HIGASHI-OSAKA, OSAKA 577-8502, JAPAN

E-mail address: owa@math.kindai.ac.jp