

**STARLIKENESS PROBLEMS  
FOR CERTAIN ANALYTIC FUNCTIONS  
CONCERNED WITH SUBORDINATIONS**

HITOSHI SHIRAISHI, SHIGEYOSHI OWA, TOSHIO HAYAMI,  
KAZUO KUROKI AND H. M. SRIVASTAVA

ABSTRACT. Let  $\mathcal{A}_n$  be the class of functions  $f(z)$  which are analytic in the open unit disk  $\mathbb{U}$  with  $f(0) = 0$ ,  $f'(0) = 1$ ,  $f''(0) = f'''(0) = \dots = f^{(n)}(0) = 0$  and  $f^{(n+1)} \neq 0$ . Applying the results due to S. S. Miller (J. Math. Anal. Appl. **65**(1978), 289-305), some interesting starlikeness problems concerned with subordinations are discussed. The results in the paper are extensions of results by M. Obradović (Hokkaido Math. J. **27**(1998), 329-335).

1. INTRODUCTION

Let  $\mathcal{H}[a_0, n]$  denote the class of functions  $p(z)$  of the form

$$p(z) = a_0 + \sum_{k=n}^{\infty} a_k z^k \quad (n = 1, 2, 3, \dots)$$

which are analytic in the open unit disk  $\mathbb{U} = \{z \in \mathbb{C} : |z| < 1\}$ , where  $a_0 \in \mathbb{C}$ .

Also, let  $\mathcal{A}_n$  denote the class of functions

$$f(z) = z + a_{n+1}z^{n+1} + a_{n+2}z^{n+2} + \dots \quad (n = 1, 2, 3, \dots)$$

that are analytic in  $\mathbb{U}$  with  $a_{n+1} \neq 0$  and  $\mathcal{A} \equiv \mathcal{A}_1$ .

If  $f(z) \in \mathcal{A}_n$  satisfies

$$\operatorname{Re} \left( \frac{zf'(z)}{f(z)} \right) > \alpha \quad (z \in \mathbb{U})$$

for some real  $\alpha$  ( $0 \leq \alpha < 1$ ), then we say that  $f(z)$  is starlike of order  $\alpha$  and written by  $f(z) \in \mathcal{S}^*(\alpha)$  and  $\mathcal{S}^* \equiv \mathcal{S}^*(0)$ .

Let  $f(z)$  and  $g(z)$  be analytic in  $\mathbb{U}$ . Then  $f(z)$  is said to be subordinate to  $g(z)$  if there exists an analytic function  $w(z)$  in  $\mathbb{U}$  satisfying  $w(0) = 0$ ,  $|w(z)| < 1$  ( $z \in \mathbb{U}$ ) and such that  $f(z) = g(w(z))$ . We denote this subordination by

$$f(z) \prec g(z) \quad (z \in \mathbb{U}).$$

In particular, if  $g(z)$  is univalent in  $\mathbb{U}$ , then the subordination

$$f(z) \prec g(z) \quad (z \in \mathbb{U})$$

---

2010 *Mathematics Subject Classification.* 30C45.

*Key words and phrases.* Analytic function, univalent function, subordination, strongly starlike function.

is equivalent to  $f(0) = g(0)$  and  $f(\mathbb{U}) \subset g(\mathbb{U})$  (cf. [3]).

The basic tool in proving our results is the following lemma due to Miller and Mocanu [2] (also [3]).

**Lemma 1.** *Let the function  $w(z)$  defined by*

$$w(z) = a_n z^n + a_{n+1} z^{n+1} + a_{n+2} z^{n+2} + \dots \quad (n = 1, 2, 3, \dots)$$

*be analytic in  $\mathbb{U}$  with  $w(0) = 0$ . If  $|w(z)|$  attains its maximum value on the circle  $|z| = r$  at a point  $z_0 \in \mathbb{U}$ , then there exists a real number  $k \geq n$  such that*

$$\frac{z_0 w'(z_0)}{w(z_0)} = k.$$

## 2. MAIN RESULT

Applying Lemma 1, we have the following lemma.

**Lemma 2.** *Let  $p(z) \in \mathcal{H}[1, n]$  satisfy the condition*

$$p(z) - \frac{1}{\mu} z p'(z) \prec 1 + \lambda z \quad (z \in \mathbb{U})$$

*for some complex number  $\mu$  ( $\operatorname{Re}(\mu) < n, \mu \neq 0$ ) and some complex number  $\lambda$  ( $0 < |\lambda| \leq 1$ ), then*

$$p(z) \prec 1 + \lambda_1 z \quad (z \in \mathbb{U}),$$

*where  $\lambda_1$  is a complex number such that*

$$|\lambda_1| = |\lambda| \frac{|\mu|}{|n - \mu|}. \quad (2.1)$$

*Proof.* We consider the function  $p(z)$  defined by

$$p(z) = 1 + \lambda_1 w(z)$$

with  $\lambda_1$  is given by (2.1). Then,  $w(z)$  is analytic in  $\mathbb{U}$  and  $w(0) = 0$ . We want to show that  $|w(z)| < 1$  ( $z \in \mathbb{U}$ ). If there exists a point  $z_0 \in \mathbb{U}$  such that  $|w(z_0)| = 1$ , then we can write  $z_0 w'(z_0) = k w(z_0)$  ( $k \geq n$ ) by Lemma 1. Putting  $w(z_0) = e^{i\theta}$ ,

we get

$$\begin{aligned}
\left| p(z_0) - \frac{1}{\mu} z_0 p'(z_0) - 1 \right| &= \left| \lambda_1 w(z_0) - \frac{1}{\mu} \lambda_1 z_0 w'(z_0) \right| \\
&= \left| \lambda_1 e^{i\theta} - \frac{1}{\mu} \lambda_1 k e^{i\theta} \right| \\
&= |\lambda_1| \frac{|\mu - k|}{|\mu|} \\
&= |\lambda_1| \frac{\sqrt{(k - \operatorname{Re}(\mu))^2 + (\operatorname{Im}(\mu))^2}}{|\mu|} \\
&\geq |\lambda_1| \frac{\sqrt{(n - \operatorname{Re}(\mu))^2 + (\operatorname{Im}(\mu))^2}}{|\mu|} \\
&= |\lambda_1| \frac{|n - \mu|}{|\mu|} = |\lambda|.
\end{aligned}$$

This contradicts the assumption of Lemma 2. Therefore, there is no  $z_0 \in \mathbb{U}$  such that  $|w(z_0)| = 1$ . This implies that  $|w(z)| < 1$  ( $z \in \mathbb{U}$ ). This completes the proof of the lemma.  $\square$

Next, we show

**Lemma 3.** *Let  $\lambda$  and  $\lambda_1$  be complex numbers such that  $0 < |\lambda_1| < |\lambda| < 1$  and let  $Q(z) \in \mathcal{H}[1, n]$  such that*

$$Q(z) \prec 1 + \lambda_1 z \quad (z \in \mathbb{U}). \quad (2.2)$$

(1) *If  $p(z) \in \mathcal{H}[1, n]$  and*

$$Q(z)[\alpha + (1 - \alpha)p(z)] \prec 1 + \lambda z \quad (z \in \mathbb{U}) \quad (2.3)$$

*for some real  $\alpha$  such that*

$$\alpha \leq \begin{cases} \frac{1 - |\lambda|}{1 + |\lambda_1|} & (0 < |\lambda| + |\lambda_1| \leq 1) \\ \frac{1 - (|\lambda|^2 + |\lambda_1|^2)}{2(1 - |\lambda_1|^2)} & (|\lambda|^2 + |\lambda_1|^2 \leq 1 \leq |\lambda| + |\lambda_1|), \end{cases} \quad (2.4)$$

*then  $\operatorname{Re}(p(z)) > 0$ .*

(2) *If  $w(z) \in \mathcal{H}[0, n]$  and*

$$Q(z)[1 + w(z)] \prec 1 + \lambda z \quad (z \in \mathbb{U}), \quad (2.5)$$

*then*

$$|w(z)| < \frac{|\lambda| + |\lambda_1|}{1 - |\lambda_1|} \leq 1, \quad (2.6)$$

*where  $|\lambda| + 2|\lambda_1| \leq 1$ .*

*The bound (2.6) and the value of  $\alpha$  given by (2.4) are best possible.*

*Proof.* To prove (1), we think about

$$g(z) = \frac{1 + \lambda z}{1 + \lambda_1 e^{i\theta} z} \quad (z \in \mathbb{U})$$

for some real  $\theta$ .

It follows that

$$\left| g(z) - \frac{1 - \lambda \bar{\lambda}_1 r^2 e^{-i\theta}}{1 - |\lambda_1|^2 r^2} \right| \leq \frac{|\lambda - \lambda_1 e^{i\theta}| r}{1 - |\lambda_1|^2 r^2}$$

for  $|z| \leq r < 1$ .

If we put  $\lambda = |\lambda| e^{i\theta_0}$ ,  $\lambda_1 = |\lambda_1| e^{i\theta_1}$  and  $\phi = \theta_0 - \theta_1 - \theta$ , we can rewrite

$$\left| g(z) - \frac{1 - |\lambda| |\lambda_1| r^2 e^{i\phi}}{1 - |\lambda_1|^2 r^2} \right| \leq \frac{||\lambda_1| - |\lambda| e^{i\phi}| r}{1 - |\lambda_1|^2 r^2} \quad (|z| \leq r < 1).$$

Hence for all  $r$  ( $0 < r < 1$ ), we obtain

$$\operatorname{Re}(g(z)) > \frac{1 - |\lambda| |\lambda_1| \cos \phi - \sqrt{|\lambda|^2 - 2|\lambda| |\lambda_1| \cos \phi + |\lambda_1|^2}}{1 - |\lambda_1|^2} \equiv h(\phi).$$

Note that

$$h'(\phi) = \frac{|\lambda| |\lambda_1| \sin \phi}{1 - |\lambda_1|^2} \left( 1 - \frac{1}{\sqrt{|\lambda|^2 - 2|\lambda| |\lambda_1| \cos \phi + |\lambda_1|^2}} \right).$$

Therefore,  $h'(\phi) = 0$  if  $\sin(\phi) = 0$  or  $\sqrt{|\lambda|^2 - 2|\lambda| |\lambda_1| \cos \phi + |\lambda_1|^2} = 1$ . Since

$$|\lambda|^2 - 2|\lambda| |\lambda_1| \cos \phi + |\lambda_1|^2 \leq (|\lambda| + |\lambda_1|)^2,$$

if  $0 < |\lambda| + |\lambda_1| \leq 1$ , then  $h'(\phi) = 0$  for  $\phi = 2k\pi$  ( $k = 0, \pm 1, \pm 2, \dots$ ) and  $\phi = (2k+1)\pi$  ( $k = 0, \pm 1, \pm 2, \dots$ ). With this conditions, we see that

$$h(\phi) \geq h(2k+1)\pi = \frac{1 - |\lambda|}{1 + |\lambda_1|} \geq \alpha.$$

If  $\sqrt{|\lambda|^2 - 2|\lambda| |\lambda_1| \cos \phi + |\lambda_1|^2} = 1$ , then  $h'(\phi) = 0$  for  $\phi = \phi_1$  such that

$$|\lambda|^2 - |\lambda| |\lambda_1| \cos \phi_1 + |\lambda_1|^2 = 1.$$

Thus, we have that

$$h(\phi) \geq h(\phi_1) = \frac{1 - (|\lambda|^2 + |\lambda_1|^2)}{2(1 - |\lambda_1|^2)} \geq \alpha$$

with  $|\lambda|^2 + |\lambda_1|^2 \leq 1 \leq |\lambda| + |\lambda_1|$ .

On the other hand, in view of (2.2) and (2.3), we have

$$\alpha + (1 - \alpha)p(z) \prec g(z) \quad (z \in \mathbb{U}).$$

So, we can lead  $\operatorname{Re}(p(z)) > 0$ .

To prove (2), note that in view of (2.2) and (2.5) we have

$$\begin{aligned} |w(z)| &= \left| \frac{(Q(z)(1+w(z))-1) - (Q(z)-1)}{Q(z)} \right| \\ &\leq \frac{|Q(z)(1+w(z))-1| + |Q(z)-1|}{|Q(z)|} \\ &\leq \frac{|\lambda| + |\lambda_1|}{|1 + \lambda_1 z|} \\ &< \frac{|\lambda| + |\lambda_1|}{1 - |\lambda_1|} \leq 1 \quad (z \in \mathbb{U}). \end{aligned}$$

Let us show the sharpness.

We take

$$Q(z) = 1 + \lambda_1 w_1(z) \quad (z \in \mathbb{U})$$

where  $w_1(z)$  is analytic in  $\mathbb{U}$  such that  $w_1(0) = 0$  and  $|w_1(z)| < 1$  ( $z \in \mathbb{U}$ ).

In the case of (1), we put

$$Q(z)[\alpha + (1-\alpha)p(z)] = 1 + \lambda w_0(z) \quad (z \in \mathbb{U})$$

where  $w_0(z)$  is analytic in  $\mathbb{U}$  such that  $w_0(0) = 0$  and  $|w_0(z)| < 1$  ( $z \in \mathbb{U}$ ). Since

$$p(z) = \frac{1}{1-\alpha} \left( \frac{1 + \lambda w_0(z)}{1 + \lambda_1 w_1(z)} - \alpha \right),$$

if we consider  $w_0(z)$  such that  $\lambda w_0(z) = -|\lambda|$  and  $\lambda_1 w_1(z) = |\lambda_1|$ , then we have

$$\alpha = \frac{1 - |\lambda|}{1 + |\lambda_1|} \quad (0 < |\lambda| + |\lambda_1| \leq 1).$$

Furthermore, if we take  $w_0(z)$  such that  $\lambda w_0(z) = |\lambda|e^{i\theta_0}$  and  $w_1(z)$  such that  $\lambda_1 w_1(z) = |\lambda_1|e^{i\theta_1}$ , then we obtain that

$$\alpha = \frac{1 - (|\lambda|^2 + |\lambda_1|^2)}{2(1 - |\lambda_1|^2)}$$

with

$$|\lambda| \sin \theta_0 - |\lambda_1| \sin \theta_1 + |\lambda||\lambda_1| \sin(\theta_0 - \theta_1) = 0$$

and

$$\begin{aligned} &2(1 - |\lambda_1|^2)|\lambda| \cos \theta_0 + 2|\lambda|^2|\lambda_1| \cos \theta_1 + 2(1 - |\lambda_1|^2)|\lambda||\lambda_1| \cos(\theta_0 - \theta_1) \\ &= (|\lambda_1|^2 - 1)(1 + |\lambda|^2 - |\lambda_1|^2). \end{aligned}$$

In the case of (2), we put

$$Q(z)[1 + w(z)] = 1 + \lambda w_0(z) \quad (z \in \mathbb{U})$$

where  $w_0(z)$  that  $w_0(0) = 0$  and  $|w_0(z)| < 1$  ( $z \in \mathbb{U}$ ).

Then, we can obtain

$$w(z) = \frac{\lambda w_0(z) - \lambda_1 w_1(z)}{1 + \lambda w_0(z)} \quad (z \in \mathbb{U}).$$

If we take

$$w(z) = \frac{|\lambda| + |\lambda_1|}{1 - |\lambda_1|} \quad (z \in \mathbb{U}),$$

then we have  $\lambda w_0(z) = |\lambda|$  and  $\lambda_1 w_1(z) = -|\lambda_1|$ . □

If we consider some real  $\lambda$ ,  $\lambda_1$  and fixed  $\alpha$  in Lemma 3, we obtain Corollary 1 due to S. Ponnusamy and V. Singh [5].

**Corollary 1.** *Let  $\lambda$  and  $\lambda_1$  be real with  $0 < \lambda_1 < \lambda < 1$  and let  $Q(z) \in \mathcal{H}[1, n]$  satisfy*

$$Q(z) \prec 1 + \lambda_1 z \quad (z \in \mathbb{U}).$$

(1) *If  $p(z) \in \mathcal{H}[1, n]$  and*

$$Q(z)[\alpha + (1 - \alpha)p(z)] \prec 1 + \lambda z \quad (z \in \mathbb{U})$$

*for some real  $\alpha$  such that*

$$\alpha = \begin{cases} \frac{1 - \lambda}{1 + \lambda_1} & (0 < \lambda + \lambda_1 \leq 1) \\ \frac{1 - (\lambda^2 + \lambda_1^2)}{2(1 - \lambda_1^2)} & (\lambda^2 + \lambda_1^2 \leq 1 \leq \lambda + \lambda_1), \end{cases} \quad (2.7)$$

*then  $\operatorname{Re}(p(z)) > 0$ .*

(2) *If  $w(z) \in \mathcal{H}[0, n]$  and*

$$Q(z)[1 + w(z)] \prec 1 + \lambda z \quad (z \in \mathbb{U}),$$

*then*

$$|w(z)| < \frac{\lambda + \lambda_1}{1 - \lambda_1} \leq 1, \quad (2.8)$$

*where  $\lambda + 2\lambda_1 \leq 1$ .*

*The bound (2.8) and the value of  $\alpha$  given by (2.7) are best possible.*

By virtue of Lemma 2, we deduce the sufficient condition for the class  $\mathcal{S}^*$ .

**Theorem 1.** *If  $f(z) \in \mathcal{A}_n$  satisfies the condition*

$$f'(z) \left( \frac{z}{f(z)} \right)^{1+\mu} \prec 1 + \lambda z \quad (z \in \mathbb{U})$$

*for some complex numbers  $\mu$  ( $\operatorname{Re}(\mu) < n$ ) and  $\lambda$  such that*

$$0 < |\lambda| \leq \frac{|n - \mu|}{\sqrt{|n - \mu|^2 + |\mu|^2}}, \text{ then } f(z) \in \mathcal{S}^*.$$

*Proof.* If we put

$$Q(z) = \left( \frac{z}{f(z)} \right)^\mu,$$

then we get

$$Q(z) - \frac{1}{\mu} Q'(z) = f'(z) \left( \frac{z}{f(z)} \right)^{1+\mu} \prec 1 + \lambda z \quad (z \in \mathbb{U})$$

for  $0 < |\lambda| \leq 1$ .

In view of Lemma 2, we obtain

$$Q(z) \prec 1 + \lambda_1 z \quad (z \in \mathbb{U}) \quad (2.9)$$

and

$$|\lambda_1| = |\lambda| \frac{|\mu|}{|n - \mu|}.$$

Then, we see that

$$\left| \arg \left( f'(z) \left( \frac{z}{f(z)} \right)^{1+\mu} \right) \right| < \arctan \left( \frac{|\lambda|}{\sqrt{1 - |\lambda|^2}} \right)$$

and

$$\left| \arg \left( \left( \frac{f(z)}{z} \right)^\mu \right) \right| = \left| \arg \left( \left( \frac{z}{f(z)} \right)^\mu \right) \right| < \arctan \left( \frac{|\lambda_1|}{\sqrt{1 - |\lambda_1|^2}} \right),$$

which give

$$\begin{aligned} \left| \arg \left( \frac{zf'(z)}{f(z)} \right) \right| &\leq \left| \arg \left( f'(z) \left( \frac{z}{f(z)} \right)^{1+\mu} \right) \right| + \left| \arg \left( \left( \frac{f(z)}{z} \right)^\mu \right) \right| \\ &< \arctan \left( \frac{|\lambda|}{\sqrt{1 - |\lambda|^2}} \right) + \arctan \left( \frac{|\lambda_1|}{\sqrt{1 - |\lambda_1|^2}} \right) \\ &= \arctan \left( \frac{\frac{|\lambda|}{\sqrt{1 - |\lambda|^2}} + \frac{|\lambda_1|}{\sqrt{1 - |\lambda_1|^2}}}{1 - \frac{|\lambda||\lambda_1|}{\sqrt{(1 - |\lambda|^2)(1 - |\lambda_1|^2)}}}} \right) \leq \frac{\pi}{2} \quad (z \in \mathbb{U}). \end{aligned}$$

This implies that  $f(z) \in \mathcal{S}^*$

On the other hand, when we have

$$1 - \frac{|\lambda||\lambda_1|}{\sqrt{(1 - |\lambda|^2)(1 - |\lambda_1|^2)}} \geq 0,$$

$\lambda$  satisfies

$$|\lambda| \leq \frac{|n - \mu|}{\sqrt{|n - \mu|^2 + |\mu|^2}} < 1.$$

□

Taking  $n = 1$ ,  $0 < \mu < 1$  and  $0 < \lambda < 1$ , we have the next corollary due to Obradović [4].

**Corollary 2.** *If  $f(z) \in \mathcal{A}$  satisfies the condition*

$$\left| f'(z) \left( \frac{z}{f(z)} \right)^{1+\mu} - 1 \right| < \lambda \quad (z \in \mathbb{U})$$

*for some  $0 < \mu < 1$  and  $0 < \lambda < 1$ , then  $f(z) \in \mathcal{S}^*$ .*

Applying Lemma 3, we derive the following theorem.

**Theorem 2.** Let  $f(z) \in \mathcal{A}_n$  satisfy the condition

$$f'(z) \left( \frac{z}{f(z)} \right)^{1+\mu} \prec 1 + \lambda z \quad (z \in \mathbb{U}) \quad (2.10)$$

for some complex number  $\mu$  ( $\operatorname{Re}(\mu) < \frac{n}{2}$ ). If the complex number  $\lambda_1$  is given by

$$|\lambda_1| = |\lambda| \frac{|\mu|}{|n - \mu|},$$

then

(1)  $f(z) \in \mathcal{S}^*(\alpha)$  where

$$\alpha \leq \begin{cases} \frac{1 - |\lambda|}{1 + |\lambda_1|} & \left( 0 < |\lambda| \leq \frac{|n - \mu|}{|n - \mu| + |\mu|} \right) \\ \frac{1 - (|\lambda|^2 + |\lambda_1|^2)}{2(1 - |\lambda_1|^2)} & \left( \frac{|n - \mu|}{|n - \mu| + |\mu|} \leq |\lambda| \leq \frac{|n - \mu|}{\sqrt{|n - \mu|^2 + |\mu|^2}} \right) \end{cases}$$

and

$$(2) \quad \left| \frac{zf'(z)}{f(z)} - 1 \right| < \frac{(|n - \mu| + |\mu|)|\lambda|}{|n - \mu| - |\mu||\lambda|} \leq 1 \quad (z \in \mathbb{U}),$$

$$\text{where } 0 < |\lambda| \leq \frac{|n - \mu|}{|n - \mu| + 2|\mu|}.$$

*Proof.* Let us define

$$Q(z) = \left( \frac{z}{f(z)} \right)^\mu,$$

$$p(z) = \frac{zf'(z)}{f(z)},$$

and

$$w(z) = \frac{zf'(z)}{f(z)} - 1.$$

Then by the equation (2.9) of Theorem 1, we have

$$Q(z) \prec 1 + \lambda_1 z \quad (z \in \mathbb{U})$$

where  $0 < |\lambda_1| = |\lambda| \frac{|\mu|}{|n - \mu|} < |\lambda| < 1$  since  $\operatorname{Re}(\mu) < \frac{n}{2}$ . Also, since the condition (2.10) is equivalent to

$$Q(z) \left[ \alpha + (1 - \alpha) \frac{p(z) - \alpha}{1 - \alpha} \right] \prec 1 + \lambda z \quad (z \in \mathbb{U})$$

with  $\alpha$  is given by (2.4) and as

$$Q(z)[1 + w(z)] \prec 1 + \lambda z \quad (z \in \mathbb{U}),$$

the statement of the theorem directly follows from Lemma 3.  $\square$

Putting  $n = 1$ ,  $0 < \mu < \frac{1}{2}$ ,  $0 < \lambda < 1$  and fixed  $\alpha$ , we get the following corollary due to Obradović [4].



**Corollary 3.** Let  $f(z) \in \mathcal{A}$  satisfy the condition

$$\left| f'(z) \left( \frac{z}{f(z)} \right)^{1+\mu} - 1 \right| < \lambda \quad (z \in \mathbb{U})$$

with some  $0 < \mu < \frac{1}{2}$ . If the real number  $\lambda_1$  is given by

$$\lambda_1 = \lambda \frac{\mu}{1-\mu},$$

then

(1)  $f(z) \in \mathcal{S}^*(\alpha)$ , where

$$\alpha = \begin{cases} \frac{1-\lambda}{1+\lambda_1} & (0 < \lambda \leq 1-\mu) \\ \frac{1-(\lambda^2+\lambda_1^2)}{2(1-\lambda_1^2)} & \left( 1-\mu \leq \lambda \leq \frac{1-\mu}{\sqrt{(1-\mu)^2+\mu^2}} \right). \end{cases}$$

(2)  $\left| \frac{zf'(z)}{f(z)} - 1 \right| < \frac{\lambda}{1-\mu-\mu\lambda} \leq 1 \quad (z \in \mathbb{U})$ ,

where  $0 < \lambda \leq \frac{1-\mu}{1+\mu}$ .

Using Lemma 2 and Lemma 3, we derive

**Theorem 3.** If  $f(z) \in \mathcal{A}_n$  satisfies

$$f'(z) \left( \frac{z}{f(z)} \right)^{1+\mu} \prec 1 + \lambda z \quad (z \in \mathbb{U})$$

and

$$F(z) = z \left[ \frac{c-\mu}{z^{c-\mu}} \int_0^z \left( \frac{t}{f(t)} \right)^\mu t^{c-\mu-1} dt \right]^{-\frac{1}{\mu}} \quad (z \in \mathbb{U})$$

for some complex numbers  $\lambda, \mu$ , and  $c$  such that  $\operatorname{Re}(c-\mu) < n$ , then

(1)  $F(z) \in \mathcal{S}^*$  for  $|c-\mu||\lambda| \leq \frac{|n-\mu||n-(c-\mu)|}{\sqrt{|n-\mu|^2+|\mu|^2}}$ .

(2)  $F(z) \in \mathcal{S}^*(\alpha)$  where

$$\alpha \leq \begin{cases} \frac{1-|\lambda_1|}{1+|\lambda_2|} & \left( 0 < |\lambda_1| \leq \frac{|n-\mu|}{|n-\mu|+|\mu|} \right) \\ \frac{1-(|\lambda_1|^2+|\lambda_2|^2)}{2(1-|\lambda_2|^2)} & \left( \frac{|n-\mu|}{|n-\mu|+|\mu|} \leq |\lambda_1| \leq \frac{|n-\mu|}{\sqrt{|n-\mu|^2+|\mu|^2}} \right), \end{cases}$$

$$|\lambda_1| = |\lambda| \frac{|c-\mu|}{|n-(c-\mu)|}, \quad |\lambda_2| = |\lambda_1| \frac{|\mu|}{|n-\mu|} \quad \text{and} \quad \operatorname{Re}(\mu) < \frac{n}{2}.$$

(3)  $\left| \frac{zF'(z)}{F(z)} - 1 \right| < \frac{(|n-\mu|+|\mu|)|c-\mu||\lambda|}{|n-(c-\mu)||n-\mu|-|c-\mu||\mu||\lambda|} \leq 1 \quad (z \in \mathbb{U})$

where  $|c-\mu||\lambda| \leq \frac{|n-(c-\mu)||n-\mu|}{|n-\mu|+2|\mu|}$  and  $\operatorname{Re}(\mu) < \frac{n}{2}$ .

*Proof.* If we put

$$Q(z) = F'(z) \left( \frac{z}{F(z)} \right)^{1+\mu},$$

after some transformations, we obtain

$$Q(z) + \frac{1}{c-\mu} z Q'(z) = f'(z) \left( \frac{z}{f(z)} \right)^{1+\mu} \prec 1 + \lambda z.$$

Therefore, spending the same technique as in the proof of Lemma 2, we have that

$$Q(z) \prec 1 + \lambda_2 z \quad (z \in \mathbb{U})$$

and

$$|\lambda_2| = \frac{|\mu| |c-\mu|}{|n-\mu| |n-(c-\mu)|}.$$

The statement of the theorem now easily follows from Theorem 1 and Theorem 2.  $\square$

Taking  $n = 1$ ,  $0 < \mu < 1$ ,  $0 < \lambda < 1$  and fixed  $\alpha$ , we obtain the next the corollary due to Obradović [4].

**Corollary 4.** *If  $f(z) \in \mathcal{A}$  satisfies*

$$\left| f'(z) \left( \frac{z}{f(z)} \right)^{1+\mu} - 1 \right| < \lambda \quad (z \in \mathbb{U})$$

and

$$F(z) = z \left[ \frac{c-\mu}{z^{c-\mu}} \int_0^z \left( \frac{t}{f(t)} \right)^\mu t^{c-\mu-1} dt \right]^{-\frac{1}{\mu}} \quad (z \in \mathbb{U})$$

for  $c-\mu > 0$ , then

$$(1) \quad F(z) \in \mathcal{S}^* \text{ for } (c-\mu)\lambda \leq \frac{(1-\mu)(1-(c-\mu))}{\sqrt{(1-\mu)^2 + \mu^2}}.$$

$$(2) \quad F(z) \in \mathcal{S}^*(\alpha) \text{ where}$$

$$\alpha = \begin{cases} \frac{1-\lambda_1}{1+\lambda_2} & (0 < \lambda_1 \leq 1-\mu) \\ \frac{1-(\lambda_1^2 + \lambda_2^2)}{2(1-\lambda_2^2)} & \left( 1-\mu \leq \lambda_1 \leq \frac{1-\mu}{\sqrt{(1-\mu)^2 + \mu^2}} \right), \end{cases}$$

$$\lambda_1 = \lambda \frac{c-\mu}{1-(c-\mu)}, \quad \lambda_2 = \lambda_1 \frac{\mu}{1-\mu} \text{ and } 0 < \mu < \frac{1}{2}.$$

$$(3) \quad \left| \frac{zF'(z)}{F(z)} - 1 \right| < \frac{(c-\mu)\lambda}{(1-(c-\mu))(1-\mu) - (c-\mu)\mu\lambda} \leq 1 \quad (z \in \mathbb{U}) \text{ where}$$

$$0 < \lambda \leq \frac{(1-(c-\mu))(1-\mu)}{(c-\mu)(1+\mu)} \text{ and } 0 < \mu < \frac{1}{2}.$$

## REFERENCES

- [1] P. L. Duren, *Univalent Functions*, Springer-verlag, New York, Berlin, Heiderberg, Tokyo (1983).
- [2] S. S. Miller and P. T. Mocanu, *Second-order differential inequalities in the complex plane*, J. Math. Anal. Appl. **65**(1978), 289–305.
- [3] S. S. Miller and P. T. Mocanu, *Differential Subordinations, Theory and Applications*. Monographs and Textbooks in Pure and Applied Mathematics, 225. Marcel Dekker, Inc., New York, 2000.
- [4] M. Obradović, *A class of univalent functions*, Hokkaido Math. J. **27**(1998), 329–335.
- [5] S. Ponnusamy and V. Singh, *Convolution properties of some classes of analytic functions*, J. Mathe. Sci. **89**(1998), 1008–1020.

HITOSHI SHIRAISHI

DEPARTMENT OF MATHEMATICS

KINKI UNIVERSITY

HIGASHI-OSAKA, OSAKA 577-8502, JAPAN

*E-mail address:* shiraishi@math.kindai.ac.jp

SHIGEYOSHI OWA

DEPARTMENT OF MATHEMATICS

KINKI UNIVERSITY

HIGASHI-OSAKA, OSAKA 577-8502, JAPAN

*E-mail address:* owa@math.kindai.ac.jp

TOSHIO HAYAMI

SCHOOL OF SCIENCE AND TECHNOLOGY

KINKI UNIVERSITY

SANDA, HYOGO 669-1337, JAPAN

*E-mail address:* ha\_ya\_to112@hotmail.com

KAZUO KUROKI

DEPARTMENT OF MATHEMATICS

KINKI UNIVERSITY

HIGASHI-OSAKA, OSAKA 577-8502, JAPAN

*E-mail address:* freedom@sakai.zaq.ne.jp

H. M. SRIVASTAVA

DEPARTMENT OF MATHEMATICS AND STATISTICS

UNIVERSITY OF VICTORIA

VICTORIA, BRITISH COLUMBIA V8W 3P4, CANADA

*E-mail address:* harimsri@math.uvic.ca